



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 10, 2021 – 11:03 am BST

PDB ID : 7A0U
Title : LIGAND FREE TYPE II E. COLI ASPARAGINASE T12S/T89S DOUBLE
MUTANT
Authors : Maggi, M.; Scotti, C.
Deposited on : 2020-08-10
Resolution : 2.26 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : 1.13
EDS : 2.23.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.1


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.26 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1377 (2.26-2.26)
Clashscore	141614	1487 (2.26-2.26)
Ramachandran outliers	138981	1449 (2.26-2.26)
Sidechain outliers	138945	1450 (2.26-2.26)
RSRZ outliers	127900	1356 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	333	
1	B	333	
1	C	333	
1	D	333	

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 19312 atoms, of which 9265 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called L-asparaginase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	316	4705	1471	2346	405	475	8	0	0	0
1	B	326	4814	1513	2389	415	489	8	0	1	0
1	C	308	4572	1439	2269	395	461	8	0	0	0
1	D	305	4541	1426	2261	389	457	8	0	0	0

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-6	MET	-	initiating methionine	UNP P00805
A	-5	HIS	-	expression tag	UNP P00805
A	-4	HIS	-	expression tag	UNP P00805
A	-3	HIS	-	expression tag	UNP P00805
A	-2	HIS	-	expression tag	UNP P00805
A	-1	HIS	-	expression tag	UNP P00805
A	0	HIS	-	expression tag	UNP P00805
A	12	SER	THR	engineered mutation	UNP P00805
A	89	SER	THR	engineered mutation	UNP P00805
B	-6	MET	-	initiating methionine	UNP P00805
B	-5	HIS	-	expression tag	UNP P00805
B	-4	HIS	-	expression tag	UNP P00805
B	-3	HIS	-	expression tag	UNP P00805
B	-2	HIS	-	expression tag	UNP P00805
B	-1	HIS	-	expression tag	UNP P00805
B	0	HIS	-	expression tag	UNP P00805
B	12	SER	THR	engineered mutation	UNP P00805
B	89	SER	THR	engineered mutation	UNP P00805
C	-6	MET	-	initiating methionine	UNP P00805
C	-5	HIS	-	expression tag	UNP P00805
C	-4	HIS	-	expression tag	UNP P00805

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-3	HIS	-	expression tag	UNP P00805
C	-2	HIS	-	expression tag	UNP P00805
C	-1	HIS	-	expression tag	UNP P00805
C	0	HIS	-	expression tag	UNP P00805
C	12	SER	THR	engineered mutation	UNP P00805
C	89	SER	THR	engineered mutation	UNP P00805
D	-6	MET	-	initiating methionine	UNP P00805
D	-5	HIS	-	expression tag	UNP P00805
D	-4	HIS	-	expression tag	UNP P00805
D	-3	HIS	-	expression tag	UNP P00805
D	-2	HIS	-	expression tag	UNP P00805
D	-1	HIS	-	expression tag	UNP P00805
D	0	HIS	-	expression tag	UNP P00805
D	12	SER	THR	engineered mutation	UNP P00805
D	89	SER	THR	engineered mutation	UNP P00805


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	191	Total O 191 191	0	0
2	B	176	Total O 176 176	0	0
2	C	180	Total O 180 180	0	0
2	D	133	Total O 133 133	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: L-asparaginase 2




- Molecule 1: L-asparaginase 2



- Molecule 1: L-asparaginase 2



- Molecule 1: L-asparaginase 2



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	150.85Å 61.86Å 142.42Å 90.00° 117.94° 90.00°	Depositor
Resolution (Å)	125.83 – 2.26 125.83 – 2.26	Depositor EDS
% Data completeness (in resolution range)	96.9 (125.83-2.26) 96.9 (125.83-2.26)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.92 (at 2.27Å)	Xtrriage
Refinement program	PHENIX 1.11.1_2575, PHENIX 1.11.1_2575	Depositor
R, R_{free}	0.171 , 0.240 0.171 , 0.240	Depositor DCC
R_{free} test set	2659 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	26.2	Xtrriage
Anisotropy	0.158	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 40.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	19312	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.70% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.62	0/2395	0.79	4/3259 (0.1%)
1	B	0.58	0/2469	0.69	0/3362
1	C	0.62	0/2340	0.71	1/3186 (0.0%)
1	D	0.55	0/2315	0.69	2/3153 (0.1%)
All	All	0.59	0/9519	0.72	7/12960 (0.1%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	272	ARG	NE-CZ-NH2	-12.44	114.08	120.30
1	D	272	ARG	NE-CZ-NH2	-8.35	116.12	120.30
1	A	225	ASP	CB-CG-OD1	7.43	124.99	118.30
1	D	272	ARG	NE-CZ-NH1	7.40	124.00	120.30
1	A	272	ARG	NE-CZ-NH1	6.23	123.42	120.30
1	A	272	ARG	CG-CD-NE	5.28	122.88	111.80
1	C	269	ARG	NE-CZ-NH1	5.24	122.92	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2359	2346	2347	14	0
1	B	2425	2389	2395	14	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	2303	2269	2279	16	0
1	D	2280	2261	2269	14	0
2	A	191	0	0	1	0
2	B	176	0	0	1	0
2	C	180	0	0	0	1
2	D	133	0	0	0	0
All	All	10047	9265	9290	54	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:37:ASN:OD1	1:D:38:ALA:N	2.30	0.64
1:C:163:THR:HB	1:C:170:THR:O	1.98	0.63
1:C:92:MET:CE	1:C:148:VAL:HG13	2.29	0.63
1:B:112:VAL:HG13	1:B:132:ALA:HB2	1.83	0.60
1:D:73:ILE:HD11	1:D:85:ILE:HD11	1.84	0.58
1:B:40:PRO:O	1:B:43:LYS:HG2	2.05	0.57
1:A:40:PRO:O	1:A:43:LYS:HG2	2.05	0.57
1:D:56:ILE:HD11	1:D:61:MET:HE2	1.88	0.56
1:C:182:ILE:HG12	1:C:187:ILE:HG12	1.88	0.56
1:C:317:GLN:O	1:C:321:GLN:HG2	2.09	0.53
1:A:227:PRO:HB3	1:C:227:PRO:HB3	1.91	0.52
1:B:112:VAL:CG1	1:B:132:ALA:HB2	2.38	0.52
1:A:164:ASN:HA	1:C:275:THR:O	2.10	0.50
1:A:303:ARG:NH2	1:C:222:ASN:HB3	2.29	0.48
1:B:21:THR:HG22	1:B:21:THR:O	2.14	0.47
1:D:261:ALA:HA	1:D:265:THR:O	2.14	0.47
1:B:19:SER:O	1:B:20:ALA:HB3	2.14	0.47
1:A:96:ALA:O	1:A:100:ASP:HB2	2.16	0.46
1:B:84:VAL:HA	1:B:110:VAL:O	2.16	0.45
1:D:5:THR:CG2	1:D:80:THR:HG21	2.46	0.45
1:D:84:VAL:HA	1:D:110:VAL:O	2.16	0.45
1:A:0:HIS:CG	1:A:0:HIS:O	2.70	0.45
1:C:92:MET:HE2	1:C:148:VAL:HG13	1.97	0.45
1:A:51:GLU:OE1	1:A:72:LYS:NZ	2.44	0.45
1:A:261:ALA:HA	1:A:265:THR:O	2.17	0.44
1:A:138:ASP:OD2	2:A:401:HOH:O	2.21	0.44
1:D:144:ARG:NH1	1:D:191:ARG:O	2.51	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:260:ALA:HB1	1:D:265:THR:HB	1.98	0.44
1:A:251:LYS:HG3	1:A:252:SER:N	2.32	0.43
1:C:6:ILE:HA	1:C:84:VAL:O	2.17	0.43
1:D:53:VAL:HG23	1:D:54:VAL:HG22	1.99	0.43
1:C:156:ASP:HB2	1:C:193:PRO:HG3	2.01	0.43
1:B:150:MET:O	1:B:151:ASN:C	2.57	0.43
1:D:298:ASN:HB2	1:D:299:PRO:CD	2.48	0.43
1:D:179:LEU:HA	1:D:191:ARG:HB2	2.00	0.43
1:D:6:ILE:HA	1:D:84:VAL:O	2.20	0.42
1:D:124:ASP:OD1	1:D:124:ASP:N	2.43	0.42
1:B:238:GLY:HA2	1:B:266:ALA:O	2.19	0.42
1:C:51:GLU:CD	1:C:72:LYS:HZ1	2.22	0.42
1:B:316:PRO:HD2	2:B:401:HOH:O	2.19	0.42
1:D:72:LYS:HE2	1:D:76:ASP:OD2	2.19	0.42
1:A:103:VAL:O	1:A:198:THR:HA	2.21	0.41
1:C:92:MET:HE3	1:C:148:VAL:HG13	1.99	0.41
1:C:214:VAL:HA	1:C:238:GLY:O	2.20	0.41
1:B:6:ILE:HA	1:B:84:VAL:O	2.20	0.41
1:C:2:PRO:HG3	1:C:137:ALA:HB1	2.01	0.41
1:A:274:PRO:HB3	1:C:274:PRO:HB3	2.03	0.41
1:C:58:SER:HB3	1:C:88:GLY:HA3	2.02	0.41
1:A:263:THR:HG22	1:A:263:THR:O	2.20	0.40
1:B:73:ILE:HD11	1:B:85:ILE:HD11	2.02	0.40
1:B:156:ASP:HB2	1:B:193:PRO:HG3	2.02	0.40
1:B:214:VAL:HA	1:B:238:GLY:O	2.22	0.40
1:A:84:VAL:HA	1:A:110:VAL:O	2.21	0.40
1:B:243:GLY:O	1:B:272:ARG:HG2	2.20	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:456:HOH:O	2:C:566:HOH:O[2_555]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	312/333 (94%)	306 (98%)	5 (2%)	1 (0%)	41	46
1	B	323/333 (97%)	310 (96%)	13 (4%)	0	100	100
1	C	304/333 (91%)	296 (97%)	8 (3%)	0	100	100
1	D	301/333 (90%)	291 (97%)	10 (3%)	0	100	100
All	All	1240/1332 (93%)	1203 (97%)	36 (3%)	1 (0%)	51	60

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	198	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	258/273 (94%)	249 (96%)	9 (4%)	36	43
1	B	265/273 (97%)	257 (97%)	8 (3%)	41	50
1	C	250/273 (92%)	240 (96%)	10 (4%)	31	37
1	D	249/273 (91%)	242 (97%)	7 (3%)	43	52
All	All	1022/1092 (94%)	988 (97%)	34 (3%)	38	46

All (34) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	19	SER
1	A	49	LYS
1	A	54	VAL
1	A	112	VAL
1	A	156	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	201	THR
1	A	222	ASN
1	A	254	PHE
1	A	317	GLN
1	B	0	HIS
1	B	22	LYS
1	B	29	LYS
1	B	112	VAL
1	B	120	SER
1	B	156	ASP
1	B	254	PHE
1	B	263	THR
1	C	12	SER
1	C	139	LYS
1	C	156	ASP
1	C	163	THR
1	C	222	ASN
1	C	226	LEU
1	C	254	PHE
1	C	272	ARG
1	C	283	GLU
1	C	314	LYS
1	D	5	THR
1	D	207	LYS
1	D	222	ASN
1	D	254	PHE
1	D	272	ARG
1	D	314	LYS
1	D	321	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	316/333 (94%)	-0.58	1 (0%) 94 94	17, 25, 43, 84	0
1	B	326/333 (97%)	-0.48	3 (0%) 84 85	19, 29, 52, 86	0
1	C	308/333 (92%)	-0.55	2 (0%) 89 89	17, 26, 44, 74	0
1	D	305/333 (91%)	-0.50	1 (0%) 94 94	22, 33, 52, 66	0
All	All	1255/1332 (94%)	-0.52	7 (0%) 89 89	17, 28, 50, 86	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	17	GLY	3.0
1	B	19	SER	2.9
1	B	0	HIS	2.9
1	C	-1	HIS	2.7
1	A	0	HIS	2.6
1	D	38	ALA	2.2
1	C	78	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.