



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 15, 2020 – 11:52 pm BST

PDB ID : 4A3I  
Title : RNA Polymerase II binary complex with DNA  
Authors : Cheung, A.C.M.; Sainsbury, S.; Cramer, P.  
Deposited on : 2011-09-30  
Resolution : 3.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1212 (4.00-3.60)
Clashscore	141614	1288 (4.00-3.60)
Ramachandran outliers	138981	1243 (4.00-3.60)
Sidechain outliers	138945	1237 (4.00-3.60)
RSRZ outliers	127900	1121 (4.00-3.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1732	
2	B	1224	
3	C	318	
4	D	221	
5	E	215	
6	F	155	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
7	G	171	<p>58% 37%</p>
8	H	146	<p>3% 59% 25% 6% 9%</p>
9	I	122	<p>2% 70% 22% 6%</p>
10	J	70	<p>44% 36% 13% 7%</p>
11	K	120	<p>61% 28% 7%</p>
12	L	70	<p>3% 31% 24% 10% 34%</p>
13	N	15	<p>40% 33% 27%</p>
14	T	27	<p>48% 11% 37%</p>

## 2 Entry composition [i](#)

There are 16 unique types of molecules in this entry. The entry contains 31768 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-DIRECTED RNA POLYMERASE II SUBUNIT RPB1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	1422	11174	7037	1954	2121	62	0	0	0

- Molecule 2 is a protein called DNA-DIRECTED RNA POLYMERASE II SUBUNIT RPB2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	1115	8859	5609	1554	1641	55	0	0	0

- Molecule 3 is a protein called DNA-DIRECTED RNA POLYMERASE II SUBUNIT RPB3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	266	2095	1317	348	417	13	0	0	0

- Molecule 4 is a protein called DNA-DIRECTED RNA POLYMERASE II SUBUNIT RPB4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	D	178	1434	887	257	288	2	0	0	0

- Molecule 5 is a protein called DNA-DIRECTED RNA POLYMERASES I, II, AND III SUBUNIT RPABC 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
5	E	214	1752	1111	309	321	11	0	0	0

- Molecule 6 is a protein called DNA-DIRECTED RNA POLYMERASES I, II, AND III SUBUNIT RPABC 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
6	F	84	679	434	115	127	3	0	0	0

- Molecule 7 is a protein called RPB7, DNA-DIRECTED RNA POLYMERASE II SUBUNIT RPB7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
7	G	171	1340	861	222	249	8	0	0	0

- Molecule 8 is a protein called DNA-DIRECTED RNA POLYMERASES I, II, AND III SUBUNIT RPABC 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
8	H	133	1068	673	180	211	4	0	0	0

- Molecule 9 is a protein called DNA-DIRECTED RNA POLYMERASE II SUBUNIT RPB9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
9	I	119	971	596	179	186	10	0	0	0

- Molecule 10 is a protein called DNA-DIRECTED RNA POLYMERASES I, II, AND III SUBUNIT RPABC 5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
10	J	65	532	339	93	94	6	0	0	0

- Molecule 11 is a protein called DNA-DIRECTED RNA POLYMERASE II SUBUNIT RPB11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
11	K	115	920	590	157	171	2	0	0	1

- Molecule 12 is a protein called DNA-DIRECTED RNA POLYMERASES I, II, AND III SUBUNIT RPABC 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
12	L	46	363	224	72	63	4	0	0	0

- Molecule 13 is a DNA chain called 5'-D(\*GP\*GP\*CP\*AP\*CP\*AP\*AP\*CP\*TP\*GP\*CP\*GP\*GP\*CP\*T)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
13	N	11	222	106	44	62	10	0	0	0

- Molecule 14 is a DNA chain called TEMPLATE DNA 27-MER.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	Br	C	N	O				P
14	T	17	350	1	166	61	105	17	0	0	0

- Molecule 15 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
15	J	1	Total	Zn	0	0
			1	1		
15	B	1	Total	Zn	0	0
			1	1		
15	I	2	Total	Zn	0	0
			2	2		
15	C	1	Total	Zn	0	0
			1	1		
15	A	2	Total	Zn	0	0
			2	2		
15	L	1	Total	Zn	0	0
			1	1		


- Molecule 16 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
16	A	1	Total	Mg	0	0
			1	1		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA-DIRECTED RNA POLYMERASE II SUBUNIT RPB1



T1095	L1092	G1286	R1391	VAL	PRO	ALA	LYS
R1100	L1195	E1297	S1392	GLY	THR	TYR	GLN
L1101	L1199	E1301	M1393	PHE	SER	SER	ASP
I1104	R1199	E1302	C1400	LEU	PRO	PRO	GLU
M1111	T1208	L1306	S1401	THR	TYR	TYR	GLN
K1112	T1208	E1307	F1402	VAL	SER	PRO	HIS
T1113	G1213	T1308	T1405	LYS	PRO	PRO	ASN
P1114	G1213	D1309	T1405	GLU	THR	TYR	GLU
S1115	E1214	T1308	A1416	GLY	SER	SER	ASN
L1116	R1215	G1310	E1417	GLU	PRO	PRO	GLU
T1117	R1215	E1310	F1417	MET	PRO	PRO	ASN
V1118	L1216	V1311	L1418	PHE	THR	TYR	SER
P1122	K1217	E1315	R1422	SER	SER	SER	PRO
G1123	V1118	V1316	E1426	PRO	PRO	PRO	THR
H1124	D1223	M1317	E1426	ALA	THR	THR	TYR
Q1130	E1234	T1325	L1430	VAL	SER	SER	SER
L1133	E1237	R1326	G1431	SER	PRO	PRO	THR
I1134	I1237	I1328	Q1432	GLY	THR	THR	SER
L1133	I1238	Y1328	I1436	ASN	PRO	PRO	PRO
I1134	R1239	T1329	G1437	SER	THR	THR	THR
T1142	C1240	N1330	F1441	MET	PRO	PRO	THR
V1146	R1241	S1331	D1442	ALA	SER	SER	THR
T1147	V1242	D1334	V1443	THR	THR	THR	PRO
I1148	V1243	I1335	V1443	GLY	SER	SER	PRO
D1155	ARG	M1336	M1444	PHE	PRO	PRO	SER
P1158	LYS	M1336	I1445	THR	THR	THR	THR
R1159	SER	L1339	I1445	THR	SER	SER	THR
D1165	LEU	G1340	V1451	ALA	PRO	PRO	PRO
E1167	ASP	I1341	R1452	THR	SER	SER	PRO
E1168	ALA	E1342	M1454	GLY	PRO	PRO	PRO
I1169	GLU	Y1349	P1455	ALA	SER	PRO	PRO
I1170	THR	K1350	GLJ	THR	PRO	PRO	PRO
O1171	GLU	E1351	GLN	TYR	SER	SER	PRO
L1172	A1264	E1254	LYS	GLY	THR	THR	PRO
H1173	E1255	D1166	ILE	GLU	PRO	PRO	PRO
F1174	E1256	E1167	ALA	ALA	PRO	PRO	PRO
S1175	D1257	E1168	THR	TYR	TYR	TYR	PRO
L1176	H1258	I1169	THR	THR	THR	THR	PRO
LEU	M1259	O1171	ILE	PRO	PRO	PRO	PRO
ASP	K1262	L1172	ASP	PHE	THR	THR	PRO
GLU	K1262	L1172	ASP	GLY	THR	THR	PRO
GLU	H1173	H1173	GLY	ALA	PRO	PRO	PRO
GLU	F1174	F1174	GLN	TYR	PRO	PRO	PRO
ALA	S1175	S1175	GLY	GLY	THR	THR	PRO
GLU	M1265	T1266	ASP	THR	THR	THR	PRO
GLU	M1267	M1267	GLY	ALA	PRO	PRO	PRO
SER	I1271	I1271	VAL	PRO	THR	THR	PRO
SER	I1279	I1279	THR	THR	THR	THR	PRO
PHE	E1280	E1280	TYR	PRO	PRO	PRO	PRO
SER	GLU	GLU	S1383	PRO	PRO	PRO	PRO
GLN	D1288	D1288	SER	THR	THR	THR	PRO
SER	H1387	H1387	ASN	SER	SER	SER	PRO
ASP	F1388	F1388	GLU	GLY	PRO	PRO	PRO
ASP	P1294	P1294	G1389	VAL	THR	THR	PRO
LEU	T1295	T1295	M1390	SER	PRO	PRO	PRO


● Molecule 2: DNA-DIRECTED RNA POLYMERASE II SUBUNIT RPB2




MET	L1170	L284	F370	R485
SER	TYR	L285	R373	V486
ASP	GLU	I90	L385	S490
LEU	S91	F92	L386	L502
ALA	F92	Y96	D894	GLY
ASN	Y96	M101	Q395	ARG
SER	M101	V102	D397	ASP
GLU	V102	V108	R398	GLY
ASN	V108	L112	Y303	LYS
HIS	L112	Y113	Q309	LEU
ASN	Y113	P114	L314	K510
SER	P114	R118	G317	L514
PRO	R118	M121	V316	T517
THR	M121	L122	F322	E526
PRO	L122	S126	V323	K418
THR	S126	L128	I324	K423
PRO	L128	F129	R327	L424
PRO	F129	V130	D332	D427
THR	V130	K134	R336	I428
THR	K134	ARG	R337	R434
PRO	ARG	THR	G338	E437
THR	THR	THR	I240	GLU
PRO	GLU	GLU	A340	ALA
THR	ILE	ILE	L341	HIS
PRO	ALA	ALA	G342	ASP
ASN	VAL	VAL	Y343	PHE
THR	PRO	PRO	K344	ASN
SER	GLY	GLY	R345	MET
PRO	ARG	ARG	F346	LYS
PRO	GLU	GLU	R347	LYS
THR	THR	THR	R261	L446
PRO	LEU	LEU	E262	A447
PRO	L69	L70	I349	I448
LEU	I70	LEU	Q350	T452
GLU	LEU	GLU	I351	L453
GLN	GLN	GLN	A352	S574
ALA	ALA	ALA	I355	A577
ALA	ALA	ALA	T267	R579
HIS	HIS	HIS	I269	V580
THR	THR	THR	T272	F581
THR	THR	THR	L273	K470
GLU	GLU	GLU	P274	K471
GLU	ASP	ASP	Y275	K472
ASP	ASP	ASP	I276	L472
ASN	ASN	ASN	K277	S475
ILE	ILE	ILE	Q278	R476
ARG	ARG	ARG	I282	V589
LYS	LYS	LYS	V283	









- Molecule 6: DNA-DIRECTED RNA POLYMERASES I, II, AND III SUBUNIT RPABC 2



- Molecule 7: RPB7, DNA-DIRECTED RNA POLYMERASE II SUBUNIT RPB7




- Molecule 8: DNA-DIRECTED RNA POLYMERASES I, II, AND III SUBUNIT RPABC 3



- Molecule 9: DNA-DIRECTED RNA POLYMERASE II SUBUNIT RPB9






- Molecule 10: DNA-DIRECTED RNA POLYMERASES I, II, AND III SUBUNIT RPABC 5

Chain J:




- Molecule 11: DNA-DIRECTED RNA POLYMERASE II SUBUNIT RPB11

Chain K:



- Molecule 12: DNA-DIRECTED RNA POLYMERASES I, II, AND III SUBUNIT RPABC 4

Chain L:




- Molecule 13: 5'-D(\*GP\*GP\*CP\*AP\*CP\*AP\*AP\*CP\*TP\*GP\*CP\*GP\*GP\*CP\*T)-3'

Chain N:



- Molecule 14: TEMPLATE DNA 27-MER

Chain T:



## 4 Data and refinement statistics i

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	221.14Å 393.18Å 282.48Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.80 – 3.80 49.80 – 3.80	Depositor EDS
% Data completeness (in resolution range)	100.0 (49.80-3.80) 100.0 (49.80-3.80)	Depositor EDS
$R_{merge}$	0.89	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.92 (at 3.77Å)	Xtriage
Refinement program	BUSTER 2.11.2	Depositor
R, $R_{free}$	0.159 , 0.191 0.184 , 0.210	Depositor DCC
$R_{free}$ test set	2395 reflections (1.99%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	111.3	Xtriage
Anisotropy	0.409	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 112.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	0.026 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.034 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	31768	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	131.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.83% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ZN, BRU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.54	0/11374	0.86	10/15383 (0.1%)
2	B	0.52	0/9029	0.81	4/12171 (0.0%)
3	C	0.49	0/2133	0.80	0/2891
4	D	0.53	0/1444	0.83	0/1935
5	E	0.48	0/1788	0.76	0/2406
6	F	0.60	0/691	0.82	0/933
7	G	0.52	0/1368	0.82	0/1844
8	H	0.51	0/1086	0.83	0/1470
9	I	0.45	0/989	0.77	0/1331
10	J	0.57	0/541	0.90	0/727
11	K	0.49	0/938	0.77	0/1267
12	L	0.57	0/365	1.00	0/485
13	N	1.12	0/249	0.97	0/382
14	T	1.20	1/369 (0.3%)	0.96	0/568
All	All	0.54	1/32364 (0.0%)	0.83	14/43793 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
14	T	19	DT	C1'-N1	5.82	1.56	1.49

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	399	HIS	N-CA-CB	7.68	124.42	110.60
1	A	56	PRO	C-N-CA	7.07	139.38	121.70
2	B	339	THR	C-N-CA	6.22	137.25	121.70
1	A	34	LYS	C-N-CA	6.11	136.96	121.70
1	A	194	ALA	C-N-CA	5.80	136.19	121.70

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	34	LYS	Mainchain,Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	11174	0	11233	252	0
2	B	8859	0	8901	166	0
3	C	2095	0	2051	59	0
4	D	1434	0	1460	36	0
5	E	1752	0	1776	37	0
6	F	679	0	701	22	0
7	G	1340	0	1357	45	0
8	H	1068	0	1040	27	0
9	I	971	0	927	17	0
10	J	532	0	542	23	0
11	K	920	0	929	28	0
12	L	363	0	386	7	0
13	N	222	0	124	4	0
14	T	350	0	191	3	0
15	A	2	0	0	0	0
15	B	1	0	0	0	0
15	C	1	0	0	0	0
15	I	2	0	0	0	0
15	J	1	0	0	0	0
15	L	1	0	0	0	0
16	A	1	0	0	0	0
All	All	31768	0	31618	640	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 640 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:867:ILE:CD1	1:A:867:ILE:CG1	1.81	1.53
1:A:37:PHE:HD1	1:A:52:GLY:HA3	1.26	1.01
1:A:768:GLN:HG2	1:A:816:HIS:HA	1.49	0.94
1:A:35:ILE:HG22	1:A:84:ILE:HG22	1.51	0.92
2:B:114:PRO:HG3	2:B:181:LEU:HD11	1.52	0.90

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1414/1732 (82%)	1207 (85%)	137 (10%)	70 (5%)	2	23
2	B	1095/1224 (90%)	929 (85%)	122 (11%)	44 (4%)	3	28
3	C	264/318 (83%)	231 (88%)	25 (10%)	8 (3%)	4	33
4	D	174/221 (79%)	152 (87%)	13 (8%)	9 (5%)	2	23
5	E	212/215 (99%)	186 (88%)	19 (9%)	7 (3%)	4	32
6	F	82/155 (53%)	75 (92%)	6 (7%)	1 (1%)	13	50
7	G	169/171 (99%)	149 (88%)	17 (10%)	3 (2%)	8	42
8	H	129/146 (88%)	102 (79%)	16 (12%)	11 (8%)	1	12
9	I	117/122 (96%)	91 (78%)	23 (20%)	3 (3%)	5	36
10	J	63/70 (90%)	52 (82%)	6 (10%)	5 (8%)	1	14
11	K	113/120 (94%)	107 (95%)	6 (5%)	0	100	100
12	L	44/70 (63%)	26 (59%)	10 (23%)	8 (18%)	0	2
All	All	3876/4564 (85%)	3307 (85%)	400 (10%)	169 (4%)	2	25

5 of 169 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	43	GLU
1	A	48	ALA
1	A	54	ASN
1	A	57	ARG
1	A	58	LEU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1240/1519 (82%)	1029 (83%)	211 (17%)	2	14
2	B	966/1061 (91%)	813 (84%)	153 (16%)	2	17
3	C	234/274 (85%)	200 (86%)	34 (14%)	3	19
4	D	160/200 (80%)	132 (82%)	28 (18%)	2	13
5	E	196/197 (100%)	176 (90%)	20 (10%)	7	31
6	F	74/137 (54%)	64 (86%)	10 (14%)	4	22
7	G	152/152 (100%)	131 (86%)	21 (14%)	3	22
8	H	117/128 (91%)	101 (86%)	16 (14%)	3	22
9	I	113/116 (97%)	101 (89%)	12 (11%)	6	30
10	J	60/65 (92%)	45 (75%)	15 (25%)	0	4
11	K	99/102 (97%)	82 (83%)	17 (17%)	2	13
12	L	40/57 (70%)	28 (70%)	12 (30%)	0	2
All	All	3451/4008 (86%)	2902 (84%)	549 (16%)	2	16

5 of 549 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	343	ILE
2	B	835	GLN
10	J	14	VAL
2	B	373	ARG

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type
2	B	498	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 41 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	255	GLN
2	B	449	ASN
8	H	139	ASN
2	B	300	HIS
2	B	350	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
14	BRU	T	22	14	15,21,22	2.68	3 (20%)	17,30,33	2.83	5 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
14	BRU	T	22	14	-	2/4/21/22	0/2/2/2

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
14	T	22	BRU	C4-C5	9.12	1.50	1.38
14	T	22	BRU	C4-N3	3.93	1.39	1.33
14	T	22	BRU	BR-C5	2.05	1.96	1.90

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
14	T	22	BRU	C4-N3-C2	9.35	123.04	115.14
14	T	22	BRU	C5-C4-N3	-4.71	118.00	123.64
14	T	22	BRU	BR-C5-C4	3.07	126.17	121.50
14	T	22	BRU	C5-C6-N1	3.00	123.85	119.97
14	T	22	BRU	C2'-C1'-N1	2.71	120.52	114.27

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
14	T	22	BRU	C3'-C4'-C5'-O5'
14	T	22	BRU	O4'-C4'-C5'-O5'

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
14	T	22	BRU	1	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 9 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
2	B	2

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	B	934:LYS	C	935:ARG	N	5.74
1	B	351:TYR	C	352:ALA	N	3.07

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	1422/1732 (82%)	-0.18	14 (0%) 82 76	64, 116, 174, 247	0
2	B	1115/1224 (91%)	-0.12	12 (1%) 80 74	67, 130, 191, 222	0
3	C	266/318 (83%)	-0.22	0 100 100	90, 119, 163, 183	0
4	D	178/221 (80%)	-0.18	0 100 100	102, 135, 182, 198	0
5	E	214/215 (99%)	-0.17	3 (1%) 75 68	90, 151, 199, 208	0
6	F	84/155 (54%)	-0.29	0 100 100	70, 95, 126, 149	0
7	G	171/171 (100%)	-0.07	0 100 100	87, 116, 155, 179	0
8	H	133/146 (91%)	0.24	5 (3%) 40 33	122, 161, 195, 205	0
9	I	119/122 (97%)	-0.06	3 (2%) 57 49	123, 158, 192, 214	0
10	J	65/70 (92%)	-0.35	0 100 100	97, 115, 153, 166	0
11	K	115/120 (95%)	-0.22	0 100 100	83, 115, 163, 181	0
12	L	46/70 (65%)	0.01	2 (4%) 35 30	103, 159, 184, 191	0
13	N	11/15 (73%)	0.44	0 100 100	203, 219, 272, 273	0
14	T	16/27 (59%)	0.44	0 100 100	170, 216, 267, 270	0
All	All	3955/4606 (85%)	-0.14	39 (0%) 82 76	64, 125, 188, 273	0

The worst 5 of 39 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
5	E	93	MET	3.8
1	A	194	ALA	3.8
8	H	139	ASN	3.7
9	I	120	GLN	3.2
2	B	733	HIS	3.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
14	BRU	T	22	20/21	0.68	0.31	221,231,236,237	0

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
16	MG	A	2458	1/1	0.96	0.30	106,106,106,106	0
15	ZN	I	1122	1/1	0.97	0.04	197,197,197,197	0
15	ZN	A	2456	1/1	0.99	0.06	146,146,146,146	0
15	ZN	B	2225	1/1	0.99	0.21	92,92,92,92	0
15	ZN	I	1121	1/1	0.99	0.11	126,126,126,126	0
15	ZN	L	1071	1/1	0.99	0.07	164,164,164,164	0
15	ZN	A	2457	1/1	0.99	0.15	89,89,89,89	0
15	ZN	C	1269	1/1	1.00	0.09	88,88,88,88	0
15	ZN	J	1066	1/1	1.00	0.24	90,90,90,90	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.