



## wwPDB EM Validation Summary Report ⓘ

Dec 12, 2022 – 07:58 am GMT

PDB ID : 5A5T  
EMDB ID : EMD-3056  
Title : Structure of mammalian eIF3 in the context of the 43S preinitiation complex  
Authors : des-Georges, A.; Dhote, V.; Kuhn, L.; Hellen, C.U.T.; Pestova, T.V.; Frank, J.; Hashem, Y.  
Deposited on : 2015-06-21  
Resolution : 6.00 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev43  
MolProbity : 4.02b-467  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
MapQ : 1.9.9  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.31.3


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*ELECTRON MICROSCOPY*

The reported resolution of this entry is 6.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1362	
2	C	843	
3	E	445	
4	F	364	
5	H	352	
6	K	218	
7	L	564	
8	M	374	

## 2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 25432 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called EUKARYOTIC TRANSLATION INITIATION FACTOR 3 SUBUNIT A.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	600	4935	3107	893	914	21	0	1

- Molecule 2 is a protein called EUKARYOTIC TRANSLATION INITIATION FACTOR 3 SUBUNIT C.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	C	558	4529	2842	805	849	33	0	1

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	577	TYR	ALA	conflict	UNP G1U971

- Molecule 3 is a protein called EUKARYOTIC TRANSLATION INITIATION FACTOR 3 SUBUNIT E.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	E	420	3466	2220	587	639	20	0	1

- Molecule 4 is a protein called EUKARYOTIC TRANSLATION INITIATION FACTOR 3 SUBUNIT F.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
4	F	272	2111	1330	359	410	12	0	0

- Molecule 5 is a protein called EUKARYOTIC TRANSLATION INITIATION FACTOR 3 SUBUNIT H.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
5	H	324	2624	1654	452	503	15	0	0

- Molecule 6 is a protein called EUKARYOTIC TRANSLATION INITIATION FACTOR 3 SUBUNIT K.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
6	K	216	1738	1109	286	330	13	0	1

- Molecule 7 is a protein called UNCHARACTERIZED PROTEIN.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
7	L	373	3110	2010	520	563	17	0	1


- Molecule 8 is a protein called EUKARYOTIC TRANSLATION INITIATION FACTOR 3 SUBUNIT M.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
8	M	366	2919	1850	494	558	17	0	1











• Molecule 6: EUKARYOTIC TRANSLATION INITIATION FACTOR 3 SUBUNIT K





• Molecule 7: UNCHARACTERIZED PROTEIN





THR  
LEU  
LYS  
MET  
GLY  
GLN  
ARG  
PRO

● Molecule 8: EUKARYOTIC TRANSLATION INITIATION FACTOR 3 SUBUNIT M



## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	87192	Depositor
Resolution determination method	Not provided	
CTF correction method	EACH PARTICLE	Depositor
Microscope	FEI/PHILIPS CM300FEG/HE	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	25	Depositor
Minimum defocus (nm)	500	Depositor
Maximum defocus (nm)	4500	Depositor
Magnification	30120	Depositor
Image detector	GATAN K2 (4k x 4k)	Depositor
Maximum map value	0.121	Depositor
Minimum map value	-0.062	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.002	Depositor
Recommended contour level	0.025	Depositor
Map size ( $\text{\AA}$ )	498.0, 498.0, 498.0	wwPDB
Map dimensions	300, 300, 300	wwPDB
Map angles ( $^\circ$ )	90.0, 90.0, 90.0	wwPDB
Pixel spacing ( $\text{\AA}$ )	1.66, 1.66, 1.66	Depositor

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.70	0/5021	1.22	40/6781 (0.6%)
2	C	0.70	0/4608	1.19	35/6219 (0.6%)
3	E	1.61	8/3539 (0.2%)	1.29	43/4788 (0.9%)
4	F	0.69	0/2149	1.23	14/2920 (0.5%)
5	H	0.71	0/2675	1.09	7/3609 (0.2%)
6	K	0.69	0/1773	1.09	5/2398 (0.2%)
7	L	0.74	0/3186	1.18	28/4298 (0.7%)
8	M	0.70	0/2964	1.20	19/4000 (0.5%)
All	All	0.88	8/25915 (0.0%)	1.20	191/35013 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	22
2	C	1	18
3	E	1	11
4	F	0	6
5	H	1	8
6	K	0	4
7	L	2	12
8	M	1	13
All	All	6	94

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	189	LEU	CB-CG	53.28	3.07	1.52
3	E	217	TRP	CD2-CE3	30.41	1.85	1.40
3	E	217	TRP	CD2-CE2	30.28	1.77	1.41
3	E	217	TRP	CE2-CZ2	28.28	1.87	1.39

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	217	TRP	CE3-CZ3	22.94	1.77	1.38

The worst 5 of 191 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	F	128	ARG	NE-CZ-NH1	14.34	127.47	120.30
3	E	251	PRO	CA-C-N	-13.42	87.67	117.20
3	E	251	PRO	C-N-CA	11.49	150.42	121.70
3	E	251	PRO	CA-C-O	11.39	147.53	120.20
3	E	189	LEU	CB-CG-CD1	11.26	130.14	111.00

5 of 6 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	C	513	LYS	CA
3	E	67	PRO	CA
5	H	223	ALA	CA
7	L	418	PRO	CA
7	L	439	PRO	CA

5 of 94 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	165	ARG	Sidechain
1	A	169	ARG	Sidechain
1	A	190	ARG	Sidechain
1	A	195	ARG	Sidechain
1	A	91	ARG	Sidechain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4935	0	5017	3	0
2	C	4529	0	4533	5	0
3	E	3466	0	3446	74	0
4	F	2111	0	2105	2	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	H	2624	0	2592	5	0
6	K	1738	0	1706	3	0
7	L	3110	0	3084	5	0
8	M	2919	0	2950	1	0
All	All	25432	0	25433	94	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 94 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:E:217:TRP:CE3	3:E:217:TRP:CZ3	1.77	1.71
3:E:217:TRP:CZ3	3:E:217:TRP:CH2	1.75	1.62
3:E:217:TRP:CE2	3:E:217:TRP:CZ2	1.87	1.60
3:E:217:TRP:CH2	3:E:217:TRP:CZ2	1.79	1.59
3:E:217:TRP:CE3	3:E:217:TRP:CD2	1.86	1.59

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	598/1362 (44%)	544 (91%)	36 (6%)	18 (3%)	4	28
2	C	556/843 (66%)	512 (92%)	28 (5%)	16 (3%)	4	29
3	E	418/445 (94%)	362 (87%)	37 (9%)	19 (4%)	2	22
4	F	270/364 (74%)	234 (87%)	20 (7%)	16 (6%)	1	17
5	H	322/352 (92%)	277 (86%)	29 (9%)	16 (5%)	2	20
6	K	214/218 (98%)	200 (94%)	12 (6%)	2 (1%)	17	56

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
7	L	371/564 (66%)	331 (89%)	26 (7%)	14 (4%)	3	24
8	M	364/374 (97%)	320 (88%)	30 (8%)	14 (4%)	3	24
All	All	3113/4522 (69%)	2780 (89%)	218 (7%)	115 (4%)	6	25

5 of 115 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	40	HIS
1	A	136	ASP
1	A	166	ASN
1	A	478	CYS
1	A	539	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	551/1245 (44%)	528 (96%)	23 (4%)	30	54
2	C	503/750 (67%)	480 (95%)	23 (5%)	27	52
3	E	384/406 (95%)	361 (94%)	23 (6%)	19	44
4	F	239/282 (85%)	231 (97%)	8 (3%)	38	61
5	H	293/311 (94%)	270 (92%)	23 (8%)	12	36
6	K	190/193 (98%)	188 (99%)	2 (1%)	73	84
7	L	342/516 (66%)	325 (95%)	17 (5%)	24	49
8	M	327/335 (98%)	309 (94%)	18 (6%)	21	47
All	All	2829/4038 (70%)	2692 (95%)	137 (5%)	29	51

5 of 137 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
7	L	439	PRO
8	M	13	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
8	M	261	ASP
3	E	33	ASN
3	E	8	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
4	F	207	HIS
5	H	328	GLN
8	M	348	GLN
7	L	546	GLN
2	C	840	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
3	E	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	E	251:PRO	C	252:HIS	N	1.68




## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-3056. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

### 6.2 Central slices [i](#)


#### 6.2.1 Primary map



X Index: 150



Y Index: 150



Z Index: 150

The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)

### 6.3.1 Primary map



X Index: 138



Y Index: 147



Z Index: 220


The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal surface views [i](#)


### 6.4.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.025. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


## 6.5 Mask visualisation

This section was not generated. No masks/segmentation were deposited.

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [i](#)



The volume at the recommended contour level is 259 nm<sup>3</sup>; this corresponds to an approximate mass of 234 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum [i](#)



\*Reported resolution corresponds to spatial frequency of  $0.167 \text{\AA}^{-1}$


## 8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

## 9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-3056 and PDB model 5A5T. Per-residue inclusion information can be found in section 3 on page 5.

### 9.1 Map-model overlay [i](#)



The images above show the 3D surface view of the map at the recommended contour level 0.025 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.




## 9.2 Q-score mapped to coordinate model [\(i\)](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model [\(i\)](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.025).



















## 9.4 Atom inclusion [i](#)



At the recommended contour level, 78% of all backbone atoms, 55% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.025) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.5485	 0.1070
A	 0.5850	 0.1150
C	 0.6152	 0.1230
E	 0.6365	 0.1200
F	 0.5489	 0.1010
H	 0.4754	 0.0830
K	 0.4361	 0.1020
L	 0.3580	 0.0800
M	 0.6145	 0.1110

