



wwPDB NMR Structure Validation Summary Report ⓘ

Oct 23, 2021 – 05:03 PM EDT

PDB ID : 1A66
Title : SOLUTION NMR STRUCTURE OF THE CORE NFATC1/DNA COM-
PLEX, 18 STRUCTURES
Authors : Zhou, P.; Sun, L.J.; Doetsch, V.; Wagner, G.; Verdine, G.L.
Deposited on : 1998-03-06

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : 2.23.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	B	12	8% (red) 92% (orange)
2	C	12	33% (yellow) 67% (orange)
3	A	178	26% (green) 57% (yellow) 9% (orange) 8% (cyan)

2 Ensemble composition and analysis

This entry contains 18 models. Model 3 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:10-A:40, A:45-A:177 (164)	0.77	3

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 3 single-model clusters were found.

Cluster number	Models
1	1, 2, 3, 5, 6, 8, 9, 11, 12, 13, 15, 17, 18
2	4, 16
Single-model clusters	7; 10; 14

3 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3595 atoms, of which 1704 are hydrogens and 0 are deuteriums.

- Molecule 1 is a DNA chain called DNA (5'-D(*CP*GP*AP*GP*GP*AP*AP*AP*AP*TP*TP*G)-3').

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	P	
1	B	12	385	119	136	52	67	11	0

- Molecule 2 is a DNA chain called DNA (5'-D(*CP*AP*AP*TP*TP*TP*TP*CP*CP*TP*CP*G)-3').

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	P	
2	C	12	376	116	139	37	73	11	0

- Molecule 3 is a protein called CORE NFATC1.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
3	A	178	2834	875	1429	267	258	5	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	ALA	engineered mutation	UNP O95644
A	2	LYS	LEU	engineered mutation	UNP O95644
A	28	ARG	HIS	engineered mutation	UNP O95644


4 Residue-property plots i

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


- Molecule 1: DNA (5'-D(*CP*GP*AP*GP*GP*AP*AP*AP*AP*TP*TP*G)-3')

Chain B: 



- Molecule 2: DNA (5'-D(*CP*AP*AP*TP*TP*TP*TP*CP*CP*TP*CP*G)-3')

Chain C: 




- Molecule 3: CORE NFATC1

Chain A: 







4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 3. Colouring as in section 4.1 above.

- Molecule 1: DNA (5'-D(*CP*GP*AP*GP*GP*AP*AP*AP*AP*TP*TP*G)-3')

Chain B: 

C316
G316
A317
G318
G319
A320
A321
A322
A323
T324
T325
G326

- Molecule 2: DNA (5'-D(*CP*AP*AP*TP*TP*TP*TP*CP*CP*TP*CP*G)-3')

Chain C:  33% 67%

C340
A341
A342
T343
T344
T345
T346
C347
C348
T349
C350
C351

- Molecule 3: CORE NFATC1

Chain A:  33% 40% 17% 8%

M1
K2
D3
V4
Q5
L6
P7
S8
H9
S10
Y13
Y14
L15
R16
I17
E18
V19
Q20
P21
K22
S23
H24
R25
R26
A27
R28
Y29
A37
V38
K39
A40
S41
A42
G43
G44
H45
P46
L47
V48
Q49
L50
E51
G52
Y53
L54
E55
N56
E57
P58
L59
M60
L61
Q62
L63
F64
L65
G66
T67

R71
L72
L73
R74
P75
H76
A77
F78
Y79
Q80
R83
R84
K87
T88
V89
S90
H94
E95
A96
I97
L98
S99
M100
T101
K102
V103
L104
E105
L106
P107
L108
L109
P110
E111
M112
S113
M114
R115
A116
V117
I118
D119
I123
L124
K125
L126
R127
N128
S129
D130
I131
K135
G136
E137

T138
D139
I140
G141
R142
K143
M144
T145
R146
V147
R148
L149
V150
F151
V152
H154
V155
P156
Q157
P158
S159
G160
R161
T162
L163
S164
L165
S169
I172
E173
Q176
R177
S178

5 Refinement protocol and experimental data overview

The models were refined using the following method: *TORSION ANGLE DYNAMICS AND SIMULATED ANNEALING*.

Of the 18 calculated structures, 18 were deposited, based on the following criterion: *NOE VIOLATION <= 0.4 ANGSTROM, DIHEDRAL ANGLE VIOLATION <= 5 DEGREE*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	refinement	3.1
DYANA-1.4	structure solution	
X-PLOR	structure solution	3.1

No chemical shift data was provided.

6 Model quality i

6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	B	1.28±0.02	2±0/281 (0.7± 0.1%)	2.45±0.01	25±1/433 (5.8± 0.3%)
2	C	1.23±0.01	1±0/263 (0.3± 0.2%)	2.26±0.01	17±1/403 (4.3± 0.3%)
3	A	1.02±0.01	0±0/1325 (0.0± 0.0%)	0.80±0.01	0±0/1794 (0.0± 0.0%)
All	All	1.10	51/33642 (0.2%)	1.49	765/47340 (1.6%)

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	B	325	DT	C5-C7	5.54	1.53	1.50	12	17
2	C	345	DT	C5-C7	5.49	1.53	1.50	1	15
1	B	324	DT	C5-C7	5.48	1.53	1.50	3	18
2	C	343	DT	C5-C7	5.11	1.53	1.50	13	1

5 of 59 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	B	326	DG	N7-C8-N9	9.71	117.95	113.10	8	18
1	B	316	DG	N7-C8-N9	9.38	117.79	113.10	16	18
2	C	351	DG	N7-C8-N9	9.23	117.71	113.10	5	18
1	B	319	DG	N7-C8-N9	9.15	117.68	113.10	17	18
1	B	318	DG	N7-C8-N9	8.94	117.57	113.10	11	18

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	B	249	136	136	15±3
2	C	237	139	139	21±3
3	A	1300	1332	1332	127±13
All	All	32148	28926	28926	2773

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 45.

5 of 996 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
3:A:61:LEU:HD13	3:A:153:VAL:HG22	1.11	1.11	14	18
3:A:15:LEU:HD13	3:A:153:VAL:HG21	1.09	1.23	10	18
3:A:37:ALA:HB2	3:A:123:ILE:HG22	1.05	1.17	14	6
3:A:79:TYR:CE2	3:A:126:LEU:HD21	1.04	1.87	13	7
3:A:47:ILE:HD12	3:A:117:VAL:HG13	1.00	1.34	17	12

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	A	164/178 (92%)	114±5 (70±3%)	37±5 (22±3%)	13±3 (8±2%)	2	14
All	All	2952/3204 (92%)	2054 (70%)	659 (22%)	239 (8%)	2	14

5 of 49 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
3	A	99	SER	18

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
3	A	110	PRO	18
3	A	77	ALA	16
3	A	18	GLU	15
3	A	52	GLY	14

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	145/156 (93%)	101±7 (70±5%)	44±7 (30±5%)	1	16
All	All	2610/2808 (93%)	1822 (70%)	788 (30%)	1	16

5 of 113 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
3	A	65	ILE	18
3	A	15	LEU	16
3	A	161	ARG	16
3	A	62	GLN	15
3	A	83	ARG	15

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided