



# wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 17, 2023 – 10:27 am GMT

PDB ID : 4A6N  
Title : STRUCTURE OF THE TETRACYCLINE DEGRADING MONOOXYGENASE TETX IN COMPLEX WITH TIGECYCLINE  
Authors : Volkers, G.; Palm, G.J.; Weiss, M.S.; Hinrichs, W.  
Deposited on : 2011-11-07  
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	398	 8% 79% 13% • 8%
2	B	398	 10% 78% 13% • 8%
2	C	398	 15% 78% 13% • 8%
2	D	398	 12% 80% 11% • 8%

## 2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 12141 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TETX2 PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	368	2886	1827	490	557	12	0	0	0

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-9	MET	-	expression tag	UNP Q93L51
A	-8	GLY	-	expression tag	UNP Q93L51
A	-7	SER	-	expression tag	UNP Q93L51
A	-6	SER	-	expression tag	UNP Q93L51
A	-5	HIS	-	expression tag	UNP Q93L51
A	-4	HIS	-	expression tag	UNP Q93L51
A	-3	HIS	-	expression tag	UNP Q93L51
A	-2	HIS	-	expression tag	UNP Q93L51
A	-1	HIS	-	expression tag	UNP Q93L51
A	0	HIS	-	expression tag	UNP Q93L51
A	1	SER	-	expression tag	UNP Q93L51
A	2	SER	-	expression tag	UNP Q93L51
A	3	GLY	-	expression tag	UNP Q93L51
A	4	LEU	-	expression tag	UNP Q93L51
A	5	VAL	-	expression tag	UNP Q93L51
A	6	PRO	-	expression tag	UNP Q93L51
A	7	ARG	-	expression tag	UNP Q93L51
A	8	GLY	-	expression tag	UNP Q93L51
A	9	SER	-	expression tag	UNP Q93L51
A	10	HIS	-	expression tag	UNP Q93L51
A	272	LYS	GLU	conflict	UNP Q93L51

- Molecule 2 is a protein called TETX2 PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	368	Total 2883	C 1826	N 489	O 556	S 12	0	0	0
2	C	367	Total 2868	C 1817	N 485	O 554	S 12	0	0	0
2	D	367	Total 2859	C 1812	N 485	O 550	S 12	0	0	0

There are 60 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	-9	MET	-	expression tag	UNP Q93L51
B	-8	GLY	-	expression tag	UNP Q93L51
B	-7	SER	-	expression tag	UNP Q93L51
B	-6	SER	-	expression tag	UNP Q93L51
B	-5	HIS	-	expression tag	UNP Q93L51
B	-4	HIS	-	expression tag	UNP Q93L51
B	-3	HIS	-	expression tag	UNP Q93L51
B	-2	HIS	-	expression tag	UNP Q93L51
B	-1	HIS	-	expression tag	UNP Q93L51
B	0	HIS	-	expression tag	UNP Q93L51
B	1	SER	-	expression tag	UNP Q93L51
B	2	SER	-	expression tag	UNP Q93L51
B	3	GLY	-	expression tag	UNP Q93L51
B	4	LEU	-	expression tag	UNP Q93L51
B	5	VAL	-	expression tag	UNP Q93L51
B	6	PRO	-	expression tag	UNP Q93L51
B	7	ARG	-	expression tag	UNP Q93L51
B	8	GLY	-	expression tag	UNP Q93L51
B	9	SER	-	expression tag	UNP Q93L51
B	10	HIS	-	expression tag	UNP Q93L51
C	-9	MET	-	expression tag	UNP Q93L51
C	-8	GLY	-	expression tag	UNP Q93L51
C	-7	SER	-	expression tag	UNP Q93L51
C	-6	SER	-	expression tag	UNP Q93L51
C	-5	HIS	-	expression tag	UNP Q93L51
C	-4	HIS	-	expression tag	UNP Q93L51
C	-3	HIS	-	expression tag	UNP Q93L51
C	-2	HIS	-	expression tag	UNP Q93L51
C	-1	HIS	-	expression tag	UNP Q93L51
C	0	HIS	-	expression tag	UNP Q93L51
C	1	SER	-	expression tag	UNP Q93L51
C	2	SER	-	expression tag	UNP Q93L51
C	3	GLY	-	expression tag	UNP Q93L51

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
C	4	LEU	-	expression tag	UNP Q93L51
C	5	VAL	-	expression tag	UNP Q93L51
C	6	PRO	-	expression tag	UNP Q93L51
C	7	ARG	-	expression tag	UNP Q93L51
C	8	GLY	-	expression tag	UNP Q93L51
C	9	SER	-	expression tag	UNP Q93L51
C	10	HIS	-	expression tag	UNP Q93L51
D	-9	MET	-	expression tag	UNP Q93L51
D	-8	GLY	-	expression tag	UNP Q93L51
D	-7	SER	-	expression tag	UNP Q93L51
D	-6	SER	-	expression tag	UNP Q93L51
D	-5	HIS	-	expression tag	UNP Q93L51
D	-4	HIS	-	expression tag	UNP Q93L51
D	-3	HIS	-	expression tag	UNP Q93L51
D	-2	HIS	-	expression tag	UNP Q93L51
D	-1	HIS	-	expression tag	UNP Q93L51
D	0	HIS	-	expression tag	UNP Q93L51
D	1	SER	-	expression tag	UNP Q93L51
D	2	SER	-	expression tag	UNP Q93L51
D	3	GLY	-	expression tag	UNP Q93L51
D	4	LEU	-	expression tag	UNP Q93L51
D	5	VAL	-	expression tag	UNP Q93L51
D	6	PRO	-	expression tag	UNP Q93L51
D	7	ARG	-	expression tag	UNP Q93L51
D	8	GLY	-	expression tag	UNP Q93L51
D	9	SER	-	expression tag	UNP Q93L51
D	10	HIS	-	expression tag	UNP Q93L51

- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula:  $C_{27}H_{33}N_9O_{15}P_2$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
3	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
3	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
3	C	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
3	D	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 4 is TIGECYCLINE (three-letter code: T1C) (formula:  $C_{29}H_{41}N_5O_8$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			42	29	5	8		
4	B	1	Total	C	N	O	0	0
			42	29	5	8		
4	C	1	Total	C	N	O	0	0
			42	29	5	8		
4	D	1	Total	C	N	O	0	0
			42	29	5	8		

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	O S	0	0
			5	4 1		
5	A	1	Total	O S	0	0
			5	4 1		
5	B	1	Total	O S	0	0
			5	4 1		
5	B	1	Total	O S	0	0
			5	4 1		
5	B	1	Total	O S	0	0
			5	4 1		
5	C	1	Total	O S	0	0
			5	4 1		
5	C	1	Total	O S	0	0
			5	4 1		

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	D	1	Total	O	S	0	0
			5	4	1		
5	D	1	Total	O	S	0	0
			5	4	1		
5	D	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is water.



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	61	Total	O	0	0
			61	61		
6	B	68	Total	O	0	0
			68	68		
6	C	35	Total	O	0	0
			35	35		
6	D	46	Total	O	0	0
			46	46		







● Molecule 2: TETX2 PROTEIN



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	68.88Å 80.79Å 87.65Å 110.84° 89.84° 93.63°	Depositor
Resolution (Å)	81.90 – 2.30 40.95 – 2.30	Depositor EDS
% Data completeness (in resolution range)	96.0 (81.90-2.30) 96.1 (40.95-2.30)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.08 (at 2.29Å)	Xtrriage
Refinement program	REFMAC 5.6.0116	Depositor
R, $R_{free}$	0.218 , 0.260 0.227 , 0.267	Depositor DCC
$R_{free}$ test set	3769 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	43.0	Xtrriage
Anisotropy	0.334	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 59.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	12141	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	72.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.33% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: T1C, FAD, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.85	0/2944	0.89	5/3987 (0.1%)
2	B	0.84	0/2941	0.90	7/3983 (0.2%)
2	C	0.70	1/2926 (0.0%)	0.80	3/3965 (0.1%)
2	D	0.70	0/2917	0.80	4/3954 (0.1%)
All	All	0.78	1/11728 (0.0%)	0.85	19/15889 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	298	TRP	CD2-CE2	5.54	1.48	1.41

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	175	ARG	NE-CZ-NH1	8.96	124.78	120.30
2	B	251	VAL	CG1-CB-CG2	8.76	124.92	110.90
2	B	175	ARG	NE-CZ-NH1	8.12	124.36	120.30
2	C	175	ARG	NE-CZ-NH1	7.27	123.94	120.30
2	D	175	ARG	NE-CZ-NH1	7.17	123.88	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2886	0	2807	32	0
2	B	2883	0	2797	36	0
2	C	2868	0	2771	31	0
2	D	2859	0	2761	33	0
3	A	53	0	31	4	0
3	B	53	0	31	8	0
3	C	53	0	31	1	0
3	D	53	0	31	2	0
4	A	42	0	39	10	0
4	B	42	0	39	7	0
4	C	42	0	40	5	0
4	D	42	0	38	15	0
5	A	10	0	0	0	0
5	B	20	0	0	0	0
5	C	10	0	0	0	0
5	D	15	0	0	0	0
6	A	61	0	0	0	0
6	B	68	0	0	2	0
6	C	35	0	0	4	0
6	D	46	0	0	5	0
All	All	12141	0	11416	154	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 154 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:389:FAD:HM73	4:A:392:T1C:H421	1.35	1.06
2:C:269:ASP:HB2	6:C:2030:HOH:O	1.59	1.02
2:D:279:HIS:CE1	6:D:2037:HOH:O	2.13	0.98
4:D:392:T1C:C72	4:D:392:T1C:H62C	2.03	0.87
4:D:392:T1C:H422	4:D:392:T1C:O3	1.81	0.81

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	364/398 (92%)	351 (96%)	12 (3%)	1 (0%)	41	50
2	B	364/398 (92%)	350 (96%)	13 (4%)	1 (0%)	41	50
2	C	363/398 (91%)	348 (96%)	14 (4%)	1 (0%)	41	50
2	D	363/398 (91%)	348 (96%)	14 (4%)	1 (0%)	41	50
All	All	1454/1592 (91%)	1397 (96%)	53 (4%)	4 (0%)	41	50

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	323	GLY
1	A	323	GLY
2	D	323	GLY
2	C	323	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	312/345 (90%)	302 (97%)	10 (3%)	39	54
2	B	310/345 (90%)	298 (96%)	12 (4%)	32	46
2	C	307/345 (89%)	295 (96%)	12 (4%)	32	46
2	D	305/345 (88%)	295 (97%)	10 (3%)	38	53
All	All	1234/1380 (89%)	1190 (96%)	44 (4%)	35	49

5 of 44 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	269	ASP
2	D	127	SER
2	C	273	ARG
2	C	300	SER
2	D	192	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 28 such sidechains are listed below:

Mol	Chain	Res	Type
2	C	38	ASN
2	D	371	ASN
2	C	190	ASN
2	D	192	GLN
2	C	130	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

19 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	SO4	D	1384	-	4,4,4	0.38	0	6,6,6	0.30	0
4	T1C	B	392	-	44,45,45	2.03	10 (22%)	53,72,72	2.26	15 (28%)
5	SO4	A	1384	-	4,4,4	0.48	0	6,6,6	0.46	0
5	SO4	A	1383	-	4,4,4	0.32	0	6,6,6	0.34	0
5	SO4	D	1383	-	4,4,4	0.35	0	6,6,6	0.18	0
5	SO4	D	1385	-	4,4,4	0.41	0	6,6,6	0.43	0
4	T1C	A	392	-	44,45,45	2.02	11 (25%)	53,72,72	1.82	12 (22%)
5	SO4	C	1384	-	4,4,4	0.44	0	6,6,6	0.32	0
5	SO4	B	1384	-	4,4,4	0.39	0	6,6,6	0.35	0
3	FAD	B	389	-	53,58,58	1.60	10 (18%)	68,89,89	1.52	12 (17%)
5	SO4	C	1383	-	4,4,4	0.55	0	6,6,6	0.34	0
5	SO4	B	1383	-	4,4,4	0.33	0	6,6,6	0.35	0
3	FAD	A	389	-	53,58,58	1.57	10 (18%)	68,89,89	1.58	13 (19%)
4	T1C	D	392	-	44,45,45	1.80	9 (20%)	53,72,72	1.98	14 (26%)
3	FAD	D	389	-	53,58,58	1.62	11 (20%)	68,89,89	1.89	20 (29%)
5	SO4	B	1385	-	4,4,4	0.36	0	6,6,6	0.26	0
3	FAD	C	389	-	53,58,58	1.44	7 (13%)	68,89,89	1.75	14 (20%)
4	T1C	C	392	-	44,45,45	1.79	10 (22%)	53,72,72	1.82	10 (18%)
5	SO4	B	1386	-	4,4,4	0.43	0	6,6,6	0.30	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FAD	C	389	-	-	3/30/50/50	0/6/6/6
4	T1C	C	392	-	-	5/22/80/80	0/4/4/4
4	T1C	B	392	-	-	0/22/80/80	0/4/4/4
3	FAD	A	389	-	-	1/30/50/50	0/6/6/6
4	T1C	D	392	-	-	7/22/80/80	0/4/4/4
3	FAD	D	389	-	-	4/30/50/50	0/6/6/6
3	FAD	B	389	-	-	1/30/50/50	0/6/6/6
4	T1C	A	392	-	-	3/22/80/80	0/4/4/4

The worst 5 of 78 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	392	T1C	C2-C21	6.89	1.61	1.47

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	392	T1C	C1C-C12	6.47	1.57	1.52
4	C	392	T1C	C2-C21	5.91	1.59	1.47
3	A	389	FAD	C9A-C5X	5.77	1.50	1.41
4	B	392	T1C	C2-C21	5.73	1.59	1.47

The worst 5 of 110 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	392	T1C	O12-C12-C1C	7.03	123.55	113.37
3	D	389	FAD	C5'-C4'-C3'	-6.85	98.97	112.20
4	C	392	T1C	C1C-C1-C2	6.35	125.83	115.75
3	A	389	FAD	C5'-C4'-C3'	-5.59	101.40	112.20
4	D	392	T1C	C1C-C1-C2	5.44	124.40	115.75

There are no chirality outliers.

5 of 24 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	C	392	T1C	C41-C4-N4-C42
4	D	392	T1C	C94-C93-N92-C92
4	D	392	T1C	C95-C93-N92-C92
4	D	392	T1C	C41-C4-N4-C43
4	D	392	T1C	C96-C93-N92-C92


There are no ring outliers.


8 monomers are involved in 45 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	392	T1C	7	0
4	A	392	T1C	10	0
3	B	389	FAD	8	0
3	A	389	FAD	4	0
4	D	392	T1C	15	0
3	D	389	FAD	2	0
3	C	389	FAD	1	0
4	C	392	T1C	5	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is


within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	368/398 (92%)	0.68	32 (8%) 10 14	43, 62, 96, 142	0
2	B	368/398 (92%)	0.79	38 (10%) 6 9	45, 63, 95, 130	0
2	C	367/398 (92%)	0.95	60 (16%) 1 2	53, 75, 111, 156	0
2	D	367/398 (92%)	0.89	46 (12%) 3 5	50, 75, 115, 144	0
All	All	1470/1592 (92%)	0.83	176 (11%) 4 6	43, 69, 108, 156	0

The worst 5 of 176 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	340	ASP	8.7
2	B	249	THR	7.3
1	A	249	THR	6.9
1	A	343	PHE	6.5
2	C	380	PHE	6.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.






Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	SO4	A	1384	5/5	0.37	0.25	109,115,142,145	0
4	T1C	C	392	42/42	0.66	0.35	77,120,163,175	0
5	SO4	C	1383	5/5	0.74	0.16	99,100,115,129	0
4	T1C	D	392	42/42	0.77	0.31	91,115,142,151	0
4	T1C	B	392	42/42	0.78	0.24	61,90,115,117	0
4	T1C	A	392	42/42	0.83	0.22	61,85,102,117	0
5	SO4	B	1385	5/5	0.88	0.12	105,109,124,129	0
5	SO4	D	1383	5/5	0.89	0.20	117,119,129,130	0
5	SO4	C	1384	5/5	0.91	0.21	90,97,107,118	0
3	FAD	B	389	53/53	0.92	0.19	33,45,58,72	0
5	SO4	B	1383	5/5	0.94	0.15	98,100,106,119	0
3	FAD	C	389	53/53	0.94	0.14	36,50,62,73	0
5	SO4	B	1386	5/5	0.94	0.24	80,96,107,108	0
5	SO4	D	1385	5/5	0.94	0.24	90,93,100,108	0
5	SO4	B	1384	5/5	0.95	0.11	89,95,103,109	0
3	FAD	A	389	53/53	0.96	0.14	30,43,53,65	0
3	FAD	D	389	53/53	0.96	0.14	39,50,61,69	0
5	SO4	A	1383	5/5	0.96	0.11	94,96,104,108	0
5	SO4	D	1384	5/5	0.98	0.08	88,88,92,98	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around T1C C 392:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around T1C D 392:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around T1C B 392:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around T1C A 392:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around FAD B 389:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around FAD C 389:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around FAD A 389:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around FAD D 389:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.