



Full wwPDB X-ray Structure Validation Report i

Oct 26, 2023 – 01:33 PM EDT

PDB ID : 3A7A
Title : Crystal structure of E. coli lipoate-protein ligase A in complex with octyl-amp and apoH-protein
Authors : Fujiwara, K.; Hosaka, H.; Nakagawa, A.
Deposited on : 2009-09-20
Resolution : 3.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1094 (3.10-3.10)
Clashscore	141614	1184 (3.10-3.10)
Ramachandran outliers	138981	1141 (3.10-3.10)
Sidechain outliers	138945	1141 (3.10-3.10)
RSRZ outliers	127900	1067 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition [\(i\)](#)

There are 4 unique types of molecules in this entry. The entry contains 7300 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Lipoate-protein ligase A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	337	Total	C 2663	N 1673	O 476	S 503	11	0	0
1	C	337	Total	C 2663	N 1673	O 476	S 503	11	0	0

- Molecule 2 is a protein called Glycine cleavage system H protein.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	127	Total	C 956	N 600	O 146	S 208	2	0	0
2	D	127	Total	C 956	N 600	O 146	S 208	2	0	0

- Molecule 3 is N-OCTANE (three-letter code: OCT) (formula: C₈H₁₈).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C 8 8	0	0
3	C	1	Total C 8 8	0	0

- Molecule 4 is ADENOSINE MONOPHOSPHATE (three-letter code: AMP) (formula: C₁₀H₁₄N₅O₇P).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C N O P 23 10 5 7 1	0	0
4	C	1	Total C N O P 23 10 5 7 1	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Lipoate-protein ligase A





- Molecule 1: Lipoate-protein ligase A



- Molecule 2: Glycine cleavage system H protein





- Molecule 2: Glycine cleavage system H protein



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	70.01 Å 102.02 Å 159.93 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 3.10 48.60 – 2.96	Depositor EDS
% Data completeness (in resolution range)	98.3 (20.00-3.10) 96.6 (48.60-2.96)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	3.56 (at 2.96 Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R , R_{free}	0.228 , 0.279 0.210 , 0.277	Depositor DCC
R_{free} test set	1215 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å ²)	75.8	Xtriage
Anisotropy	0.554	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 23.2	EDS
L-test for twinning ²	$< L > = 0.49$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7300	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.85% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:287:PRO:CD	1:C:288:ALA:H	2.35	0.40
1:A:38:ARG:HD3	1:A:196:HIS:CE1	2.57	0.40
1:A:147:PHE:C	1:A:147:PHE:HD2	2.24	0.40
1:C:55:CYS:HB2	1:C:164:TYR:O	2.21	0.40
1:C:287:PRO:O	1:C:288:ALA:C	2.58	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:1:SER:CA	1:C:274:HIS:NE2[4_555]	2.05	0.15

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	335/337 (99%)	307 (92%)	20 (6%)	8 (2%)	6 27
1	C	335/337 (99%)	294 (88%)	32 (10%)	9 (3%)	5 25
2	B	125/128 (98%)	103 (82%)	16 (13%)	6 (5%)	2 14
2	D	125/128 (98%)	112 (90%)	12 (10%)	1 (1%)	19 54
All	All	920/930 (99%)	816 (89%)	80 (9%)	24 (3%)	5 26

All (24) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	224	LYS
1	C	201	GLU
1	C	316	VAL
1	A	41	ASP
1	A	232	ALA

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	302	TYR
1	C	41	ASP
1	C	81	LEU
2	D	97	ALA
1	A	231	PHE
1	A	260	THR
2	B	36	LEU
2	B	63	VAL
2	B	82	ASP
2	B	85	SER
1	C	230	ASN
1	C	231	PHE
1	C	232	ALA
2	B	4	PRO
2	B	21	ASP
1	A	222	PRO
1	C	2	THR
1	A	288	ALA
1	C	287	PRO

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	282/282 (100%)	264 (94%)	18 (6%)	17 48
1	C	282/282 (100%)	261 (93%)	21 (7%)	13 42
2	B	102/103 (99%)	98 (96%)	4 (4%)	32 65
2	D	102/103 (99%)	98 (96%)	4 (4%)	32 65
All	All	768/770 (100%)	721 (94%)	47 (6%)	18 49

All (47) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	26	GLN
1	A	32	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	64	ASN
1	A	147	PHE
1	A	149	HIS
1	A	181	ARG
1	A	188	THR
1	A	230	ASN
1	A	234	THR
1	A	255	LEU
1	A	277	ARG
1	A	280	VAL
1	A	285	LEU
1	A	297	LEU
1	A	312	GLU
1	A	314	LEU
1	A	323	LYS
1	A	326	ARG
2	B	56	ASP
2	B	89	GLU
2	B	105	LYS
2	B	127	ASP
1	C	2	THR
1	C	25	ARG
1	C	26	GLN
1	C	32	ARG
1	C	53	LYS
1	C	83	ASN
1	C	97	LYS
1	C	121	ASN
1	C	140	ARG
1	C	143	LYS
1	C	147	PHE
1	C	149	HIS
1	C	169	LYS
1	C	178	THR
1	C	224	LYS
1	C	233	GLU
1	C	255	LEU
1	C	272	LYS
1	C	307	LEU
1	C	314	LEU
1	C	326	ARG
2	D	11	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	D	68	ASP
2	D	82	ASP
2	D	116	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	106	ASN
1	A	230	ASN
1	A	267	HIS
1	A	298	GLN
2	B	13	HIS
2	B	81	ASN
1	C	106	ASN
1	C	155	ASN
1	C	267	HIS
1	C	274	HIS
1	C	279	GLN
2	D	31	HIS
2	D	81	ASN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and


Mol	Chain	Res	Type	Atoms
4	A	402	AMP	C5'-O5'-P-O2P
4	A	402	AMP	C5'-O5'-P-O3P
4	C	402	AMP	C3'-C4'-C5'-O5'
4	C	402	AMP	O4'-C4'-C5'-O5'
4	A	402	AMP	C3'-C4'-C5'-O5'
3	C	401	OCT	C4-C5-C6-C7
4	A	402	AMP	C5'-O5'-P-O1P
3	A	401	OCT	C1-C2-C3-C4
3	C	401	OCT	C2-C3-C4-C5
3	A	401	OCT	C4-C5-C6-C7
3	A	401	OCT	C2-C3-C4-C5
3	C	401	OCT	C3-C4-C5-C6
4	A	402	AMP	O4'-C4'-C5'-O5'

There are no ring outliers.

3 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	401	OCT	2	0
4	C	402	AMP	3	0
3	A	401	OCT	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	337/337 (100%)	-0.44	0 100 100	27, 36, 50, 54	0
1	C	337/337 (100%)	-0.37	1 (0%) 94 88	28, 43, 61, 63	0
2	B	127/128 (99%)	-0.36	0 100 100	52, 61, 96, 98	0
2	D	127/128 (99%)	-0.11	3 (2%) 59 37	54, 65, 98, 104	0
All	All	928/930 (99%)	-0.36	4 (0%) 92 84	27, 45, 84, 104	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	128	GLU	4.0
2	D	127	ASP	3.9
1	C	1	SER	3.3
2	D	126	GLU	2.6

6.2 Non-standard residues in protein, DNA, RNA chains i

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates i


There are no monosaccharides in this entry.


6.4 Ligands i

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	OCT	A	401	8/8	0.97	0.24	32,32,33,34	0
3	OCT	C	401	8/8	0.97	0.30	44,44,45,46	0
4	AMP	C	402	23/23	0.97	0.15	47,47,47,47	0
4	AMP	A	402	23/23	0.98	0.16	34,36,36,36	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.