



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2024 – 05:25 PM EST

PDB ID : 3AC0
Title : Crystal structure of Beta-glucosidase from *Kluyveromyces marxianus* in complex with glucose
Authors : Yoshida, E.; Hidaka, M.; Fushinobu, S.; Katayama, T.; Kumagai, H.
Deposited on : 2009-12-25
Resolution : 2.54 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.54 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1284 (2.56-2.52)
Clashscore	141614	1332 (2.56-2.52)
Ramachandran outliers	138981	1315 (2.56-2.52)
Sidechain outliers	138945	1315 (2.56-2.52)
RSRZ outliers	127900	1272 (2.56-2.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	845	
1	B	845	
1	C	845	
1	D	845	

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 27663 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-glucosidase I.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	841	Total 6587	C 4184	N 1107	O 1281	S 15	0	0	0
1	B	825	Total 6462	C 4104	N 1091	O 1253	S 14	0	0	0
1	C	831	Total 6514	C 4143	N 1093	O 1263	S 15	0	0	0
1	D	829	Total 6488	C 4128	N 1092	O 1253	S 15	0	0	0

- Molecule 2 is beta-D-glucopyranose (three-letter code: BGC) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	Total 12	C 6	O 6	0	0
2	B	1	Total 12	C 6	O 6	0	0

Continued on next page...

Continued from previous page...



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	C	1	Total	C	O	0	0
			12	6	6		
2	D	1	Total	C	O	0	0
			12	6	6		

- Molecule 3 is water.



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	448	Total	O	0	0
			448	448		
3	B	446	Total	O	0	0
			446	446		
3	C	345	Total	O	0	0
			345	345		
3	D	325	Total	O	0	0
			325	325		




• Molecule 1: Beta-glucosidase I



• Molecule 1: Beta-glucosidase I





4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	245.85Å 148.41Å 119.64Å 90.00° 112.84° 90.00°	Depositor
Resolution (Å)	50.00 – 2.54 42.98 – 2.54	Depositor EDS
% Data completeness (in resolution range)	99.5 (50.00-2.54) 99.6 (42.98-2.54)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.11 (at 2.54Å)	Xtrriage
Refinement program	REFMAC 5.5.0072	Depositor
R, R_{free}	0.169 , 0.242 0.165 , 0.233	Depositor DCC
R_{free} test set	6526 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	36.7	Xtrriage
Anisotropy	0.011	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 46.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	27663	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BGC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.89	0/6723	0.82	3/9089 (0.0%)
1	B	0.94	3/6591 (0.0%)	0.86	7/8907 (0.1%)
1	C	0.83	1/6646 (0.0%)	0.81	4/8981 (0.0%)
1	D	0.82	3/6622 (0.0%)	0.80	2/8952 (0.0%)
All	All	0.87	7/26582 (0.0%)	0.82	16/35929 (0.0%)

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	816	GLU	CG-CD	6.68	1.61	1.51
1	B	264	GLU	CD-OE2	-5.95	1.19	1.25
1	D	269	GLU	CG-CD	5.79	1.60	1.51
1	D	264	GLU	CD-OE2	-5.70	1.19	1.25
1	B	264	GLU	CD-OE1	-5.21	1.20	1.25

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	579	LEU	CA-CB-CG	8.51	134.86	115.30
1	B	249	ARG	NE-CZ-NH2	-8.23	116.18	120.30
1	A	26	ASP	CB-CG-OD1	6.59	124.23	118.30
1	B	769	ARG	NE-CZ-NH2	-6.26	117.17	120.30
1	A	42	ARG	NE-CZ-NH2	-5.96	117.32	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6587	0	6490	96	0
1	B	6462	0	6387	66	0
1	C	6514	0	6427	103	0
1	D	6488	0	6417	113	0
2	A	12	0	12	2	0
2	B	12	0	12	1	0
2	C	12	0	12	1	0
2	D	12	0	12	3	0
3	A	448	0	0	19	0
3	B	446	0	0	11	0
3	C	345	0	0	12	0
3	D	325	0	0	10	0
All	All	27663	0	25769	372	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 372 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:827:ASP:HB3	3:B:1558:HOH:O	1.46	1.12
1:D:562:GLU:HG2	3:D:1448:HOH:O	1.60	1.01
1:B:633:GLU:HG2	3:B:999:HOH:O	1.63	0.97
1:A:737:ILE:HD11	1:A:796:LEU:HD23	1.45	0.96
1:C:763:LEU:HD11	1:C:818:LEU:HD13	1.47	0.94

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	837/845 (99%)	792 (95%)	44 (5%)	1 (0%)	51	65
1	B	815/845 (96%)	776 (95%)	38 (5%)	1 (0%)	51	65
1	C	821/845 (97%)	773 (94%)	46 (6%)	2 (0%)	47	60
1	D	823/845 (97%)	773 (94%)	50 (6%)	0	100	100
All	All	3296/3380 (98%)	3114 (94%)	178 (5%)	4 (0%)	51	65

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	524	GLY
1	B	297	ASN
1	C	329	ASN
1	C	524	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	714/717 (100%)	685 (96%)	29 (4%)	30	41
1	B	700/717 (98%)	664 (95%)	36 (5%)	24	32
1	C	707/717 (99%)	662 (94%)	45 (6%)	17	23
1	D	703/717 (98%)	669 (95%)	34 (5%)	25	34
All	All	2824/2868 (98%)	2680 (95%)	144 (5%)	24	32

5 of 144 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	366	SER
1	D	845	LEU
1	D	411	SER
1	D	530	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	576	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 33 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	443	HIS
1	D	480	GLN
1	D	764	ASN
1	B	525	GLN
1	B	338	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BGC	B	5002	-	12,12,12	0.81	0	17,17,17	2.16	4 (23%)
2	BGC	A	5001	-	12,12,12	0.69	0	17,17,17	1.59	3 (17%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BGC	C	5003	-	12,12,12	0.60	0	17,17,17	0.75	0
2	BGC	D	5004	-	12,12,12	0.84	0	17,17,17	1.66	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	B	5002	-	-	0/2/22/22	0/1/1/1
2	BGC	A	5001	-	-	0/2/22/22	0/1/1/1
2	BGC	C	5003	-	-	0/2/22/22	0/1/1/1
2	BGC	D	5004	-	-	0/2/22/22	0/1/1/1

There are no bond length outliers.

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	5002	BGC	C1-O5-C5	5.56	124.15	113.66
2	B	5002	BGC	O1-C1-O5	-4.95	95.51	110.38
2	A	5001	BGC	C1-O5-C5	4.50	122.15	113.66
2	D	5004	BGC	O1-C1-O5	-3.25	100.63	110.38
2	D	5004	BGC	C3-C4-C5	-3.18	104.57	110.24

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	5002	BGC	1	0
2	A	5001	BGC	2	0
2	C	5003	BGC	1	0
2	D	5004	BGC	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	841/845 (99%)	-0.27	10 (1%) 79 84	15, 31, 60, 77	0
1	B	825/845 (97%)	-0.18	28 (3%) 45 52	15, 26, 76, 97	0
1	C	831/845 (98%)	-0.00	50 (6%) 21 25	18, 38, 87, 105	0
1	D	829/845 (98%)	0.04	29 (3%) 44 51	19, 43, 80, 99	0
All	All	3326/3380 (98%)	-0.10	117 (3%) 44 51	15, 35, 80, 105	0

The worst 5 of 117 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	406	ALA	5.2
1	A	543	GLY	5.0
1	C	403	ALA	4.9
1	D	403	ALA	4.9
1	D	402	ALA	4.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	BGC	A	5001	12/12	0.96	0.19	20,28,32,37	0
2	BGC	D	5004	12/12	0.97	0.19	32,36,38,41	0
2	BGC	C	5003	12/12	0.98	0.16	28,34,38,42	0
2	BGC	B	5002	12/12	0.98	0.12	25,28,30,33	0

6.5 Other polymers [i](#)

There are no such residues in this entry.