

wwPDB EM Validation Summary Report (i)

Nov 9, 2022 – 01:39 AM JST

PDB ID : 6ACK EMDB ID : EMD-9594

Title: Trypsin-cleaved and low pH-treated SARS-CoV spike glycoprotein and ACE2

complex, ACE2-bound conformation 3

Authors : Gui, M.; Song, W.

Deposited on : 2018-07-26

Resolution : 4.50 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev43

MolProbity : 4.02b-467

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

MapQ: 1.9.9

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.31.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 4.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	${ m EM\ structures} \ (\#{ m Entries})$
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion <40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain						
			39%						
1	A	1203	54%	33% · 11%					
			37%						
1	В	1203	57%	30% • 11%					
			41%						
1	С	1203	55%	31% • 12%					
			99%						
2	D	603	72%	27% •					



2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 29715 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Spike glycoprotein.

Mol	Chain	Residues	Atoms				AltConf	Trace	
1	A	1065	Total	С	N	О	S	0	0
1	A	1005	8302	5304	1374	1579	45	U	
1	В	1065	Total	С	N	О	S	0	0
1	Б	1005	8302	5304	1374	1579	45	0	
1	С	1057	Total	С	N	О	S	0	0
1	C	1057	8241	5264	1364	1568	45	0	

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1197	SER	-	expression tag	UNP P59594
A	1198	HIS	-	expression tag	UNP P59594
A	1199	PRO	-	expression tag	UNP P59594
A	1200	GLN	-	expression tag	UNP P59594
A	1201	PHE	-	expression tag	UNP P59594
A	1202	GLU	-	expression tag	UNP P59594
A	1203	LYS	-	expression tag	UNP P59594
В	1197	SER	-	expression tag	UNP P59594
В	1198	HIS	-	expression tag	UNP P59594
В	1199	PRO	-	expression tag	UNP P59594
В	1200	GLN	-	expression tag	UNP P59594
В	1201	PHE	-	expression tag	UNP P59594
В	1202	GLU	-	expression tag	UNP P59594
В	1203	LYS	-	expression tag	UNP P59594
С	1197	SER	-	expression tag	UNP P59594
С	1198	HIS	-	expression tag	UNP P59594
С	1199	PRO	-	expression tag	UNP P59594
С	1200	GLN	-	expression tag	UNP P59594
С	1201	PHE	-	expression tag	UNP P59594
С	1202	GLU		expression tag	UNP P59594
С	1203	LYS	_	expression tag	UNP P59594

• Molecule 2 is a protein called Angiotensin-converting enzyme 2.



Mol	Chain	Residues		${f Atoms}$				AltConf	Trace
2	D	597	Total 4870	C 3115	N 806	O 920	S 29	0	0

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference	
D	616	HIS	-	expression tag	UNP Q9BYF1	
D	617	HIS	-	expression tag	UNP Q9BYF1	
D	618	HIS	-	expression tag	UNP Q9BYF1	
D	619	HIS	-	expression tag	UNP Q9BYF1	
D	620	HIS	-	expression tag	UNP Q9BYF1	
D	621	HIS	-	expression tag	UNP Q9BYF1	




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Spike glycoprotein









• Molecule 1: Spike glycoprotein






• Molecule 1: Spike glycoprotein











• Molecule 2: Angiotensin-converting enzyme 2











4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	56553	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE	Depositor
	CORRECTION	
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{Å}^2)$	50	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	29.588	Depositor
Minimum map value	-14.552	Depositor
Average map value	-0.001	Depositor
Map value standard deviation	0.908	Depositor
Recommended contour level	8.0	Depositor
Map size (Å)	380.16, 380.16, 380.16	wwPDB
Map dimensions	288, 288, 288	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.32, 1.32, 1.32	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	nd lengths	Bond angles		
WIOI		RMSZ	# Z > 5	RMSZ	# Z >5	
1	A	0.54	$2/8499 \ (0.0\%)$	0.73	5/11568 (0.0%)	
1	В	0.53	1/8499 (0.0%)	0.73	2/11568 (0.0%)	
1	С	0.59	5/8435 (0.1%)	0.76	7/11477 (0.1%)	
2	D	0.34	0/5007	0.58	0/6803	
All	All	0.53	8/30440 (0.0%)	0.72	14/41416 (0.0%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	$\mid \# ext{Planarity outliers} \mid$
1	A	0	13
1	В	0	14
1	С	0	18
All	All	0	45

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(\mathbf{\mathring{A}})$	$Ideal(\AA)$
1	A	725	CYS	CB-SG	-9.21	1.66	1.82
1	A	731	CYS	CB-SG	-8.50	1.67	1.82
1	С	677	TYR	CE2-CZ	-8.19	1.27	1.38
1	В	725	CYS	CB-SG	-6.59	1.71	1.82
1	С	676	ALA	C-O	-6.38	1.11	1.23

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
1	С	677	TYR	CB-CG-CD2	-6.45	117.13	121.00
1	С	674	ILE	CB-CA-C	-6.36	98.89	111.60
1	В	557	ASP	CB-CG-OD1	6.35	124.02	118.30
1	С	644	THR	O-C-N	5.97	132.25	122.70

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
1	С	644	THR	CA-C-N	-5.78	104.49	117.20

There are no chirality outliers.

5 of 45 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	506	ALA	Peptide
1	A	629	PHE	Peptide
1	A	632	GLN	Peptide
1	A	633	ALA	Peptide
1	A	95	SER	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	8302	0	8082	286	0
1	В	8302	0	8082	259	0
1	С	8241	0	8009	293	0
2	D	4870	0	4643	99	0
All	All	29715	0	28816	884	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 884 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:C:646:TYR:HB2	1:C:677:TYR:CE2	1.47	1.48
1:C:646:TYR:CB	1:C:677:TYR:CE2	2.02	1.41
1:C:646:TYR:O	1:C:680:SER:OG	1.62	1.15
1:C:646:TYR:HB2	1:C:677:TYR:CD2	1.84	1.11
1:C:646:TYR:CA	1:C:677:TYR:CE2	2.36	1.07

There are no symmetry-related clashes.



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	A	1057/1203 (88%)	874 (83%)	180 (17%)	3 (0%)	41	76
1	В	1057/1203 (88%)	852 (81%)	201 (19%)	4 (0%)	34	72
1	С	1045/1203 (87%)	843 (81%)	192 (18%)	10 (1%)	15	54
2	D	595/603 (99%)	564 (95%)	31 (5%)	0	100	100
All	All	3754/4212 (89%)	3133 (84%)	604 (16%)	17 (0%)	32	68

5 of 17 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	554	ASP
1	В	560	ASP
1	В	1072	PRO
1	С	645	SER
1	В	632	GLN

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	A	922/1048 (88%)	916 (99%)	6 (1%)	84	90
1	В	922/1048 (88%)	917 (100%)	5 (0%)	88	93
1	С	914/1048 (87%)	903 (99%)	11 (1%)	71	84
2	D	527/533 (99%)	521 (99%)	6 (1%)	73	85
All	All	3285/3677 (89%)	3257 (99%)	28 (1%)	79	87



5 of 28 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	С	333	LYS
2	D	436	ILE
1	С	674	ILE
2	D	114	LYS
1	С	646	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 64 such sidechains are listed below:

Mol	Chain	Res	Type
1	С	1005	ASN
2	D	33	ASN
1	В	699	ASN
1	В	692	ASN
2	D	53	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.




6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-9594. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections (i)


6.1.1 Primary map




The images above show the map projected in three orthogonal directions.

6.2 Central slices (i)


6.2.1 Primary map







Y Index: 144




Z Index: 144





The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)

6.3.1 Primary map







X Index: 146


Y Index: 149

Z Index: 141

The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal surface views (i)

6.4.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 8.0. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.




6.5 Mask visualisation (i)


This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency

A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure

6.5.1 emd_9594_msk_1.map (i)








7 Map analysis (i)

This section contains the results of statistical analysis of the map.


7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



7.2 Volume estimate (i)




The volume at the recommended contour level is $119~\mathrm{nm}^3$; this corresponds to an approximate mass of $108~\mathrm{kDa}$.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



7.3 Rotationally averaged power spectrum (i)




^{*}Reported resolution corresponds to spatial frequency of 0.222 $\rm \mathring{A}^{-1}$



8 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC (i)



*Reported resolution corresponds to spatial frequency of 0.222 $\rm \AA^{-1}$



8.2 Resolution estimates (i)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
resolution estimate (A)	0.143	0.5	Half-bit
Reported by author	4.50	-	-
Author-provided FSC curve	4.43	6.79	4.48
Unmasked-calculated*	-	-	-


^{*}Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.



9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-9594 and PDB model 6ACK. Per-residue inclusion information can be found in section 3 on page 5.


9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 8.0 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.




9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (8.0).



9.4 Atom inclusion (i)



At the recommended contour level, 51% of all backbone atoms, 36% of all non-hydrogen atoms, are inside the map.



9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (8.0) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.3592	0.2390
A	0.4339	0.2750
В	0.4372	0.2630
С	0.4106	0.2560
D	0.0115	0.1050



