



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 18, 2021 – 08:01 AM BST

PDB ID : 7ACM  
Title : Crystal structure of E. coli HTH-type transcriptional regulator RcdA in complex with TMAO at 1.76 Å resolution  
Authors : Pietrzyk-Brzezinska, A.J.; Cociurovscaia, A.  
Deposited on : 2020-09-11  
Resolution : 1.76 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.76 Å.



Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	2466 (1.76-1.76)
Ramachandran outliers	138981	2437 (1.76-1.76)
Sidechain outliers	138945	2437 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	178	 85% 12%
1	B	178	 87% 12%

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 3162 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HTH-type transcriptional regulator RcdA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	173	1418	905	235	265	13	0	7	0
1	B	178	1446	918	248	267	13	0	4	0

- Molecule 2 is trimethylamine oxide (three-letter code: TMO) (formula:  $C_3H_9NO$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	5	3	1	1	0	0
2	B	1	10	6	2	2	0	1

- Molecule 3 is water.


<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
3	A	149	Total 149	O 149	0	0
3	B	134	Total 134	O 134	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.


- Molecule 1: HTH-type transcriptional regulator RcdA

Chain A:  85% 12%



- Molecule 1: HTH-type transcriptional regulator RcdA

Chain B:  87% 12%



## 4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	39.76Å 74.13Å 51.71Å 90.00° 95.19° 90.00°	Depositor
Resolution (Å)	42.29 – 1.76	Depositor
% Data completeness (in resolution range)	98.0 (42.29-1.76)	Depositor
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	PHENIX 1.9_1692	Depositor
R, $R_{free}$	0.158 , 0.203	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	3162	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TMO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.37	0/1464	0.52	0/1977
1	B	0.36	0/1480	0.53	0/1998
All	All	0.36	0/2944	0.52	0/3975

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1418	0	1426	17	0
1	B	1446	0	1453	17	0
2	A	5	0	9	0	0
2	B	10	0	18	0	0
3	A	149	0	0	5	0
3	B	134	0	0	4	0
All	All	3162	0	2906	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:58:GLU:HG3	3:A:314:HOH:O	1.79	0.81
1:A:139[B]:GLU:HG2	1:A:140:PRO:HD2	1.67	0.74
1:B:6:ASP:H	1:B:48:TYR:HE1	1.35	0.73
1:A:58:GLU:CG	3:A:314:HOH:O	2.40	0.66
1:B:3:ARG:NE	1:B:5:ASN:OD1	2.28	0.66
1:B:6:ASP:N	1:B:48:TYR:HE1	1.97	0.61
1:B:29[A]:VAL:HG23	1:B:52:ILE:HD11	1.89	0.54
1:B:162:LYS:NZ	3:B:308:HOH:O	2.44	0.51
1:B:9:ARG:NE	1:B:48:TYR:OH	2.39	0.49
1:A:14:ILE:HG23	1:A:59:ALA:HA	1.95	0.48
1:B:50:SER:OG	1:B:51:GLY:N	2.47	0.48
1:B:171:ARG:NH1	3:B:302:HOH:O	2.23	0.48
1:B:14:ILE:HG23	1:B:59:ALA:HA	1.94	0.48
1:A:139[B]:GLU:CD	1:A:141:GLY:H	2.17	0.47
1:B:53:ASP:HB3	1:B:118:LEU:HD12	1.97	0.47
1:A:110:TYR:HA	1:A:123[B]:MET:HE1	1.97	0.47
1:A:174:GLU:OE1	3:A:301:HOH:O	2.21	0.46
1:B:9:ARG:NH2	3:B:311:HOH:O	2.48	0.46
1:B:106[B]:MET:HG2	1:B:126:TRP:HH2	1.81	0.45
1:A:29:VAL:HG12	1:A:52:ILE:HD11	1.98	0.44
1:B:104:GLU:O	1:B:108:GLN:HG3	2.18	0.44
1:A:139[B]:GLU:HG2	1:A:140:PRO:CD	2.42	0.43
1:B:26:ILE:O	1:B:29[A]:VAL:HG22	2.18	0.43
1:B:65:GLU:OE1	3:B:301:HOH:O	2.21	0.43
1:A:62:SER:O	1:A:66:ILE:HG12	2.19	0.43
1:A:27:HIS:NE2	1:A:112:LEU:HD11	2.35	0.42
1:A:49:PHE:HE1	3:A:314:HOH:O	2.02	0.42
1:A:123[B]:MET:HG2	1:B:159:THR:HG21	2.01	0.42
1:A:92:ILE:HG21	1:A:153[B]:MET:HG3	2.00	0.41
1:B:106[B]:MET:O	1:B:109:LEU:HB3	2.20	0.41
1:A:10:ARG:NH1	3:A:314:HOH:O	2.54	0.41
1:A:149:PHE:CE2	1:A:153[A]:MET:HG3	2.55	0.41
1:A:26[B]:ILE:HD13	1:A:112:LEU:HD22	2.03	0.40

There are no symmetry-related clashes.



## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	178/178 (100%)	177 (99%)	1 (1%)	0	100	100
1	B	180/178 (101%)	177 (98%)	3 (2%)	0	100	100
All	All	358/356 (101%)	354 (99%)	4 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	155/152 (102%)	155 (100%)	0	100	100
1	B	155/152 (102%)	154 (99%)	1 (1%)	86	79
All	All	310/304 (102%)	309 (100%)	1 (0%)	92	89

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	3	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	TMO	B	201[A]	-	4,4,4	0.22	0	6,6,6	0.31	0
2	TMO	B	201[B]	-	4,4,4	0.24	0	6,6,6	0.24	0
2	TMO	A	201	-	4,4,4	0.18	0	6,6,6	0.25	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

### 6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.