



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 28, 2022 – 12:04 pm GMT

PDB ID : 7ACW  
Title : LID/HID (LMW SLP and HMW SLP interacting domains) from *C. difficile* (R7404 strain)  
Authors : Barwinska-Sendra, A.; Lanzoni-Mangutchi, P.; Basle, A.; Salgado, P.S.  
Deposited on : 2020-09-11  
Resolution : 1.50 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.27  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0267  
CCP4 : 7.1.010 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.27


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2936 (1.50-1.50)
Clashscore	141614	3144 (1.50-1.50)
Ramachandran outliers	138981	3066 (1.50-1.50)
Sidechain outliers	138945	3064 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	A	78	88% (green), 12% (grey)
1	C	78	83% (green), 6% (yellow), 10% (grey)
2	B	50	78% (green), 22% (grey)
2	D	50	70% (green), 1% (yellow), 28% (grey)

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 1816 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called S-layer protein.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
1	A	69	542	344	86	112	0	0	0
1	C	70	549	348	87	114	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	240	MET	-	initiating methionine	UNP Q9AEM2
C	240	MET	-	initiating methionine	UNP Q9AEM2

- Molecule 2 is a protein called S-layer protein.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	39	322	204	57	61	0	0	0
2	D	36	296	189	50	57	0	0	0

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	initiating methionine	UNP Q9AEM2
B	42	LEU	-	expression tag	UNP Q9AEM2
B	43	GLU	-	expression tag	UNP Q9AEM2
B	44	HIS	-	expression tag	UNP Q9AEM2
B	45	HIS	-	expression tag	UNP Q9AEM2
B	46	HIS	-	expression tag	UNP Q9AEM2
B	47	HIS	-	expression tag	UNP Q9AEM2
B	48	HIS	-	expression tag	UNP Q9AEM2
B	49	HIS	-	expression tag	UNP Q9AEM2
D	0	MET	-	initiating methionine	UNP Q9AEM2

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
D	42	LEU	-	expression tag	UNP Q9AEM2
D	43	GLU	-	expression tag	UNP Q9AEM2
D	44	HIS	-	expression tag	UNP Q9AEM2
D	45	HIS	-	expression tag	UNP Q9AEM2
D	46	HIS	-	expression tag	UNP Q9AEM2
D	47	HIS	-	expression tag	UNP Q9AEM2
D	48	HIS	-	expression tag	UNP Q9AEM2
D	49	HIS	-	expression tag	UNP Q9AEM2

- Molecule 3 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	31	Total O 31 31	0	0
3	B	23	Total O 23 23	0	0
3	C	45	Total O 45 45	0	0
3	D	8	Total O 8 8	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: S-layer protein

Chain A:  88% 12%





- Molecule 1: S-layer protein

Chain C:  83% 6% 10%




- Molecule 2: S-layer protein

Chain B:  78% 22%



- Molecule 2: S-layer protein

Chain D:  70% 28%



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	73.27Å 56.70Å 61.81Å 90.00° 122.71° 90.00°	Depositor
Resolution (Å)	52.01 – 1.50 52.01 – 1.50	Depositor EDS
% Data completeness (in resolution range)	96.5 (52.01-1.50) 94.8 (52.01-1.50)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.46 (at 1.50Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, $R_{free}$	0.164 , 0.204 0.178 , 0.215	Depositor DCC
$R_{free}$ test set	1687 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	19.2	Xtriage
Anisotropy	0.567	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	(Not available) , (Not available)	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	1816	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.26% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.45	0/550	0.63	0/743
1	C	0.47	0/557	0.60	0/753
2	B	0.41	0/329	0.62	0/445
2	D	0.39	0/301	0.55	0/406
All	All	0.44	0/1737	0.61	0/2347

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	542	0	531	0	0
1	C	549	0	538	4	0
2	B	322	0	319	0	0
2	D	296	0	301	2	0
3	A	31	0	0	0	0
3	B	23	0	0	0	0
3	C	45	0	0	0	0
3	D	8	0	0	0	0
All	All	1816	0	1689	4	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (4) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:254:ASP:OD2	1:C:301:LYS:NZ	2.23	0.72
1:C:244:THR:HG22	1:C:246:ALA:H	1.74	0.53
1:C:253:VAL:HG22	2:D:17:ALA:HB2	2.00	0.44
1:C:253:VAL:CG2	2:D:17:ALA:HB2	2.48	0.44

There are no symmetry-related clashes.

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	67/78 (86%)	67 (100%)	0	0	100	100
1	C	68/78 (87%)	68 (100%)	0	0	100	100
2	B	37/50 (74%)	37 (100%)	0	0	100	100
2	D	34/50 (68%)	34 (100%)	0	0	100	100
All	All	206/256 (80%)	206 (100%)	0	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	60/69 (87%)	60 (100%)	0	100	100
1	C	61/69 (88%)	61 (100%)	0	100	100
2	B	37/45 (82%)	37 (100%)	0	100	100
2	D	34/45 (76%)	34 (100%)	0	100	100
All	All	192/228 (84%)	192 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	35	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.