



# Full wwPDB X-ray Structure Validation Report ⓘ

May 22, 2020 – 12:09 am BST

PDB ID : 5AK6  
Title : ligand complex structure of soluble epoxide hydrolase  
Authors : Oster, L.; Tapani, S.; Xue, Y.; Kack, H.  
Deposited on : 2015-03-02  
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1479 (2.16-2.16)
Clashscore	141614	1585 (2.16-2.16)
Ramachandran outliers	138981	1560 (2.16-2.16)
Sidechain outliers	138945	1559 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	A	549	

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 4683 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called BIFUNCTIONAL EPOXIDE HYDROLASE 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	541	4286	2748	722	780	36	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP P34913

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	5	4	1	0	0
2	A	1	5	4	1	0	0

- Molecule 3 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			7	4	3		
3	A	1	Total	C	O	0	0
			7	4	3		


- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is 2-[(5-CHLORO-2-PYRIDYL)SULFANYL]ETHANOL (three-letter code:

YPN) (formula: C<sub>7</sub>H<sub>8</sub>ClNOS).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	Cl	N	O			S
5	A	1	Total	C	Cl	N	O	S	0	0
			11	7	1	1	1	1		
5	A	1	Total	C	Cl	N	O	S	0	0
			11	7	1	1	1	1		
5	A	1	Total	C	Cl	N	O	S	0	0
			11	7	1	1	1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	334	Total O 334 334	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: BIFUNCTIONAL EPOXIDE HYDROLASE 2



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	91.74Å 91.74Å 245.27Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	80.48 – 2.15 79.45 – 2.12	Depositor EDS
% Data completeness (in resolution range)	100.0 (80.48-2.15) 98.4 (79.45-2.12)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.64 (at 2.12Å)	Xtrriage
Refinement program	BUSTER 2.11.1	Depositor
R, $R_{free}$	0.204 , 0.248 0.356 , 0.374	Depositor DCC
$R_{free}$ test set	1774 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	25.7	Xtrriage
Anisotropy	0.235	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 44.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.82	EDS
Total number of atoms	4683	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.65% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, PEG, YPN, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.55	0/4390	0.73	1/5947 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	292	ASP	N-CA-C	-5.79	95.36	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4286	0	4288	30	13
2	A	10	0	0	0	0
3	A	14	0	20	3	0
4	A	6	0	8	0	0
5	A	33	0	24	2	0
6	A	334	0	0	4	8
All	All	4683	0	4340	32	16

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.



All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:331:PHE:HB3	1:A:341:VAL:HG22	1.52	0.91
1:A:41:PHE:HA	1:A:67:MET:HE1	1.57	0.86
1:A:384:GLN:HE22	5:A:1553:YPN:H5	1.54	0.72
1:A:422:VAL:HG11	1:A:428:LEU:HD22	1.78	0.65
1:A:309:CYS:SG	1:A:312:VAL:HG13	2.41	0.60
1:A:350:GLU:HG3	6:A:2204:HOH:O	2.03	0.57
1:A:266:GLY:HA3	1:A:335:ASP:HB3	1.84	0.57
1:A:41:PHE:HA	1:A:67:MET:CE	2.34	0.56
1:A:22:VAL:HG11	1:A:95:ALA:HB2	1.88	0.55
1:A:233:ASN:ND2	6:A:2133:HOH:O	2.40	0.55
3:A:1550:PEG:H31	6:A:2333:HOH:O	2.07	0.54
3:A:1551:PEG:H21	5:A:1555:YPN:N4	2.23	0.53
1:A:341:VAL:HG13	1:A:355:VAL:HG23	1.90	0.51
1:A:27:GLU:HA	1:A:32:LEU:HD12	1.94	0.50
1:A:341:VAL:HG13	1:A:355:VAL:CG2	2.42	0.49
1:A:69:GLU:HG3	1:A:73:LYS:HZ2	1.76	0.49
1:A:3:LEU:HD13	1:A:181:VAL:HG22	1.95	0.48
1:A:495:LYS:HD2	1:A:521:ASP:HA	1.96	0.48
1:A:83:PRO:HG2	1:A:86:PHE:HB2	1.95	0.48
1:A:484:ILE:HB	1:A:511:ILE:HG12	1.97	0.46
1:A:410:ARG:HD2	1:A:416:VAL:HG22	1.98	0.46
1:A:69:GLU:HG3	1:A:73:LYS:NZ	2.31	0.46
1:A:355:VAL:O	1:A:488:ALA:HA	2.16	0.46
1:A:168:PHE:O	1:A:172:THR:HG23	2.17	0.44
1:A:170:LEU:HD13	1:A:177:PRO:HD3	2.00	0.43
1:A:107:GLN:HG2	6:A:2053:HOH:O	2.17	0.43
1:A:122:LEU:HD22	1:A:169:LEU:HD22	2.00	0.42
1:A:15:ALA:HB1	1:A:99:ARG:HG2	2.02	0.42
1:A:370:SER:HB2	1:A:373:GLU:HG3	2.02	0.41
1:A:70:ASN:HA	1:A:73:LYS:HZ3	1.85	0.41
1:A:536:GLN:HB2	3:A:1550:PEG:H41	2.03	0.40
1:A:87:SER:HB3	1:A:90:GLU:HB2	2.03	0.40

All (16) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:68:GLU:OE1	1:A:72:ARG:NH2[8_435]	1.28	0.92
6:A:2093:HOH:O	6:A:2312:HOH:O[8_545]	1.64	0.56

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:327:SER:OG	6:A:2071:HOH:O[12_544]	1.65	0.55
1:A:239:HIS:O	1:A:241:TYR:N[12_544]	1.69	0.51
1:A:68:GLU:OE1	1:A:72:ARG:CZ[8_435]	1.73	0.47
1:A:239:HIS:N	1:A:241:TYR:O[12_544]	1.84	0.36
1:A:155:GLN:NE2	6:A:2290:HOH:O[12_544]	1.92	0.28
1:A:327:SER:N	6:A:2151:HOH:O[12_544]	1.94	0.26
1:A:237:MET:O	6:A:2140:HOH:O[12_544]	1.96	0.24
1:A:236:ASP:OD1	1:A:323:LYS:NZ[12_544]	2.04	0.16
1:A:256:GLY:N	1:A:324:LEU:O[12_544]	2.06	0.14
1:A:65:PRO:O	1:A:69:GLU:OE2[8_435]	2.08	0.12
1:A:236:ASP:O	1:A:323:LYS:CD[12_544]	2.10	0.10
6:A:2077:HOH:O	6:A:2205:HOH:O[12_544]	2.12	0.08
6:A:2134:HOH:O	6:A:2196:HOH:O[12_544]	2.15	0.05
1:A:324:LEU:O	6:A:2147:HOH:O[12_544]	2.15	0.05

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	537/549 (98%)	523 (97%)	13 (2%)	1 (0%)	47 46

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	268	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	468/474 (99%)	444 (95%)	24 (5%)	24	20

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	ARG
1	A	68	GLU
1	A	71	CYS
1	A	72	ARG
1	A	75	SER
1	A	81	CYS
1	A	110	LEU
1	A	159	VAL
1	A	181	VAL
1	A	216	VAL
1	A	224	THR
1	A	233	ASN
1	A	249	ARG
1	A	253	VAL
1	A	306	GLU
1	A	341	VAL
1	A	353	ARG
1	A	410	ARG
1	A	419	MET
1	A	428	LEU
1	A	448	GLN
1	A	460	ARG
1	A	508	GLU
1	A	547	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	138	GLN
1	A	233	ASN
1	A	384	GLN
1	A	452	GLN
1	A	513	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	YPN	A	1553	-	11,11,11	0.50	0	13,13,13	0.92	0
3	PEG	A	1550	-	6,6,6	0.71	0	5,5,5	1.08	0
2	SO4	A	1548	-	4,4,4	0.39	0	6,6,6	0.18	0
5	YPN	A	1554	-	11,11,11	0.34	0	13,13,13	0.76	0
4	GOL	A	1552	-	5,5,5	0.71	0	5,5,5	0.61	0
5	YPN	A	1555	-	11,11,11	0.75	0	13,13,13	0.87	0
3	PEG	A	1551	-	6,6,6	0.52	0	5,5,5	1.09	0
2	SO4	A	1549	-	4,4,4	0.41	0	6,6,6	0.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	YPN	A	1553	-	-	0/4/4/4	0/1/1/1
3	PEG	A	1550	-	-	3/4/4/4	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	YPN	A	1554	-	-	0/4/4/4	0/1/1/1
4	GOL	A	1552	-	-	1/4/4/4	-
5	YPN	A	1555	-	-	3/4/4/4	0/1/1/1
3	PEG	A	1551	-	-	4/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	1555	YPN	C2-C3-S8-C9
5	A	1555	YPN	N4-C3-S8-C9
5	A	1555	YPN	O11-C10-C9-S8
3	A	1551	PEG	O1-C1-C2-O2
3	A	1550	PEG	O1-C1-C2-O2
3	A	1551	PEG	O2-C3-C4-O4
3	A	1551	PEG	C1-C2-O2-C3
3	A	1550	PEG	C4-C3-O2-C2
3	A	1550	PEG	O2-C3-C4-O4
3	A	1551	PEG	C4-C3-O2-C2
4	A	1552	GOL	O1-C1-C2-O2

There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	1553	YPN	1	0
3	A	1550	PEG	2	0
5	A	1555	YPN	1	0
3	A	1551	PEG	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.