



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 18, 2023 – 07:13 pm BST

PDB ID : 8AK4
Title : Structure of the C-terminally truncated NAD⁺-dependent DNA ligase from the poly-extremophile *Deinococcus radiodurans*
Authors : Fernandes, A.; Williamson, A.K.; Matias, P.M.; Moe, E.
Deposited on : 2022-07-29
Resolution : 3.36 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.36 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1558 (3.42-3.30)
Clashscore	141614	1627 (3.42-3.30)
Ramachandran outliers	138981	1599 (3.42-3.30)
Sidechain outliers	138945	1598 (3.42-3.30)
RSRZ outliers	127900	1507 (3.42-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	600	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3932 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA ligase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	513	3926	2462	705	748	11	0	0	0

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Mn	0	0
			2	2		


- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	4	Total	Zn	0	0
			4	4		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA ligase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 2 21	Depositor
Cell constants a, b, c, α , β , γ	70.84Å 73.43Å 149.94Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	74.97 – 3.36 74.97 – 3.36	Depositor EDS
% Data completeness (in resolution range)	67.0 (74.97-3.36) 67.0 (74.97-3.36)	Depositor EDS
R_{merge}	0.22	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.82 (at 3.33Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.215 , 0.275 0.214 , 0.271	Depositor DCC
R_{free} test set	394 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	109.9	Xtrriage
Anisotropy	0.114	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 71.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	0.047 for k,h,-l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3932	wwPDB-VP
Average B, all atoms (Å ²)	103.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.07% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.24	0/4002	0.49	0/5433

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	438	ASN	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3926	0	3907	38	0
2	A	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	4	0	0	0	0
All	All	3932	0	3907	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:406:PRO:HG2	1:A:409:ALA:HB2	1.69	0.73
1:A:312:PHE:HA	1:A:317:PRO:HA	1.74	0.68
1:A:436:CYS:HB3	1:A:444:GLN:NE2	2.14	0.62
1:A:453:VAL:HG13	1:A:460:VAL:HB	1.81	0.61
1:A:436:CYS:SG	1:A:437:PRO:HD2	2.43	0.59
1:A:438:ASN:HB3	1:A:439:PRO:HD3	1.84	0.58
1:A:548:LEU:HD23	1:A:573:LEU:HD23	1.86	0.57
1:A:445:SER:O	1:A:449:ILE:HG12	2.08	0.53
1:A:193:ALA:HA	1:A:394:ILE:HD11	1.90	0.53
1:A:101:LEU:HG	1:A:273:ILE:HD11	1.92	0.52
1:A:138:TYR:HB2	1:A:179:LEU:HB3	1.91	0.52
1:A:541:PHE:HZ	1:A:555:ILE:HG22	1.75	0.52
1:A:561:LEU:HD22	1:A:565:ILE:HG21	1.92	0.51
1:A:418:CYS:HB3	1:A:421:CYS:SG	2.50	0.50
1:A:270:LEU:HD13	1:A:276:ALA:HA	1.94	0.50
1:A:436:CYS:HB3	1:A:444:GLN:HE22	1.77	0.49
1:A:521:ILE:HG13	1:A:534:ALA:HB1	1.94	0.49
1:A:518:TRP:HA	1:A:521:ILE:HD11	1.95	0.48
1:A:438:ASN:HD22	1:A:578:MET:HA	1.79	0.48
1:A:479:ASP:HB2	1:A:581:LEU:HD13	1.95	0.48
1:A:349:PRO:HG2	1:A:368:LEU:HB3	1.95	0.47
1:A:156:GLU:HG3	1:A:219:ARG:HH12	1.80	0.47
1:A:231:LEU:HD23	1:A:231:LEU:HA	1.75	0.46
1:A:438:ASN:CB	1:A:439:PRO:HD3	2.46	0.46
1:A:379:LEU:HD11	1:A:385:VAL:HG11	1.99	0.45
1:A:341:VAL:HG11	1:A:441:CYS:SG	2.57	0.44
1:A:358:ILE:O	1:A:360:GLY:N	2.51	0.44
1:A:87:THR:HB	1:A:182:ARG:NH2	2.33	0.43
1:A:350:LEU:HD11	1:A:365:LYS:HB3	2.00	0.43
1:A:512:SER:HA	1:A:515:LYS:HG3	1.99	0.43
1:A:385:VAL:HG21	1:A:397:ILE:HD11	2.01	0.43
1:A:279:TYR:O	1:A:283:MET:HG2	2.18	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:167:ASP:O	1:A:262:PRO:HD3	2.20	0.42
1:A:368:LEU:HD12	1:A:368:LEU:HA	1.87	0.42
1:A:329:GLU:HG2	1:A:388:ARG:HB3	2.02	0.42
1:A:301:LEU:O	1:A:307:GLN:NE2	2.48	0.41
1:A:541:PHE:CZ	1:A:555:ILE:HG22	2.55	0.41
1:A:332:THR:HB	1:A:356:ARG:HG3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	511/600 (85%)	465 (91%)	40 (8%)	6 (1%)	13 44

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	437	PRO
1	A	166	PRO
1	A	438	ASN
1	A	359	GLU
1	A	369	HIS
1	A	360	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	407/478 (85%)	396 (97%)	11 (3%)	44 72

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	207	LEU
1	A	231	LEU
1	A	236	TYR
1	A	323	TYR
1	A	353	LEU
1	A	370	ASN
1	A	403	ASP
1	A	416	THR
1	A	500	LYS
1	A	521	ILE
1	A	570	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	513/600 (85%)	0.39	24 (4%) 31 34	71, 100, 140, 189	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	387	VAL	3.4
1	A	388	ARG	3.1
1	A	396	GLN	3.1
1	A	386	LEU	3.0
1	A	531	GLN	2.7
1	A	179	LEU	2.6
1	A	379	LEU	2.6
1	A	138	TYR	2.5
1	A	593	GLN	2.4
1	A	542	GLY	2.4
1	A	397	ILE	2.3
1	A	334	LEU	2.3
1	A	137	TYR	2.3
1	A	261	PHE	2.3
1	A	516	PRO	2.2
1	A	81	GLU	2.2
1	A	563	GLY	2.2
1	A	181	VAL	2.1
1	A	218	LEU	2.1
1	A	537	LEU	2.1
1	A	394	ILE	2.0
1	A	462	GLY	2.0
1	A	492	ALA	2.0
1	A	385	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MN	A	801	1/1	0.95	0.26	59,59,59,59	0
3	ZN	A	806	1/1	0.97	0.25	67,67,67,67	0
3	ZN	A	803	1/1	0.98	0.30	59,59,59,59	0
3	ZN	A	805	1/1	0.98	0.36	30,30,30,30	1
2	MN	A	802	1/1	0.98	0.31	61,61,61,61	0
3	ZN	A	804	1/1	0.99	0.29	48,48,48,48	0

6.5 Other polymers [i](#)

There are no such residues in this entry.