



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 25, 2020 – 05:32 PM BST

PDB ID : 5AN7  
Title : Structure of the engineered retro-aldolase RA95.5-8F with a bound 1,3-diketone inhibitor  
Authors : Obexer, R.; Mittl, P.R.E.; Hilvert, D.  
Deposited on : 2015-09-04  
Resolution : 1.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1619 (1.14-1.06)
Clashscore	141614	1671 (1.14-1.06)
Ramachandran outliers	138981	1615 (1.14-1.06)
Sidechain outliers	138945	1613 (1.14-1.06)
RSRZ outliers	127900	1588 (1.14-1.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	258	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	LLK	A	5001	-	X	-	-

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4799 atoms, of which 2286 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RA95.5-8F.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	244	4474	1413	2272	364	414	11	0	28	1

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	P		
2	A	1	5	4	1	0	0
2	A	1	5	4	1	0	0

- Molecule 3 is (2E)-1-(6-methoxynaphthalen-2-yl)but-2-en-1-one (three-letter code: LLK) (formula: C<sub>15</sub>H<sub>14</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
3	A	1	31	15	14	2	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	284	284	284	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: RA95.5-8F



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	75.11Å 84.43Å 37.64Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.60 – 1.10 37.64 – 1.17	Depositor EDS
% Data completeness (in resolution range)	89.2 (37.60-1.10) 99.0 (37.64-1.17)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.13 (at 1.17Å)	Xtrriage
Refinement program	SHELXL	Depositor
R, $R_{free}$	0.136 , 0.188 0.146 , 0.182	Depositor DCC
$R_{free}$ test set	3884 reflections (4.81%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	10.7	Xtrriage
Anisotropy	0.228	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.43 , 90.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	4799	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.56% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: LLK, MHO, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.72	1/2226 (0.0%)	1.43	46/3000 (1.5%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1248	GLU	C-N	10.21	1.51	1.33

All (46) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1036	ARG	NE-CZ-NH1	10.29	125.44	120.30
1	A	1150[A]	ARG	NE-CZ-NH1	9.73	125.16	120.30
1	A	1150[B]	ARG	NE-CZ-NH1	9.73	125.16	120.30
1	A	1195	ARG	CD-NE-CZ	9.67	137.14	123.60
1	A	1171	ARG	NE-CZ-NH1	8.84	124.72	120.30
1	A	1093	TYR	CB-CG-CD1	8.72	126.23	121.00
1	A	1223	ARG	CD-NE-CZ	8.00	134.80	123.60
1	A	1223	ARG	NE-CZ-NH1	7.73	124.17	120.30
1	A	1036	ARG	NE-CZ-NH2	-7.60	116.50	120.30
1	A	1023	HIS	CA-CB-CG	-7.40	101.02	113.60
1	A	1026	ARG	NE-CZ-NH1	7.38	123.99	120.30
1	A	1248	GLU	O-C-N	7.37	135.72	123.20
1	A	1065	ASP	CB-CG-OD1	7.29	124.86	118.30
1	A	1188[A]	GLU	OE1-CD-OE2	-7.18	114.69	123.30
1	A	1188[B]	GLU	OE1-CD-OE2	-7.18	114.69	123.30
1	A	1150[A]	ARG	NE-CZ-NH2	-6.69	116.95	120.30
1	A	1150[B]	ARG	NE-CZ-NH2	-6.69	116.95	120.30
1	A	1076	TYR	CB-CG-CD1	6.39	124.83	121.00
1	A	1054	ARG	CD-NE-CZ	6.32	132.45	123.60
1	A	1238	ARG	NE-CZ-NH2	6.15	123.38	120.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1188[A]	GLU	N-CA-CB	6.05	121.50	110.60
1	A	1188[B]	GLU	N-CA-CB	6.05	121.50	110.60
1	A	1223	ARG	NH1-CZ-NH2	-5.90	112.91	119.40
1	A	1092[A]	SER	N-CA-CB	5.81	119.22	110.50
1	A	1092[B]	SER	N-CA-CB	5.81	119.22	110.50
1	A	1248	GLU	CA-C-N	-5.76	104.67	116.20
1	A	1148	TYR	CB-CG-CD2	5.74	124.45	121.00
1	A	1052	TYR	CG-CD2-CE2	5.64	125.81	121.30
1	A	1152	TYR	CB-CG-CD2	5.59	124.35	121.00
1	A	1018	ARG	CD-NE-CZ	5.50	131.29	123.60
1	A	1086	GLU	OE1-CD-OE2	-5.44	116.78	123.30
1	A	1238	ARG	CG-CD-NE	-5.39	100.47	111.80
1	A	1069	TYR	CB-CG-CD1	5.37	124.22	121.00
1	A	1092[A]	SER	CB-CA-C	-5.36	99.92	110.10
1	A	1092[B]	SER	CB-CA-C	-5.36	99.92	110.10
1	A	1207	LYS	CD-CE-NZ	5.35	124.00	111.70
1	A	1175	ARG	CD-NE-CZ	5.33	131.06	123.60
1	A	1162	ASP	CB-CG-OD1	5.29	123.06	118.30
1	A	1223	ARG	NE-CZ-NH2	5.23	122.91	120.30
1	A	1025	SER	N-CA-CB	-5.21	102.69	110.50
1	A	1055	LYS	CA-CB-CG	5.17	124.78	113.40
1	A	1022[A]	VAL	C-N-CA	5.14	134.56	121.70
1	A	1022[B]	VAL	C-N-CA	5.14	134.56	121.70
1	A	1243	ILE	CB-CG1-CD1	5.13	128.28	113.90
1	A	1076	TYR	CB-CG-CD2	-5.07	117.96	121.00
1	A	1093	TYR	CB-CG-CD2	-5.02	117.99	121.00

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2202	2272	2248	41	0
2	A	10	0	0	0	0
3	A	17	14	13	6	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	284	0	0	24	0
All	All	2513	2286	2261	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1188[A]:GLU:CD	4:A:2226:HOH:O	1.82	1.18
3:A:5001:LLK:C13	4:A:2122:HOH:O	1.95	1.14
1:A:1051[A]:TYR:OH	4:A:2122:HOH:O	1.65	1.13
1:A:1188[A]:GLU:CG	4:A:2226:HOH:O	1.94	1.13
1:A:1188[A]:GLU:OE1	4:A:2226:HOH:O	1.69	1.09
1:A:1199[A]:SER:OG	4:A:2240:HOH:O	1.68	1.08
1:A:1188[A]:GLU:HB3	4:A:2226:HOH:O	1.51	1.08
1:A:1188[A]:GLU:CB	4:A:2226:HOH:O	2.02	1.07
1:A:1071[A]:LYS:HG3	4:A:2136:HOH:O	1.50	1.07
1:A:1051[B]:TYR:OH	3:A:5001:LLK:O1	1.78	1.00
1:A:1051[B]:TYR:HE1	1:A:1053[B]:LEU:HD22	1.25	0.96
1:A:1146[B]:LEU:HD12	1:A:1158[B]:ILE:HD11	1.49	0.94
1:A:1083[A]:LYS:NZ	4:A:2122:HOH:O	1.98	0.91
1:A:1051[B]:TYR:CE1	1:A:1053[B]:LEU:HD22	2.06	0.89
1:A:1237:MHO:HG2	4:A:2123:HOH:O	1.90	0.70
1:A:1184:PHE:O	3:A:5001:LLK:H2	1.97	0.65
1:A:1092[A]:SER:OG	4:A:2157:HOH:O	2.01	0.64
1:A:1098[C]:LYS:HE3	4:A:2162:HOH:O	1.97	0.64
1:A:1006:LYS:HE3	4:A:2005:HOH:O	1.99	0.61
1:A:1237:MHO:HG3	4:A:2121:HOH:O	2.00	0.60
1:A:1067[B]:ILE:HD13	1:A:1095[B]:MET:SD	2.42	0.59
1:A:1055:LYS:HE2	4:A:2130:HOH:O	2.02	0.58
1:A:1067[B]:ILE:CD1	1:A:1095[B]:MET:SD	2.92	0.57
1:A:1107[B]:ILE:HD11	4:A:2094:HOH:O	2.04	0.57
1:A:1051[B]:TYR:O	1:A:1051[B]:TYR:HD1	1.88	0.56
3:A:5001:LLK:H14	4:A:2150:HOH:O	2.08	0.54
1:A:1051[B]:TYR:C	1:A:1051[B]:TYR:CD1	2.81	0.53
1:A:1195:ARG:HD3	4:A:2239:HOH:O	2.10	0.52
1:A:1182:MET:HA	1:A:1188[A]:GLU:O	2.10	0.51
1:A:1051[B]:TYR:O	1:A:1051[B]:TYR:CD1	2.64	0.51
3:A:5001:LLK:H7	4:A:2151:HOH:O	2.10	0.50
1:A:1026:ARG:NH2	1:A:1030[B]:ILE:HD13	2.28	0.48

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1098[B]:LYS:HD3	4:A:2162:HOH:O	2.14	0.47
1:A:1168[B]:ILE:HD13	4:A:2003:HOH:O	2.15	0.47
1:A:1066:PRO:HG2	1:A:1095[B]:MET:SD	2.55	0.46
1:A:1062:VAL:HG12	1:A:1062:VAL:O	2.15	0.46
1:A:1217:ASN:HB3	4:A:2259:HOH:O	2.14	0.46
1:A:1063:GLU:N	1:A:1063:GLU:OE1	2.50	0.45
1:A:1208[B]:VAL:O	1:A:1208[B]:VAL:HG13	2.18	0.44
1:A:1083[B]:LYS:NZ	4:A:2150:HOH:O	2.50	0.43
1:A:1219:ILE:HG23	1:A:1230:PHE:CZ	2.53	0.43
1:A:1207:LYS:H	1:A:1228:ASN:HD22	1.67	0.42
1:A:1056:SER:HB2	3:A:5001:LLK:C10	2.51	0.41
1:A:1083[B]:LYS:HA	1:A:1110:ASN:O	2.20	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	267/258 (104%)	265 (99%)	2 (1%)	0	<b>100</b>   <b>100</b>

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	249/234 (106%)	241 (97%)	8 (3%)	39 7

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1023	HIS
1	A	1025	SER
1	A	1054	ARG
1	A	1092[A]	SER
1	A	1092[B]	SER
1	A	1202	PRO
1	A	1217	ASN
1	A	1247	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1041	ASN
1	A	1228	ASN
1	A	1239	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	MHO	A	1237	1	7,8,9	1.67	1 (14%)	4,9,11	6.01	3 (75%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	MHO	A	1237	1	-	3/6/7/9	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1237	MHO	OD1-SD	3.96	1.61	1.50

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1237	MHO	OD1-SD-CE	7.31	121.06	106.25
1	A	1237	MHO	OD1-SD-CG	-7.18	86.24	106.03
1	A	1237	MHO	CE-SD-CG	6.05	111.47	97.71

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	1237	MHO	CB-CG-SD-CE
1	A	1237	MHO	CB-CG-SD-OD1
1	A	1237	MHO	CA-CB-CG-SD

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	1237	MHO	2	0

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	PO4	A	4002	-	4,4,4	1.10	0	6,6,6	1.05	0
2	PO4	A	4003	-	4,4,4	1.23	0	6,6,6	0.73	0
3	LLK	A	5001	-	18,18,18	2.18	6 (33%)	24,24,24	3.23	13 (54%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	LLK	A	5001	-	-	4/9/9/9	0/2/2/2

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	5001	LLK	C8-C11	-5.03	1.41	1.49
3	A	5001	LLK	C12-C13	4.72	1.48	1.30
3	A	5001	LLK	C1-C2	3.07	1.43	1.36
3	A	5001	LLK	C10-C9	2.65	1.42	1.36
3	A	5001	LLK	C6-C5	-2.57	1.36	1.42
3	A	5001	LLK	C4-C3	2.46	1.41	1.37

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	5001	LLK	C13-C12-C11	8.16	134.89	123.00
3	A	5001	LLK	C14-C13-C12	-7.34	110.86	125.34
3	A	5001	LLK	C10-C5-C4	-4.74	113.65	122.02
3	A	5001	LLK	C8-C7-C6	-4.09	114.92	121.24
3	A	5001	LLK	C4-C5-C6	4.04	124.71	118.96
3	A	5001	LLK	O3-C3-C4	-3.57	114.69	124.43
3	A	5001	LLK	C7-C6-C5	3.48	123.92	118.96
3	A	5001	LLK	C9-C8-C11	-2.92	113.96	120.88
3	A	5001	LLK	C1-C6-C7	-2.79	117.10	122.02

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	5001	LLK	C9-C10-C5	-2.69	116.66	120.82
3	A	5001	LLK	C3-C4-C5	-2.56	116.01	119.70
3	A	5001	LLK	C9-C8-C7	2.41	122.10	119.23
3	A	5001	LLK	C8-C11-C12	2.20	121.70	119.23

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	5001	LLK	C11-C12-C13-C14
3	A	5001	LLK	C4-C3-O3-C31
3	A	5001	LLK	C2-C3-O3-C31
3	A	5001	LLK	C8-C11-C12-C13

There are no ring outliers.

1 monomer is involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	5001	LLK	6	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	243/258 (94%)	-0.00	4 (1%) 72 68	8, 13, 38, 71	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1062	VAL	12.7
1	A	1216	PRO	8.5
1	A	1217	ASN	3.1
1	A	1023	HIS	2.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	MHO	A	1237	9/10	0.91	0.13	8,29,46,46	1

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	LLK	A	5001	17/17	0.90	0.15	15,31,78,78	31
2	PO4	A	4003	5/5	0.99	0.05	15,16,27,28	0
2	PO4	A	4002	5/5	0.99	0.06	12,14,18,20	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.