



wwPDB X-ray Structure Validation Summary Report ⓘ

May 21, 2020 – 03:10 pm BST

PDB ID : 1AQC
Title : X11 PTB DOMAIN-10MER PEPTIDE COMPLEX
Authors : Lee, C.-H.; Zhang, Z.; Kuriyan, J.
Deposited on : 1997-07-28
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	172	41% (green), 32% (yellow), 24% (grey), 0% (orange), 0% (red)
1	B	172	38% (green), 30% (yellow), 29% (grey), 0% (orange), 0% (red)
2	C	10	60% (green), 20% (yellow), 20% (orange), 0% (red)
2	D	10	70% (green), 30% (yellow), 0% (orange), 0% (red)

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2351 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called X11.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	130	1016	642	174	190	1	9	0	0	0
1	B	122	944	599	166	171	1	7	0	0	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	354	MSE	MET	CONFLICT	UNP Q02410
A	355	MSE	MET	CONFLICT	UNP Q02410
A	366	MSE	MET	CONFLICT	UNP Q02410
A	385	MSE	MET	CONFLICT	UNP Q02410
A	408	MSE	MET	CONFLICT	UNP Q02410
A	409	MSE	MET	CONFLICT	UNP Q02410
A	429	MSE	MET	CONFLICT	UNP Q02410
A	458	MSE	MET	CONFLICT	UNP Q02410
B	354	MSE	MET	CONFLICT	UNP Q02410
B	355	MSE	MET	CONFLICT	UNP Q02410
B	366	MSE	MET	CONFLICT	UNP Q02410
B	385	MSE	MET	CONFLICT	UNP Q02410
B	408	MSE	MET	CONFLICT	UNP Q02410
B	409	MSE	MET	CONFLICT	UNP Q02410
B	429	MSE	MET	CONFLICT	UNP Q02410
B	458	MSE	MET	CONFLICT	UNP Q02410

- Molecule 2 is a protein called PEPTIDE.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	10	90	62	12	16	0	0	0
2	D	10	86	59	11	16	0	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	102	Total 102	O 102	0	0
3	C	4	Total 4	O 4	0	0
3	B	92	Total 92	O 92	0	0
3	D	17	Total 17	O 17	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	74.40Å 74.40Å 157.10Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	6.00 – 2.30 28.08 – 2.30	Depositor EDS
% Data completeness (in resolution range)	69.0 (6.00-2.30) 90.9 (28.08-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.92 (at 2.31Å)	Xtrriage
Refinement program	X-PLOR 3.8	Depositor
R, R_{free}	0.210 , 0.303 0.260 , 0.344	Depositor DCC
R_{free} test set	1822 reflections (9.84%)	wwPDB-VP
Wilson B-factor (Å ²)	37.1	Xtrriage
Anisotropy	0.109	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 60.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	2351	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.66	0/1019	0.78	0/1358
1	B	0.67	1/944 (0.1%)	0.77	1/1256 (0.1%)
2	C	0.65	0/94	0.74	0/126
2	D	0.67	0/90	0.59	0/122
All	All	0.67	1/2147 (0.0%)	0.77	1/2862 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	458	MSE	CG-SE	-5.15	1.77	1.95

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	494	PRO	N-CA-CB	5.09	109.41	103.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1016	0	1020	50	0
1	B	944	0	959	50	0
2	C	90	0	77	4	0
2	D	86	0	66	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	102	0	0	5	0
3	B	92	0	0	9	0
3	C	4	0	0	0	0
3	D	17	0	0	0	0
All	All	2351	0	2122	97	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 23.

The worst 5 of 97 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:359:GLU:O	1:B:363:ARG:HG2	1.83	0.78
1:B:391:PHE:CZ	1:B:393:LEU:HD22	2.22	0.74
2:C:8:THR:HG22	2:C:12:PHE:HE2	1.55	0.72
1:B:391:PHE:HZ	1:B:393:LEU:HD22	1.54	0.71
1:A:393:LEU:HG	1:A:395:GLN:H	1.56	0.69

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	124/172 (72%)	118 (95%)	5 (4%)	1 (1%)	19	23
1	B	114/172 (66%)	106 (93%)	6 (5%)	2 (2%)	8	7
2	C	8/10 (80%)	6 (75%)	2 (25%)	0	100	100
2	D	8/10 (80%)	8 (100%)	0	0	100	100
All	All	254/364 (70%)	238 (94%)	13 (5%)	3 (1%)	13	14

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	432	ARG
1	B	413	LEU
1	B	432	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	109/140 (78%)	104 (95%)	5 (5%)	27	38
1	B	99/140 (71%)	90 (91%)	9 (9%)	9	11
2	C	9/9 (100%)	7 (78%)	2 (22%)	1	1
2	D	8/9 (89%)	8 (100%)	0	100	100
All	All	225/298 (76%)	209 (93%)	16 (7%)	14	19

5 of 16 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	343	LEU
1	B	358	GLN
1	B	422	ILE
2	C	9	TYR
1	B	424	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	424	ASN
1	B	473	GLN
1	A	477	GLN
1	A	395	GLN
1	A	469	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.