



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 24, 2024 – 08:13 AM EDT

PDB ID : 6ARJ
Title : Crystal structure of CARM1 with EPZ022302 and SAH
Authors : Boriack-Sjodin, P.A.; Jin, L.
Deposited on : 2017-08-22
Resolution : 1.92 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 2.37.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 1.92 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	7937 (1.94-1.90)
Clashscore	141614	8644 (1.94-1.90)
Ramachandran outliers	138981	8530 (1.94-1.90)
Sidechain outliers	138945	8530 (1.94-1.90)
RSRZ outliers	127900	7793 (1.94-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	349	 92% 7%
1	B	349	 85% 13%
1	C	349	 90% 8%
1	D	349	 91% 7%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 12491 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone-arginine methyltransferase CARM1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	344	2819	1816	468	519	16	0	8	0
1	B	344	2812	1811	468	517	16	0	7	0
1	C	344	2778	1791	461	511	15	0	2	0
1	D	343	2765	1784	456	510	15	0	2	0

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	131	SER	-	expression tag	UNP Q86X55
A	132	ILE	-	expression tag	UNP Q86X55
A	133	ALA	-	expression tag	UNP Q86X55
B	131	SER	-	expression tag	UNP Q86X55
B	132	ILE	-	expression tag	UNP Q86X55
B	133	ALA	-	expression tag	UNP Q86X55
C	131	SER	-	expression tag	UNP Q86X55
C	132	ILE	-	expression tag	UNP Q86X55
C	133	ALA	-	expression tag	UNP Q86X55
D	131	SER	-	expression tag	UNP Q86X55
D	132	ILE	-	expression tag	UNP Q86X55
D	133	ALA	-	expression tag	UNP Q86X55

- Molecule 2 is S-ADENOSYL-L-HOMOCYSTEINE (three-letter code: SAH) (formula: C₁₄H₂₀N₆O₅S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			26	14	6	5	1		
2	B	1	Total	C	N	O	S	0	0
			26	14	6	5	1		
2	C	1	Total	C	N	O	S	0	0
			26	14	6	5	1		
2	D	1	Total	C	N	O	S	0	0
			26	14	6	5	1		

- Molecule 3 is methyl 2-[2-{2-chloro-5-[(2R)-2-hydroxy-3-(methylamino)propoxy]phenyl}-6-(3,5-dimethyl-1,2-oxazol-4-yl)-5-methylpyrimidin-4-yl]-2,7-diazaspiro[3.5]nonane-7-carboxylate (three-letter code: BW4) (formula: C₂₉H₃₇ClN₆O₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	Cl	N			O
3	A	1	Total 82	C 58	Cl 2	N 12	O 10	0	1
3	B	1	Total 82	C 58	Cl 2	N 12	O 10	0	1
3	C	1	Total 82	C 58	Cl 2	N 12	O 10	0	1
3	D	1	Total 82	C 58	Cl 2	N 12	O 10	0	1

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	1	Total 6	O 3	0	0
4	B	1	Total 6	O 3	0	0
4	C	1	Total 6	O 3	0	0
4	D	1	Total 6	O 3	0	0
4	D	1	Total 6	O 3	0	0

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	C	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	230	Total	O	0	0
			230	230		
6	B	245	Total	O	0	0
			245	245		
6	C	188	Total	O	0	0
			188	188		
6	D	187	Total	O	0	0
			187	187		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Histone-arginine methyltransferase CARM1

Chain A: 





- Molecule 1: Histone-arginine methyltransferase CARM1

Chain B: 




- Molecule 1: Histone-arginine methyltransferase CARM1

Chain C: 



- Molecule 1: Histone-arginine methyltransferase CARM1

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	75.16Å 98.80Å 208.29Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.11 – 1.92 46.07 – 1.92	Depositor EDS
% Data completeness (in resolution range)	99.8 (48.11-1.92) 99.8 (46.07-1.92)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.00 (at 1.92Å)	Xtrriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.175 , 0.212 0.175 , 0.211	Depositor DCC
R_{free} test set	5909 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	24.3	Xtrriage
Anisotropy	0.046	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 44.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	12491	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 34.41 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.8537e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BW4, SAH, SO4, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.53	0/2899	0.66	1/3925 (0.0%)
1	B	0.56	0/2888	0.67	0/3910
1	C	0.48	0/2848	0.62	0/3858
1	D	0.51	0/2835	0.62	0/3841
All	All	0.52	0/11470	0.64	1/15534 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	247	LEU	CA-CB-CG	5.04	126.90	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	464[B]	SER	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2819	0	2765	10	0
1	B	2812	0	2763	29	0
1	C	2778	0	2726	17	0
1	D	2765	0	2711	13	0
2	A	26	0	19	0	0
2	B	26	0	19	0	0
2	C	26	0	19	0	0
2	D	26	0	19	0	0
3	A	82	0	0	2	0
3	B	82	0	0	2	0
3	C	82	0	0	1	0
3	D	82	0	0	0	0
4	A	6	0	8	0	0
4	B	6	0	8	0	0
4	C	6	0	8	0	0
4	D	12	0	16	1	0
5	C	5	0	0	0	0
6	A	230	0	0	5	0
6	B	245	0	0	7	0
6	C	188	0	0	5	0
6	D	187	0	0	4	0
All	All	12491	0	11081	74	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 74 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:502[B]:BW4: CBD	3:A:502[B]:BW4: CAY	2.25	1.14
3:A:502[A]:BW4: CAW	3:A:502[A]:BW4: CBD	2.29	1.11
1:A:341[B]:ASP: OD1	6:A:601:HOH: O	1.75	1.03
1:C:424:SER:HB2	6:C:737:HOH: O	1.70	0.90
1:A:327:ARG:HD2	6:C:606:HOH: O	1.82	0.79

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	350/349 (100%)	341 (97%)	8 (2%)	1 (0%)	41	31
1	B	349/349 (100%)	339 (97%)	9 (3%)	1 (0%)	41	31
1	C	344/349 (99%)	334 (97%)	9 (3%)	1 (0%)	41	31
1	D	343/349 (98%)	333 (97%)	10 (3%)	0	100	100
All	All	1386/1396 (99%)	1347 (97%)	36 (3%)	3 (0%)	47	38

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	266	GLU
1	B	266	GLU
1	C	266	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	308/303 (102%)	301 (98%)	7 (2%)	50	43
1	B	307/303 (101%)	301 (98%)	6 (2%)	55	49
1	C	302/303 (100%)	291 (96%)	11 (4%)	35	24
1	D	301/303 (99%)	295 (98%)	6 (2%)	55	49

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	1218/1212 (100%)	1188 (98%)	30 (2%)	47 39

5 of 30 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	140	ARG
1	D	257	GLU
1	C	294	LEU
1	D	354	LYS
1	D	140	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 31 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	161	ASN
1	D	306	GLN
1	C	180	HIS
1	D	455	GLN
1	D	178	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

18 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SAH	B	501	-	23,28,28	0.78	0	22,40,40	1.57	4 (18%)
4	GOL	A	503	-	5,5,5	0.26	0	5,5,5	0.34	0
2	SAH	C	501	-	23,28,28	1.12	3 (13%)	22,40,40	1.37	3 (13%)
3	BW4	C	502[A]	-	40,45,45	2.79	6 (15%)	48,66,66	2.45	15 (31%)
5	SO4	C	504	-	4,4,4	0.44	0	6,6,6	0.12	0
3	BW4	A	502[B]	-	40,45,45	2.64	6 (15%)	48,66,66	2.43	15 (31%)
3	BW4	D	502[B]	-	40,45,45	2.70	5 (12%)	48,66,66	2.40	15 (31%)
3	BW4	B	502[B]	-	40,45,45	2.49	5 (12%)	48,66,66	2.62	15 (31%)
4	GOL	C	503	-	5,5,5	0.23	0	5,5,5	0.33	0
4	GOL	D	503	-	5,5,5	0.19	0	5,5,5	0.50	0
3	BW4	A	502[A]	-	40,45,45	2.59	5 (12%)	48,66,66	2.46	13 (27%)
3	BW4	D	502[A]	-	40,45,45	2.68	5 (12%)	48,66,66	2.39	15 (31%)
3	BW4	B	502[A]	-	40,45,45	2.54	5 (12%)	48,66,66	2.60	15 (31%)
3	BW4	C	502[B]	-	40,45,45	2.79	6 (15%)	48,66,66	2.43	15 (31%)
4	GOL	B	503	-	5,5,5	0.22	0	5,5,5	0.43	0
4	GOL	D	504	-	5,5,5	0.27	0	5,5,5	0.34	0
2	SAH	A	501	-	23,28,28	0.95	2 (8%)	22,40,40	1.69	4 (18%)
2	SAH	D	501	-	23,28,28	1.01	2 (8%)	22,40,40	1.55	4 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SAH	B	501	-	-	0/11/31/31	0/3/3/3
4	GOL	A	503	-	-	0/4/4/4	-
2	SAH	C	501	-	-	0/11/31/31	0/3/3/3
3	BW4	C	502[A]	-	-	2/18/48/48	0/5/5/5
3	BW4	A	502[B]	-	-	2/18/48/48	0/5/5/5
3	BW4	D	502[B]	-	-	2/18/48/48	0/5/5/5
3	BW4	B	502[B]	-	-	2/18/48/48	0/5/5/5
4	GOL	C	503	-	-	2/4/4/4	-
4	GOL	D	503	-	-	4/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BW4	A	502[A]	-	-	2/18/48/48	0/5/5/5
3	BW4	D	502[A]	-	-	0/18/48/48	0/5/5/5
3	BW4	B	502[A]	-	-	2/18/48/48	0/5/5/5
3	BW4	C	502[B]	-	-	2/18/48/48	0/5/5/5
4	GOL	B	503	-	-	2/4/4/4	-
4	GOL	D	504	-	-	0/4/4/4	-
2	SAH	A	501	-	-	1/11/31/31	0/3/3/3
2	SAH	D	501	-	-	0/11/31/31	0/3/3/3

The worst 5 of 50 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	502[B]	BW4	CBI-C4	-14.88	1.34	1.50
3	C	502[A]	BW4	CBI-C4	-14.81	1.34	1.50
3	A	502[B]	BW4	CBI-C4	-13.68	1.35	1.50
3	D	502[B]	BW4	CBI-C4	-13.62	1.36	1.50
3	D	502[A]	BW4	CBI-C4	-13.52	1.36	1.50

The worst 5 of 133 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	502[B]	BW4	OBC-CBA-NAX	11.11	120.55	111.61
3	B	502[A]	BW4	OBC-CBA-NAX	11.04	120.50	111.61
3	D	502[B]	BW4	OBC-CBA-NAX	10.75	120.26	111.61
3	D	502[A]	BW4	OBC-CBA-NAX	10.66	120.19	111.61
3	A	502[A]	BW4	OBC-CBA-NAX	10.60	120.14	111.61

There are no chirality outliers.

5 of 23 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	502[A]	BW4	OBB-CBA-OBC-CBD
3	A	502[A]	BW4	NAX-CBA-OBC-CBD
3	A	502[B]	BW4	OBB-CBA-OBC-CBD
3	A	502[B]	BW4	NAX-CBA-OBC-CBD
3	C	502[A]	BW4	OBB-CBA-OBC-CBD



There are no ring outliers.



6 monomers are involved in 6 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	502[A]	BW4	1	0
3	A	502[B]	BW4	1	0
3	B	502[B]	BW4	1	0
3	A	502[A]	BW4	1	0
3	B	502[A]	BW4	1	0
4	D	504	GOL	1	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


















5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	344/349 (98%)	-0.35	2 (0%) 89 90	14, 22, 33, 58	0
1	B	344/349 (98%)	-0.36	3 (0%) 84 85	13, 21, 33, 60	0
1	C	344/349 (98%)	-0.32	2 (0%) 89 90	20, 27, 39, 77	0
1	D	343/349 (98%)	-0.29	1 (0%) 94 94	20, 28, 40, 56	0
All	All	1375/1396 (98%)	-0.33	8 (0%) 89 90	13, 25, 38, 77	0

The worst 5 of 8 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	134	ARG	4.0
1	B	134	ARG	3.5
1	C	346	ARG	2.5
1	B	427	PHE	2.5
1	B	304[A]	MET	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	GOL	D	504	6/6	0.80	0.22	50,51,54,56	0
4	GOL	B	503	6/6	0.84	0.17	56,58,60,61	0
4	GOL	A	503	6/6	0.84	0.11	47,51,54,59	0
4	GOL	D	503	6/6	0.85	0.17	60,62,64,68	0
4	GOL	C	503	6/6	0.85	0.15	48,51,52,58	0
5	SO4	C	504	5/5	0.86	0.27	82,83,87,87	0
3	BW4	D	502[A]	41/41	0.95	0.10	21,24,41,44	41
3	BW4	D	502[B]	41/41	0.95	0.10	21,25,44,46	41
3	BW4	A	502[A]	41/41	0.95	0.13	15,16,34,39	41
3	BW4	A	502[B]	41/41	0.95	0.13	16,18,40,45	41
3	BW4	B	502[A]	41/41	0.96	0.10	14,18,36,39	41
3	BW4	B	502[B]	41/41	0.96	0.10	14,17,35,38	41
3	BW4	C	502[A]	41/41	0.96	0.11	22,25,46,48	41
3	BW4	C	502[B]	41/41	0.96	0.11	22,24,41,44	41
2	SAH	C	501	26/26	0.97	0.08	21,23,24,25	0
2	SAH	D	501	26/26	0.97	0.07	23,26,29,29	0
2	SAH	A	501	26/26	0.98	0.09	14,15,17,18	0
2	SAH	B	501	26/26	0.98	0.07	15,16,18,19	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around BW4 D 502 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around BW4 D 502 (B):**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around BW4 A 502 (A):


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around BW4 A 502 (B):**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)













Electron density around SAH C 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around SAH D 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around SAH A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around SAH B 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers

There are no such residues in this entry.