



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 6, 2023 – 06:22 PM EST

PDB ID : 6ATI
Title : HLA-DRB1*1402 in complex with Vimentin-64Cit59-71
Authors : Scally, S.W.; Ting, Y.T.; Rossjohn, J.
Deposited on : 2017-08-29
Resolution : 1.98 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.98 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	11647 (2.00-1.96)
Clashscore	141614	1014 (1.98-1.98)
Ramachandran outliers	138981	1006 (1.98-1.98)
Sidechain outliers	138945	1006 (1.98-1.98)
RSRZ outliers	127900	11410 (2.00-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	189	 2% 89% 6% 5%
1	D	189	 3% 90% 5% 5%
2	B	200	 4% 86% 9% 5%
2	E	200	 2% 88% 6% 5%
3	C	13	 85% 8% 8%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	13	 85% 8% 8%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7232 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA class II histocompatibility antigen, DR alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	180	1486	962	240	279	5	0	1	0
1	D	180	1482	960	240	277	5	0	1	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	182	THR	-	expression tag	UNP P01903
A	183	SER	-	expression tag	UNP P01903
A	184	GLY	-	expression tag	UNP P01903
A	185	ASP	-	expression tag	UNP P01903
A	186	ASP	-	expression tag	UNP P01903
A	187	ASP	-	expression tag	UNP P01903
A	188	ASP	-	expression tag	UNP P01903
A	189	LYS	-	expression tag	UNP P01903
D	182	THR	-	expression tag	UNP P01903
D	183	SER	-	expression tag	UNP P01903
D	184	GLY	-	expression tag	UNP P01903
D	185	ASP	-	expression tag	UNP P01903
D	186	ASP	-	expression tag	UNP P01903
D	187	ASP	-	expression tag	UNP P01903
D	188	ASP	-	expression tag	UNP P01903
D	189	LYS	-	expression tag	UNP P01903

- Molecule 2 is a protein called MHC class II antigen.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	190	1551	973	279	294	5	0	0	0
2	E	190	1561	978	282	296	5	0	0	0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-1	GLY	-	expression tag	UNP A0A0A1I7H6
B	0	SER	-	expression tag	UNP A0A0A1I7H6
B	191	THR	-	expression tag	UNP A0A0A1I7H6
B	192	GLY	-	expression tag	UNP A0A0A1I7H6
B	193	GLY	-	expression tag	UNP A0A0A1I7H6
B	194	ASP	-	expression tag	UNP A0A0A1I7H6
B	195	ASP	-	expression tag	UNP A0A0A1I7H6
B	196	ASP	-	expression tag	UNP A0A0A1I7H6
B	197	ASP	-	expression tag	UNP A0A0A1I7H6
B	198	LYS	-	expression tag	UNP A0A0A1I7H6
E	-1	GLY	-	expression tag	UNP A0A0A1I7H6
E	0	SER	-	expression tag	UNP A0A0A1I7H6
E	191	THR	-	expression tag	UNP A0A0A1I7H6
E	192	GLY	-	expression tag	UNP A0A0A1I7H6
E	193	GLY	-	expression tag	UNP A0A0A1I7H6
E	194	ASP	-	expression tag	UNP A0A0A1I7H6
E	195	ASP	-	expression tag	UNP A0A0A1I7H6
E	196	ASP	-	expression tag	UNP A0A0A1I7H6
E	197	ASP	-	expression tag	UNP A0A0A1I7H6
E	198	LYS	-	expression tag	UNP A0A0A1I7H6

- Molecule 3 is a protein called Vimentin-64Cit59-71.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	13	Total	C	N	O	0	0	0
			100	61	21	18			
3	F	13	Total	C	N	O	0	0	0
			100	61	21	18			

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	D	1	Total	C	N	O	0	0
			14	8	1	5		
4	D	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	229	Total	O	0	0
			229	229		
5	B	200	Total	O	0	0
			200	200		
5	D	233	Total	O	0	0
			233	233		
5	E	197	Total	O	0	0
			197	197		
5	C	17	Total	O	0	0
			17	17		
5	F	20	Total	O	0	0
			20	20		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HLA class II histocompatibility antigen, DR alpha chain

Chain A: 




- Molecule 1: HLA class II histocompatibility antigen, DR alpha chain

Chain D: 




- Molecule 2: MHC class II antigen

Chain B: 




- Molecule 2: MHC class II antigen

Chain E: 




- Molecule 3: Vimentin-64Cit59-71

Chain C: 



- Molecule 3: Vimentin-64Cit59-71



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	67.27Å 76.87Å 94.91Å 90.00° 109.68° 90.00°	Depositor
Resolution (Å)	33.63 – 1.98 33.63 – 1.98	Depositor EDS
% Data completeness (in resolution range)	99.6 (33.63-1.98) 96.4 (33.63-1.98)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.22 (at 1.98Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.190 , 0.233 0.192 , 0.233	Depositor DCC
R_{free} test set	3221 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	14.4	Xtrriage
Anisotropy	0.283	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 50.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.147 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	7232	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, CIR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.29	0/1534	0.46	0/2092
1	D	0.29	0/1530	0.45	0/2087
2	B	0.28	0/1594	0.46	0/2169
2	E	0.29	0/1604	0.46	0/2181
3	C	0.24	0/88	0.48	0/116
3	F	0.24	0/88	0.45	0/116
All	All	0.29	0/6438	0.46	0/8761

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	C	0	1
3	F	0	1
All	All	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	C	6	CIR	Mainchain
3	F	6	CIR	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1486	0	1418	10	0
1	D	1482	0	1414	6	0
2	B	1551	0	1447	13	0
2	E	1561	0	1462	7	0
3	C	100	0	103	1	0
3	F	100	0	103	1	0
4	A	28	0	26	0	0
4	D	28	0	26	1	0
5	A	229	0	0	1	0
5	B	200	0	0	2	0
5	C	17	0	0	0	0
5	D	233	0	0	2	0
5	E	197	0	0	0	0
5	F	20	0	0	0	0
All	All	7232	0	5999	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:50:ARG:NH1	5:D:601:HOH:O	2.24	0.70
1:A:3:GLU:HG2	2:B:16:HIS:HB3	1.81	0.62
1:D:67:LYS:NZ	5:D:606:HOH:O	2.36	0.58
2:E:133:ARG:HD3	2:E:171:TYR:CZ	2.39	0.56
2:B:18:PHE:HB2	2:B:23:ARG:HB2	1.87	0.56
1:A:118:ASN:HB2	1:A:166:GLU:HB2	1.89	0.54
1:A:3:GLU:HG3	2:B:18:PHE:CE2	2.42	0.54
2:B:116:VAL:HG22	2:B:160:MET:HG2	1.90	0.54
2:E:116:VAL:HG22	2:E:160:MET:HG2	1.90	0.53
1:D:4:GLU:HG2	1:D:5:HIS:CD2	2.44	0.53
1:D:118:ASN:HB2	1:D:166:GLU:HB2	1.90	0.53
2:E:18:PHE:HB2	2:E:23:ARG:HB2	1.93	0.51
2:B:107:GLN:O	2:B:110:GLN:HB2	2.11	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:91:VAL:HG23	1:A:176:LYS:HB3	1.92	0.50
2:B:25:ARG:NH1	5:B:209:HOH:O	2.43	0.50
1:A:3:GLU:N	5:A:609:HOH:O	2.45	0.49
1:D:78:ASN:HD22	4:D:500:NAG:H83	1.77	0.48
2:B:130:ARG:NH2	2:B:174:GLN:OE1	2.47	0.48
1:A:4:GLU:HG2	1:A:5:HIS:CD2	2.49	0.47
2:E:107:GLN:O	2:E:110:GLN:HB2	2.15	0.47
2:B:19:ASN:ND2	5:B:205:HOH:O	2.36	0.46
2:B:129:VAL:HB	2:B:159:VAL:HG21	2.00	0.42
2:E:123:TYR:CG	2:E:124:PRO:HA	2.54	0.42
2:B:53:LEU:HD12	2:B:53:LEU:HA	1.89	0.42
1:A:94:ASN:HB3	1:A:106:ILE:HD11	2.02	0.41
1:A:29:ASP:HB3	2:B:153:TRP:CE2	2.55	0.41
2:B:81:HIS:O	2:B:85:VAL:HG23	2.20	0.41
2:E:71:ARG:NH2	3:F:7:SER:O	2.52	0.41
2:E:90:THR:OG1	2:E:91:VAL:N	2.53	0.41
1:A:97:VAL:HG21	1:A:178:TRP:HZ2	1.85	0.41
1:D:97:VAL:HG21	1:D:178:TRP:HZ2	1.86	0.41
2:B:71:ARG:NH2	3:C:7:SER:O	2.55	0.40
1:A:176:LYS:HA	1:A:176:LYS:HD2	1.90	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	179/189 (95%)	178 (99%)	1 (1%)	0	100	100
1	D	179/189 (95%)	178 (99%)	1 (1%)	0	100	100
2	B	188/200 (94%)	184 (98%)	4 (2%)	0	100	100
2	E	188/200 (94%)	184 (98%)	4 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	C	10/13 (77%)	10 (100%)	0	0	100	100
3	F	10/13 (77%)	10 (100%)	0	0	100	100
All	All	754/804 (94%)	744 (99%)	10 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	166/173 (96%)	166 (100%)	0	100	100
1	D	165/173 (95%)	165 (100%)	0	100	100
2	B	170/179 (95%)	170 (100%)	0	100	100
2	E	172/179 (96%)	172 (100%)	0	100	100
3	C	9/9 (100%)	9 (100%)	0	100	100
3	F	9/9 (100%)	9 (100%)	0	100	100
All	All	691/722 (96%)	691 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	CIR	F	6	3	9,10,11	3.16	3 (33%)	6,11,13	1.57	2 (33%)
3	CIR	C	6	3	9,10,11	3.31	3 (33%)	6,11,13	2.34	4 (66%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CIR	F	6	3	-	1/8/9/11	-
3	CIR	C	6	3	-	1/8/9/11	-

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	6	CIR	C7-N6	8.03	1.45	1.34
3	F	6	CIR	C7-N6	7.51	1.44	1.34
3	C	6	CIR	C7-N8	4.90	1.44	1.33
3	F	6	CIR	C7-N8	4.88	1.44	1.33
3	F	6	CIR	C3-CA	-2.13	1.50	1.53
3	C	6	CIR	C3-CA	-2.12	1.50	1.53

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	6	CIR	N8-C7-N6	3.91	120.98	116.85
3	F	6	CIR	N8-C7-N6	2.50	119.49	116.85
3	C	6	CIR	O7-C7-N8	-2.48	118.97	123.22
3	C	6	CIR	C5-N6-C7	-2.44	119.83	122.73
3	C	6	CIR	O7-C7-N6	-2.28	120.05	121.74
3	F	6	CIR	O7-C7-N8	-2.05	119.70	123.22

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	F	6	CIR	CA-C3-C4-C5
3	C	6	CIR	CA-C3-C4-C5

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	D	501	1	14,14,15	0.22	0	17,19,21	0.49	0
4	NAG	D	500	1	14,14,15	0.23	0	17,19,21	0.35	0
4	NAG	A	501	1	14,14,15	0.25	0	17,19,21	0.40	0
4	NAG	A	500	1	14,14,15	0.37	0	17,19,21	0.36	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	D	501	1	-	4/6/23/26	0/1/1/1
4	NAG	D	500	1	-	4/6/23/26	0/1/1/1
4	NAG	A	501	1	-	0/6/23/26	0/1/1/1
4	NAG	A	500	1	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	500	NAG	C8-C7-N2-C2
4	A	500	NAG	O7-C7-N2-C2
4	D	500	NAG	C8-C7-N2-C2
4	D	500	NAG	O7-C7-N2-C2
4	D	501	NAG	C8-C7-N2-C2
4	D	501	NAG	O7-C7-N2-C2
4	D	501	NAG	O5-C5-C6-O6
4	D	501	NAG	C4-C5-C6-O6
4	D	500	NAG	C4-C5-C6-O6
4	D	500	NAG	O5-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	500	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	180/189 (95%)	-0.12	3 (1%) 70 71	9, 17, 41, 55	0
1	D	180/189 (95%)	-0.08	5 (2%) 53 55	10, 17, 41, 57	0
2	B	190/200 (95%)	-0.11	7 (3%) 41 44	10, 17, 43, 76	0
2	E	190/200 (95%)	-0.13	5 (2%) 56 58	10, 16, 42, 77	0
3	C	12/13 (92%)	-0.25	0 100 100	12, 18, 22, 35	0
3	F	12/13 (92%)	-0.28	0 100 100	14, 16, 25, 37	0
All	All	764/804 (95%)	-0.11	20 (2%) 56 58	9, 17, 42, 77	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	109	LEU	4.0
1	A	100	ARG	4.0
2	E	191	THR	3.8
2	B	107	GLN	3.8
1	D	100	ARG	3.5
2	E	168	GLY	3.1
2	B	108	PRO	3.0
1	D	171	ASP	2.7
2	E	107	GLN	2.7
2	B	2	ASP	2.5
1	A	182	THR	2.4
2	B	191	THR	2.4
1	D	157	THR	2.3
1	D	3	GLU	2.3
2	B	111	HIS	2.3
1	D	129	THR	2.3
2	B	134	ASN	2.1
2	E	110	GLN	2.1
1	A	154	LEU	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	E	111	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	CIR	C	6	11/12	0.96	0.10	9,13,17,18	0
3	CIR	F	6	11/12	0.96	0.10	11,14,18,22	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	D	500	14/15	0.75	0.23	37,50,52,54	0
4	NAG	D	501	14/15	0.79	0.16	32,39,43,46	0
4	NAG	A	500	14/15	0.80	0.22	33,39,43,45	0
4	NAG	A	501	14/15	0.84	0.20	24,38,45,47	0

6.5 Other polymers [i](#)

There are no such residues in this entry.