



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 19, 2024 – 05:18 AM EDT

PDB ID : 4AUP
Title : Tuber borchii Phospholipase A2
Authors : Cavazzini, D.; Meschi, F.; Corsini, R.; Bolchi, A.; Rossi, G.-L.; Einsle, O.; Ottonello, S.
Deposited on : 2012-05-18
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	123	
1	B	123	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	ACT	A	1212	-	-	X	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2198 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PHOSPHOLIPASE A2 GROUP XIII.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	120	Total 983	C 616	N 179	O 183	S 5	0	0	0
1	B	123	Total 1001	C 628	N 182	O 186	S 5	0	0	0

- Molecule 2 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	Total 4	C 2	O 2	0	0
2	B	1	Total 4	C 2	O 2	0	0

- Molecule 3 is THIOCYANATE ION (three-letter code: SCN) (formula: CNS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	S		
3	B	1	3	1	1	1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	94	94	94	0	0
4	B	109	109	109	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: PHOSPHOLIPASE A2 GROUP XIII

Chain A: 



- Molecule 1: PHOSPHOLIPASE A2 GROUP XIII

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	34.50Å 66.84Å 117.38Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.41 – 1.90 19.41 – 1.90	Depositor EDS
% Data completeness (in resolution range)	98.3 (19.41-1.90) 98.5 (19.41-1.90)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.81 (at 1.90Å)	Xtrriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.210 , 0.273 0.212 , 0.277	Depositor DCC
R_{free} test set	1112 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å ²)	25.0	Xtrriage
Anisotropy	0.030	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 46.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2198	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, SCN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.94	1/1009 (0.1%)	1.06	5/1357 (0.4%)
1	B	0.96	0/1027	0.90	4/1382 (0.3%)
All	All	0.95	1/2036 (0.0%)	0.98	9/2739 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	123	TRP	CD2-CE2	5.65	1.48	1.41

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	206	ARG	NE-CZ-NH2	-9.46	115.57	120.30
1	A	195	ARG	NE-CZ-NH2	-8.04	116.28	120.30
1	B	96	ASP	CB-CG-OD1	7.83	125.35	118.30
1	A	131	SER	N-CA-C	-7.63	90.39	111.00
1	A	189	TRP	CB-CA-C	6.21	122.82	110.40
1	B	206	ARG	NE-CZ-NH1	6.15	123.37	120.30
1	A	188	SER	C-N-CA	5.61	135.73	121.70
1	B	133	ASP	CB-CG-OD1	5.36	123.13	118.30
1	A	206	ARG	NE-CZ-NH1	5.03	122.81	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	129	SER	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	983	0	933	24	1
1	B	1001	0	954	1	0
2	A	4	0	3	6	0
2	B	4	0	3	0	0
3	B	3	0	0	0	0
4	A	94	0	0	2	1
4	B	109	0	0	1	0
All	All	2198	0	1893	25	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:117:ASN:OD1	2:A:1212:ACT:H1	1.47	1.11
1:A:187:GLU:HB3	1:A:190:LYS:HB2	1.56	0.88
1:A:186:LEU:HG	1:A:187:GLU:H	1.48	0.79
1:A:142:ASP:OD2	4:A:2045:HOH:O	2.06	0.73
1:A:186:LEU:HG	1:A:187:GLU:N	2.13	0.61
1:A:138:PHE:CE1	1:A:190:LYS:HG3	2.35	0.61
1:A:165:ARG:NH1	4:A:2064:HOH:O	2.34	0.60
1:A:187:GLU:HB3	1:A:190:LYS:CB	2.31	0.60
1:A:117:ASN:OD1	2:A:1212:ACT:CH3	2.37	0.59
1:A:138:PHE:HE1	1:A:190:LYS:HG3	1.69	0.58
1:A:186:LEU:CG	1:A:187:GLU:H	2.18	0.55
1:A:188:SER:N	1:A:189:TRP:HB3	2.22	0.54
1:A:188:SER:H	1:A:189:TRP:HB3	1.73	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:117:ASN:CG	2:A:1212:ACT:H1	2.24	0.51
1:A:134:ARG:HA	1:A:138:PHE:O	2.10	0.50
1:A:185:GLY:H	1:A:186:LEU:HB3	1.76	0.50
1:B:114:ARG:NH2	4:B:2020:HOH:O	2.47	0.47
1:A:186:LEU:O	1:A:187:GLU:C	2.53	0.47
1:A:117:ASN:HB3	2:A:1212:ACT:CH3	2.46	0.46
1:A:118:PRO:HB2	1:A:120:ASN:HD21	1.80	0.46
1:A:117:ASN:HD22	1:A:118:PRO:HA	1.83	0.43
1:A:118:PRO:HB2	1:A:120:ASN:ND2	2.34	0.42
1:A:117:ASN:CB	2:A:1212:ACT:H1	2.50	0.41
1:A:117:ASN:HB3	2:A:1212:ACT:H1	2.01	0.41
1:A:187:GLU:CB	1:A:190:LYS:HB2	2.38	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:188:SER:CB	4:A:2037:HOH:O[4_465]	2.08	0.12

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	118/123 (96%)	111 (94%)	2 (2%)	5 (4%)	3 0
1	B	121/123 (98%)	118 (98%)	2 (2%)	1 (1%)	19 9
All	All	239/246 (97%)	229 (96%)	4 (2%)	6 (2%)	5 1

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	186	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	189	TRP
1	B	129	SER
1	A	129	SER
1	A	130	LYS
1	A	187	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	104/105 (99%)	99 (95%)	5 (5%)	25 16
1	B	105/105 (100%)	103 (98%)	2 (2%)	57 53
All	All	209/210 (100%)	202 (97%)	7 (3%)	38 29

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	120	ASN
1	A	121	LEU
1	A	130	LYS
1	A	190	LYS
1	A	196	LYS
1	B	141	LEU
1	B	205	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	120	ASN
1	A	139	ASN
1	A	171	ASN
1	B	139	ASN
1	B	164	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ACT	B	1213	-	3,3,3	0.71	0	3,3,3	1.24	0
3	SCN	B	1212	-	1,2,2	0.62	0	0,1,1	-	-
2	ACT	A	1212	-	3,3,3	1.03	0	3,3,3	2.97	2 (66%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	A	1212	ACT	O-C-CH3	-3.98	106.21	122.53
2	A	1212	ACT	OXT-C-CH3	3.12	128.15	115.05

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1212	ACT	6	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	120/123 (97%)	0.50	8 (6%) 17 20	15, 29, 59, 78	0
1	B	123/123 (100%)	0.07	3 (2%) 59 62	15, 24, 39, 60	0
All	All	243/246 (98%)	0.29	11 (4%) 33 36	15, 26, 49, 78	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	158	HIS	6.1
1	A	188	SER	4.7
1	A	184	SER	4.5
1	A	186	LEU	4.4
1	A	185	GLY	4.2
1	B	129	SER	3.9
1	B	130	LYS	3.7
1	A	130	LYS	3.1
1	A	189	TRP	3.0
1	A	187	GLU	2.7
1	B	196	LYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ACT	A	1212	4/4	0.91	0.18	16,23,24,26	0
2	ACT	B	1213	4/4	0.96	0.08	29,31,32,34	0
3	SCN	B	1212	3/3	0.99	0.06	16,16,18,19	0

6.5 Other polymers

There are no such residues in this entry.