



wwPDB X-ray Structure Validation Summary Report ⓘ

May 26, 2020 – 05:48 pm BST

PDB ID : 3AW9
Title : Structure of UDP-galactose 4-epimerase mutant
Authors : Sakuraba, H.; Kawai, T.; Yoneda, K.; Ohshima, T.
Deposited on : 2011-03-15
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	308	
1	B	308	
1	C	308	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7500 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NAD-dependent epimerase/dehydratase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	304	Total 2346	C 1502	N 407	O 428	S 9	0	0	0
1	B	300	Total 2331	C 1494	N 403	O 425	S 9	0	0	0
1	C	299	Total 2324	C 1490	N 402	O 423	S 9	0	0	0

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	32	ILE	-	SEE REMARK 999	UNP A3MUJ4
A	33	VAL	-	SEE REMARK 999	UNP A3MUJ4
A	34	GLN	-	SEE REMARK 999	UNP A3MUJ4
A	36	ASP	-	SEE REMARK 999	UNP A3MUJ4
A	37	THR	-	SEE REMARK 999	UNP A3MUJ4
A	38	GLY	-	SEE REMARK 999	UNP A3MUJ4
A	39	GLY	-	SEE REMARK 999	UNP A3MUJ4
B	1032	ILE	-	SEE REMARK 999	UNP A3MUJ4
B	1033	VAL	-	SEE REMARK 999	UNP A3MUJ4
B	1034	GLN	-	SEE REMARK 999	UNP A3MUJ4
B	1036	ASP	-	SEE REMARK 999	UNP A3MUJ4
B	1037	THR	-	SEE REMARK 999	UNP A3MUJ4
B	1038	GLY	-	SEE REMARK 999	UNP A3MUJ4
B	1039	GLY	-	SEE REMARK 999	UNP A3MUJ4
C	2032	ILE	-	SEE REMARK 999	UNP A3MUJ4
C	2033	VAL	-	SEE REMARK 999	UNP A3MUJ4
C	2034	GLN	-	SEE REMARK 999	UNP A3MUJ4
C	2036	ASP	-	SEE REMARK 999	UNP A3MUJ4
C	2037	THR	-	SEE REMARK 999	UNP A3MUJ4
C	2038	GLY	-	SEE REMARK 999	UNP A3MUJ4
C	2039	GLY	-	SEE REMARK 999	UNP A3MUJ4


- Molecule 2 is GALACTOSE-URIDINE-5'-DIPHOSPHATE (three-letter code: GDU)

(formula: $C_{15}H_{24}N_2O_{17}P_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
2	A	1	Total	C	N	O	P	0	0
			36	15	2	17	2		
2	B	1	Total	C	N	O	P	0	0
			36	15	2	17	2		
2	C	1	Total	C	N	O	P	0	0
			36	15	2	17	2		

- Molecule 3 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: $C_{21}H_{27}N_7O_{14}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	B	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	C	1	Total	C	N	O	P	0	0
			44	21	7	14	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	88	Total	O	0	0
			88	88		
4	B	103	Total	O	0	0
			103	103		
4	C	68	Total	O	0	0
			68	68		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: NAD-dependent epimerase/dehydratase



- Molecule 1: NAD-dependent epimerase/dehydratase



- Molecule 1: NAD-dependent epimerase/dehydratase



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	88.62Å 113.64Å 217.27Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.30 30.54 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.7 (50.00-2.30) 99.7 (30.54-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	15.10 (at 2.31Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.204 , 0.236 0.203 , 0.234	Depositor DCC
R_{free} test set	2467 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	32.3	Xtrriage
Anisotropy	0.893	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 39.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7500	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GDU, NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.61	0/2394	0.66	0/3256
1	B	0.65	0/2378	0.71	2/3233 (0.1%)
1	C	0.63	0/2371	0.69	3/3223 (0.1%)
All	All	0.63	0/7143	0.69	5/9712 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	2016	LEU	CA-CB-CG	6.13	129.39	115.30
1	B	1043	LEU	CA-CB-CG	5.62	128.22	115.30
1	C	2043	LEU	CA-CB-CG	5.19	127.25	115.30
1	C	2236	VAL	CB-CA-C	-5.19	101.53	111.40
1	B	1016	LEU	CA-CB-CG	5.06	126.94	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	2077	GLU	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2346	0	2371	33	0
1	B	2331	0	2358	23	0
1	C	2324	0	2351	32	0
2	A	36	0	22	2	0
2	B	36	0	22	1	0
2	C	36	0	22	2	0
3	A	44	0	26	2	0
3	B	44	0	26	2	0
3	C	44	0	26	2	0
4	A	88	0	0	4	0
4	B	103	0	0	2	0
4	C	68	0	0	0	0
All	All	7500	0	7224	87	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 87 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:2192:ASP:OD1	1:C:2194:THR:HG22	1.44	1.15
1:A:258:SER:HB2	4:A:364:HOH:O	1.57	1.01
1:A:234:VAL:HG23	1:A:291:SER:HB3	1.60	0.83
1:C:2014:SER:OG	1:C:2167:ARG:NH1	2.13	0.81
1:B:1241:GLN:HG3	4:B:164:HOH:O	1.80	0.81

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	300/308 (97%)	292 (97%)	8 (3%)	0	100	100
1	B	294/308 (96%)	287 (98%)	6 (2%)	1 (0%)	41	50
1	C	293/308 (95%)	285 (97%)	8 (3%)	0	100	100
All	All	887/924 (96%)	864 (97%)	22 (2%)	1 (0%)	51	64

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	1076	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	246/251 (98%)	241 (98%)	5 (2%)	55	72
1	B	246/251 (98%)	240 (98%)	6 (2%)	49	66
1	C	245/251 (98%)	236 (96%)	9 (4%)	34	48
All	All	737/753 (98%)	717 (97%)	20 (3%)	44	61

5 of 20 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	1226	LEU
1	B	1254	ARG
1	C	2236	VAL
1	B	1043	LEU
1	B	1190	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	227	ASN
1	C	2081	HIS
1	B	1015	HIS
1	A	34	GLN
1	B	1185	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAD	A	901	-	42,48,48	1.31	2 (4%)	50,73,73	1.33	4 (8%)
2	GDU	B	1801	-	31,38,38	1.91	4 (12%)	41,58,58	2.38	9 (21%)
2	GDU	C	2801	-	31,38,38	1.88	5 (16%)	41,58,58	2.29	12 (29%)
3	NAD	B	1901	-	42,48,48	1.53	4 (9%)	50,73,73	1.22	4 (8%)
2	GDU	A	801	-	31,38,38	1.77	3 (9%)	41,58,58	1.53	8 (19%)
3	NAD	C	2901	-	42,48,48	1.42	4 (9%)	50,73,73	1.28	6 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAD	A	901	-	-	4/26/62/62	0/5/5/5
2	GDU	B	1801	-	-	12/21/59/59	0/3/3/3
2	GDU	C	2801	-	-	8/21/59/59	0/3/3/3
3	NAD	B	1901	-	-	6/26/62/62	0/5/5/5
2	GDU	A	801	-	-	5/21/59/59	0/3/3/3
3	NAD	C	2901	-	-	6/26/62/62	0/5/5/5

The worst 5 of 22 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1801	GDU	C6-N1	7.33	1.44	1.35
3	B	1901	NAD	C2N-N1N	6.47	1.42	1.35
2	A	801	GDU	C6-N1	6.23	1.43	1.35
2	C	2801	GDU	C6-N1	6.09	1.43	1.35
3	C	2901	NAD	C2N-N1N	5.75	1.42	1.35

The worst 5 of 43 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1801	GDU	O3B-C1'-C2'	8.91	124.70	108.38
2	C	2801	GDU	O3B-C1'-C2'	6.68	120.63	108.38
2	B	1801	GDU	O5'-C1'-O3B	-5.03	104.79	111.36
3	B	1901	NAD	N3A-C2A-N1A	-5.01	120.84	128.68
2	C	2801	GDU	O5'-C1'-C2'	-4.95	99.87	110.35

There are no chirality outliers.

5 of 41 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	1801	GDU	C2D-C1D-N1-C6
2	B	1801	GDU	O4D-C1D-N1-C6
2	B	1801	GDU	C5D-O5D-PA-O2A
2	B	1801	GDU	PB-O3A-PA-O5D
2	B	1801	GDU	O5'-C1'-O3B-PB



There are no ring outliers.


6 monomers are involved in 11 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	901	NAD	2	0
2	B	1801	GDU	1	0
2	C	2801	GDU	2	0
3	B	1901	NAD	2	0
2	A	801	GDU	2	0
3	C	2901	NAD	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	304/308 (98%)	0.14	16 (5%) 26 33	18, 35, 51, 64	0
1	B	300/308 (97%)	0.03	14 (4%) 31 38	17, 31, 51, 66	0
1	C	299/308 (97%)	0.00	6 (2%) 65 71	17, 35, 53, 63	0
All	All	903/924 (97%)	0.06	36 (3%) 38 45	17, 33, 51, 66	0

The worst 5 of 36 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1075	THR	5.3
1	B	1071	VAL	4.4
1	B	1076	THR	4.4
1	A	75	THR	4.1
1	B	1036	ASP	4.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	GDU	A	801	36/36	0.78	0.35	47,53,59,60	11
2	GDU	C	2801	36/36	0.82	0.28	10,37,41,45	11
2	GDU	B	1801	36/36	0.85	0.28	23,37,43,45	11
3	NAD	C	2901	44/44	0.96	0.12	23,27,33,36	0
3	NAD	A	901	44/44	0.97	0.12	23,29,38,39	0
3	NAD	B	1901	44/44	0.97	0.11	24,28,34,35	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around GDU C 2801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around GDU B 1801:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around NAD C 2901:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NAD A 901:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.