



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 7, 2020 – 10:24 AM BST

PDB ID : 3B4V  
Title : X-Ray structure of Activin in complex with FSTL3  
Authors : Thompson, T.B.  
Deposited on : 2007-10-24  
Resolution : 2.48 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.48 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5857 (2.50-2.46)
Clashscore	141614	6594 (2.50-2.46)
Ramachandran outliers	138981	6469 (2.50-2.46)
Sidechain outliers	138945	6471 (2.50-2.46)
RSRZ outliers	127900	5738 (2.50-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	116	<div style="display: flex; align-items: center;"> <div style="width: 7%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 85%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 9%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 6%; height: 10px; background-color: grey;"></div> </div> <p style="text-align: center;">85%      9%      6%</p>
1	B	116	<div style="display: flex; align-items: center;"> <div style="width: 6%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 87%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 11%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 6%; height: 10px; background-color: grey;"></div> </div> <p style="text-align: center;">87%      11%      •</p>
1	E	116	<div style="display: flex; align-items: center;"> <div style="width: 7%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 76%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 16%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 8%; height: 10px; background-color: grey;"></div> </div> <p style="text-align: center;">76%      16%      •      8%</p>
1	F	116	<div style="display: flex; align-items: center;"> <div style="width: 3%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 85%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 12%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 6%; height: 10px; background-color: grey;"></div> </div> <p style="text-align: center;">85%      12%      ••</p>
2	C	237	<div style="display: flex; align-items: center;"> <div style="width: 7%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 79%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 8%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 13%; height: 10px; background-color: grey;"></div> </div> <p style="text-align: center;">79%      8%      13%</p>
2	D	237	<div style="display: flex; align-items: center;"> <div style="width: 3%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 79%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 11%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 10%; height: 10px; background-color: grey;"></div> </div> <p style="text-align: center;">79%      11%      10%</p>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	G	237	
2	H	237	

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 9726 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Inhibin beta A chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	109	Total 855	C 540	N 147	O 155	S 13	0	0	0
1	B	114	Total 886	C 558	N 153	O 162	S 13	0	0	0
1	E	107	Total 845	C 534	N 145	O 153	S 13	0	0	0
1	F	114	Total 880	C 553	N 152	O 162	S 13	0	0	0

- Molecule 2 is a protein called follistatin-like 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	206	Total 1494	C 896	N 287	O 282	S 29	0	0	0
2	D	213	Total 1533	C 919	N 296	O 289	S 29	0	0	0
2	G	206	Total 1490	C 894	N 286	O 281	S 29	0	0	0
2	H	211	Total 1521	C 913	N 293	O 286	S 29	0	0	0

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	G	1	Total C O 4 2 2	0	0

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	C	1	Total O S 5 4 1	0	0

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
5	C	1	14	8	1	5	0	0
5	G	1	14	8	1	5	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
6	A	19	19	19	0	0
6	B	19	19	19	0	0
6	C	16	16	16	0	0
6	D	36	36	36	0	0
6	E	15	15	15	0	0
6	F	16	16	16	0	0
6	G	22	22	22	0	0
6	H	34	34	34	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Inhibin beta A chain




- Molecule 1: Inhibin beta A chain




- Molecule 1: Inhibin beta A chain




- Molecule 1: Inhibin beta A chain






- Molecule 2: follistatin-like 3











- Molecule 2: follistatin-like 3



- Molecule 2: follistatin-like 3



- Molecule 2: follistatin-like 3





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	63.61Å 71.38Å 100.20Å 98.55° 90.64° 90.11°	Depositor
Resolution (Å)	31.78 – 2.48 31.01 – 2.48	Depositor EDS
% Data completeness (in resolution range)	98.6 (31.78-2.48) 95.5 (31.01-2.48)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.68 (at 2.48Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.227 , 0.279 0.244 , 0.270	Depositor DCC
$R_{free}$ test set	3144 reflections (5.14%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	52.9	Xtriage
Anisotropy	0.023	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.27 , 2.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.407 for h,-k,-l	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	9726	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.67% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, EDO, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.51	0/876	0.58	0/1178
1	B	0.45	0/907	0.52	0/1218
1	E	0.49	0/866	0.57	0/1164
1	F	0.83	3/901 (0.3%)	0.54	0/1211
2	C	0.42	0/1521	0.51	0/2060
2	D	0.53	2/1563 (0.1%)	0.55	0/2121
2	G	0.56	2/1517 (0.1%)	0.56	1/2055 (0.0%)
2	H	0.72	3/1550 (0.2%)	0.54	0/2103
All	All	0.58	10/9701 (0.1%)	0.55	1/13110 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	D	0	1

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	6	GLY	N-CA	17.18	1.71	1.46
2	H	6	PRO	CA-CB	14.27	1.82	1.53
2	H	6	PRO	N-CA	14.20	1.71	1.47
2	G	74	ASP	CG-OD2	11.19	1.51	1.25
2	G	74	ASP	CG-OD1	10.22	1.48	1.25
1	F	6	GLY	C-O	10.10	1.39	1.23
2	H	6	PRO	CG-CD	8.52	1.78	1.50
2	D	33	ALA	C-N	8.41	1.53	1.34
2	D	33	ALA	C-O	6.07	1.34	1.23
1	F	6	GLY	CA-C	5.33	1.60	1.51

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	G	74	ASP	CB-CG-OD1	-8.71	110.46	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	D	33	ALA	Mainchain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	855	0	801	9	0
1	B	886	0	843	10	0
1	E	845	0	797	15	0
1	F	880	0	825	17	0
2	C	1494	0	1407	8	0
2	D	1533	0	1438	11	0
2	G	1490	0	1401	13	0
2	H	1521	0	1425	15	0
3	A	4	0	6	0	0
3	C	4	0	6	0	0
3	G	4	0	6	0	0
4	C	5	0	0	0	0
5	C	14	0	13	1	0
5	G	14	0	13	0	0
6	A	19	0	0	0	0
6	B	19	0	0	0	0
6	C	16	0	0	0	0
6	D	36	0	0	0	0
6	E	15	0	0	0	0
6	F	16	0	0	0	0
6	G	22	0	0	2	0
6	H	34	0	0	0	0
All	All	9726	0	8981	77	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (77) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:6:PRO:CA	2:H:6:PRO:CB	1.82	1.57
2:H:6:PRO:CD	2:H:6:PRO:CG	1.78	1.53
2:H:6:PRO:CA	2:H:6:PRO:N	1.71	1.53
1:F:6:GLY:CA	1:F:6:GLY:N	1.71	1.50
2:H:6:PRO:CD	2:H:6:PRO:CA	2.63	0.76
2:H:6:PRO:CA	2:H:6:PRO:CG	2.69	0.70
2:H:6:PRO:CD	2:H:6:PRO:CB	2.71	0.66
1:F:45:PRO:HG2	1:F:56:LEU:HD21	1.84	0.60
2:H:6:PRO:C	2:H:6:PRO:CB	2.68	0.60
2:H:6:PRO:N	2:H:6:PRO:CG	2.62	0.58
2:C:159:ASP:OD1	2:C:161:THR:HB	2.03	0.57
2:D:106:LEU:O	2:D:118:ASP:HA	2.04	0.57
1:B:86:LEU:HD23	1:B:110:VAL:HA	1.87	0.56
1:E:110:VAL:HG23	1:F:59:HIS:NE2	2.21	0.56
1:E:45:PRO:HG2	1:E:56:LEU:HD21	1.88	0.56
1:F:6:GLY:C	1:F:6:GLY:N	2.55	0.55
1:A:18:VAL:HG21	1:B:62:VAL:CG1	2.37	0.55
2:C:117:ARG:HB2	2:C:121:GLU:HG3	1.89	0.55
1:A:82:VAL:HG21	1:B:82:VAL:HG11	1.90	0.54
1:A:110:VAL:HG23	1:B:59:HIS:NE2	2.23	0.53
1:F:55:SER:HB3	2:G:24:LEU:HG	1.90	0.53
2:C:189:ASN:OD1	2:C:212:ARG:NH1	2.42	0.52
2:H:11:VAL:HG21	2:H:55:ILE:HG12	1.91	0.52
2:C:98:ASP:OD1	2:C:100:SER:HB3	2.11	0.51
1:F:65:HIS:HD2	6:G:248:HOH:O	1.93	0.49
1:A:62:VAL:CG1	1:B:18:VAL:HG21	2.42	0.49
2:G:159:ASP:OD1	2:G:161:THR:HB	2.12	0.49
2:C:71:ASP:OD2	2:C:71:ASP:N	2.44	0.49
2:H:43:THR:HG23	2:H:69:CYS:SG	2.53	0.48
2:G:31:THR:HG23	2:G:34:GLU:H	1.79	0.48
1:F:30:ILE:CG2	2:G:167:VAL:HG11	2.43	0.48
1:E:38:ASN:HB2	1:E:110:VAL:HB	1.96	0.48
2:D:117:ARG:NH2	2:D:121:GLU:OE1	2.48	0.47
1:E:59:HIS:NE2	1:F:110:VAL:HG23	2.30	0.47
2:H:119:GLU:OE2	2:H:123:ARG:NH1	2.48	0.47
2:D:11:VAL:HG21	2:D:55:ILE:HG12	1.97	0.47
1:F:97:GLY:O	2:G:210:GLY:HA2	2.15	0.47

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:55:SER:CB	2:G:24:LEU:HG	2.46	0.46
1:E:18:VAL:HG13	1:E:23:ILE:HD11	1.98	0.46
2:D:156:CYS:HA	2:D:165:HIS:O	2.15	0.46
1:F:100:ILE:HG22	2:G:159:ASP:HA	1.98	0.46
2:C:126:ARG:HD3	2:C:133:LEU:O	2.15	0.45
1:E:18:VAL:HG22	1:F:66:TYR:CE1	2.50	0.45
2:D:43:THR:HG23	2:D:69:CYS:SG	2.56	0.45
1:E:82:VAL:HG11	1:F:82:VAL:HG11	1.97	0.45
1:E:87:ARG:HB2	1:E:88:PRO:HD2	1.97	0.45
1:F:31:ALA:HB2	2:G:167:VAL:HG22	1.98	0.45
1:B:9:ASN:OD1	1:B:9:ASN:N	2.44	0.44
1:E:18:VAL:HG21	1:F:62:VAL:CG1	2.48	0.44
1:A:55:SER:HB3	2:D:25:VAL:O	2.18	0.43
1:F:65:HIS:CD2	6:G:248:HOH:O	2.71	0.43
2:H:106:LEU:O	2:H:118:ASP:HA	2.18	0.43
2:C:187:ASN:O	5:C:239:NAG:H82	2.18	0.43
2:G:43:THR:HG23	2:G:69:CYS:SG	2.59	0.43
1:B:25:TRP:CG	2:D:57:LEU:HD22	2.53	0.43
2:G:176:VAL:HA	2:G:177:PRO:HD3	1.86	0.43
2:H:12:CYS:HB2	2:H:30:VAL:HG21	2.00	0.43
1:E:2:LEU:HD23	1:E:10:ILE:HD13	1.99	0.43
2:G:102:LEU:HD13	2:G:119:GLU:HG2	2.01	0.43
1:A:72:SER:HA	1:A:73:PRO:HA	1.91	0.42
2:C:73:CYS:O	2:C:76:VAL:HG22	2.19	0.42
1:A:59:HIS:NE2	1:B:110:VAL:HG23	2.35	0.42
2:D:85:ARG:NE	2:D:94:GLU:OE2	2.52	0.42
2:D:50:HIS:HB3	2:D:54:LYS:HB3	2.02	0.42
1:E:103:LYS:HB3	1:E:103:LYS:HE2	1.70	0.42
2:D:15:GLN:OE1	2:D:20:ALA:HB1	2.19	0.41
2:H:7:ALA:HA	2:H:8:PRO:HD2	1.86	0.41
1:A:18:VAL:HG22	1:B:66:TYR:CE1	2.56	0.41
1:E:2:LEU:HD12	1:E:2:LEU:HA	1.94	0.41
2:G:156:CYS:HA	2:G:165:HIS:O	2.20	0.41
1:B:45:PRO:HG2	1:B:56:LEU:HD21	2.01	0.41
1:E:25:TRP:HB3	1:E:29:ILE:HD12	2.02	0.41
2:H:150:CYS:HA	2:H:151:PRO:HD2	1.90	0.41
1:A:58:PHE:HA	2:D:60:PHE:CE2	2.55	0.41
1:E:41:GLU:HG3	1:F:77:LEU:HD11	2.02	0.40
1:E:72:SER:HA	1:E:73:PRO:HA	1.86	0.40
2:G:14:LEU:HB2	2:G:26:LEU:HD13	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	103/116 (89%)	99 (96%)	3 (3%)	1 (1%)	15	26
1	B	108/116 (93%)	102 (94%)	6 (6%)	0	100	100
1	E	101/116 (87%)	97 (96%)	3 (3%)	1 (1%)	15	26
1	F	109/116 (94%)	103 (94%)	6 (6%)	0	100	100
2	C	202/237 (85%)	195 (96%)	7 (4%)	0	100	100
2	D	211/237 (89%)	201 (95%)	10 (5%)	0	100	100
2	G	202/237 (85%)	198 (98%)	4 (2%)	0	100	100
2	H	207/237 (87%)	200 (97%)	7 (3%)	0	100	100
All	All	1243/1412 (88%)	1195 (96%)	46 (4%)	2 (0%)	47	66

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	38	ASN
1	E	38	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	95/102 (93%)	95 (100%)	0	100	100
1	B	99/102 (97%)	97 (98%)	2 (2%)	55	77
1	E	95/102 (93%)	94 (99%)	1 (1%)	73	88

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	F	97/102 (95%)	97 (100%)	0	100	100
2	C	166/191 (87%)	162 (98%)	4 (2%)	49	72
2	D	168/191 (88%)	162 (96%)	6 (4%)	35	58
2	G	165/191 (86%)	160 (97%)	5 (3%)	41	65
2	H	167/191 (87%)	158 (95%)	9 (5%)	22	40
All	All	1052/1172 (90%)	1025 (97%)	27 (3%)	46	70

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	21	LYS
1	B	54	SER
2	C	21	THR
2	C	66	CYS
2	C	67	LEU
2	C	152	ARG
2	D	14	LEU
2	D	66	CYS
2	D	67	LEU
2	D	161	THR
2	D	167	VAL
2	D	189	ASN
1	E	85	LYS
2	G	66	CYS
2	G	67	LEU
2	G	168	VAL
2	G	195	SER
2	G	216	SER
2	H	14	LEU
2	H	21	THR
2	H	24	LEU
2	H	26	LEU
2	H	66	CYS
2	H	67	LEU
2	H	76	VAL
2	H	115	THR
2	H	161	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	F	65	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
5	NAG	G	238	2	14,14,15	0.55	0	17,19,21	1.64	2 (11%)
3	EDO	C	240	-	3,3,3	0.51	0	2,2,2	0.18	0
3	EDO	G	239	-	3,3,3	0.51	0	2,2,2	0.17	0
4	SO4	C	238	-	4,4,4	0.14	0	6,6,6	0.17	0
5	NAG	C	239	2	14,14,15	0.80	0	17,19,21	1.41	2 (11%)
3	EDO	A	117	-	3,3,3	0.50	0	2,2,2	0.21	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	C	240	-	-	1/1/1/1	-
5	NAG	G	238	2	-	5/6/23/26	0/1/1/1
5	NAG	C	239	2	-	4/6/23/26	0/1/1/1
3	EDO	A	117	-	-	1/1/1/1	-
3	EDO	G	239	-	-	1/1/1/1	-

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	G	238	NAG	C2-N2-C7	4.14	128.79	122.90
5	C	239	NAG	O5-C5-C6	3.75	113.08	107.20
5	G	238	NAG	O5-C5-C6	3.40	112.53	107.20
5	C	239	NAG	C3-C4-C5	-2.11	106.48	110.24

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	G	238	NAG	C3-C2-N2-C7
5	C	239	NAG	O5-C5-C6-O6
5	C	239	NAG	C4-C5-C6-O6
5	G	238	NAG	C4-C5-C6-O6
5	G	238	NAG	O5-C5-C6-O6
3	C	240	EDO	O1-C1-C2-O2
5	C	239	NAG	C8-C7-N2-C2
5	C	239	NAG	O7-C7-N2-C2
5	G	238	NAG	C8-C7-N2-C2
3	A	117	EDO	O1-C1-C2-O2
5	G	238	NAG	O7-C7-N2-C2
3	G	239	EDO	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	C	239	NAG	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	109/116 (93%)	0.47	8 (7%) 15 14	17, 29, 65, 69	0
1	B	114/116 (98%)	0.37	7 (6%) 21 21	14, 32, 54, 59	0
1	E	107/116 (92%)	0.32	8 (7%) 14 13	18, 28, 66, 71	0
1	F	114/116 (98%)	0.21	3 (2%) 56 58	16, 32, 52, 56	0
2	C	206/237 (86%)	0.50	17 (8%) 11 10	23, 46, 73, 83	0
2	D	213/237 (89%)	0.20	6 (2%) 53 55	16, 40, 65, 70	0
2	G	206/237 (86%)	0.42	14 (6%) 17 17	26, 45, 70, 80	0
2	H	211/237 (89%)	0.25	9 (4%) 35 37	20, 38, 66, 72	0
All	All	1280/1412 (90%)	0.34	72 (5%) 24 25	14, 37, 68, 83	0

All (72) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	48	ILE	5.4
1	B	49	ALA	5.2
1	B	7	LYS	5.1
1	B	8	VAL	4.8
1	A	1	GLY	4.7
2	G	92	ARG	4.4
2	C	218	ALA	4.3
2	C	92	ARG	4.3
1	F	1	GLY	4.3
1	A	47	HIS	4.1
2	C	76	VAL	4.1
2	H	48	LEU	4.0
2	H	8	PRO	4.0
2	G	117	ARG	3.9
2	C	13	TRP	3.9
1	B	48	ILE	3.8

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	E	1	GLY	3.6
1	A	2	LEU	3.5
2	H	7	ALA	3.5
1	F	49	ALA	3.4
2	C	37	ALA	3.3
2	G	86	MET	3.3
2	C	9	GLY	3.3
1	E	73	PRO	3.2
1	B	2	LEU	3.2
2	G	100	SER	3.1
1	E	76	ASN	3.1
1	F	48	ILE	3.1
2	G	26	LEU	3.1
2	D	8	PRO	3.0
2	G	72	SER	3.0
2	G	85	ARG	3.0
2	C	20	ALA	3.0
2	C	212	ARG	3.0
2	C	65	HIS	2.9
1	E	2	LEU	2.7
2	D	33	ALA	2.7
2	C	67	LEU	2.7
2	C	66	CYS	2.7
1	E	47	HIS	2.7
2	H	67	LEU	2.6
2	C	33	ALA	2.6
1	A	78	LYS	2.6
2	G	19	GLU	2.6
2	G	20	ALA	2.5
2	H	87	LEU	2.5
1	A	75	ALA	2.5
2	C	77	GLU	2.5
2	C	91	PRO	2.5
2	C	18	GLN	2.5
2	G	116	TYR	2.5
2	G	49	THR	2.4
2	H	218	ALA	2.4
2	C	100	SER	2.4
2	D	89	GLY	2.3
1	A	74	PHE	2.3
2	D	75	GLY	2.3
2	H	92	ARG	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	G	18	GLN	2.3
1	E	5	ASP	2.3
1	E	48	ILE	2.2
2	H	6	PRO	2.2
2	G	65	HIS	2.2
2	C	193	ILE	2.2
2	D	91	PRO	2.2
2	H	65	HIS	2.1
1	B	1	GLY	2.1
2	D	9	GLY	2.1
1	A	46	SER	2.1
2	G	91	PRO	2.1
1	B	6	GLY	2.1
1	E	78	LYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	NAG	G	238	14/15	0.69	0.17	76,82,83,83	0
5	NAG	C	239	14/15	0.78	0.16	73,79,81,81	0
3	EDO	C	240	4/4	0.86	0.25	62,65,66,67	0
3	EDO	A	117	4/4	0.88	0.28	60,60,61,62	0
3	EDO	G	239	4/4	0.89	0.29	72,73,73,74	0
4	SO4	C	238	5/5	0.93	0.17	78,79,80,80	0

## 6.5 Other polymers

There are no such residues in this entry.