



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 30, 2021 – 11:07 am BST

PDB ID : 7B80
Title : DeAMPylation complex of monomeric FICD and AMPylated BiP (state 2)
Authors : Perera, L.A.; Ron, D.
Deposited on : 2020-12-12
Resolution : 1.87 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.87 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	10282 (1.90-1.86)
Ramachandran outliers	138981	10152 (1.90-1.86)
Sidechain outliers	138945	10152 (1.90-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	B	523	
2	A	343	

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 7575 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Endoplasmic reticulum chaperone BiP.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	B	523	4103	2580	704	809	10	0	9	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	27	SER	-	expression tag	UNP G3I8R9
B	229	ALA	THR	engineered mutation	UNP G3I8R9
B	461	PHE	VAL	engineered mutation	UNP G3I8R9

- Molecule 2 is a protein called Protein adenylyltransferase FICD.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	A	340	2798	1776	486	522	14	0	7	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	103	SER	-	expression tag	UNP Q9BVA6
A	258	ASP	LEU	engineered mutation	UNP Q9BVA6
A	363	ALA	HIS	engineered mutation	UNP Q9BVA6

- Molecule 3 is ADENOSINE MONOPHOSPHATE (three-letter code: AMP) (formula: C₁₀H₁₄N₅O₇P) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	B	1	22	10	5	6	1	0	0

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
4	B	1	1	1	0	0
4	A	1	1	1	0	0

- Molecule 5 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total O P 5 4 1	0	0

- Molecule 6 is POTASSIUM ION (three-letter code: K) (formula: K).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	B	3	Total K 3 3	0	0

- Molecule 7 is 3,6,9,12,15,18-HEXA-OXAICOSANE-1,20-DIOL (three-letter code: P33) (formula: C₁₄H₃₀O₈).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			19	12	7		
7	A	1	Total	C	O	0	0
			19	12	7		

- Molecule 8 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			7	4	3		
8	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 9 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	B	325	Total	O	0	0
			325	325		
9	A	265	Total	O	0	0
			265	265		

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: Endoplasmic reticulum chaperone BiP



- Molecule 2: Protein adenylyltransferase FICD



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	95.00Å 103.89Å 104.79Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	52.40 – 1.87	Depositor
% Data completeness (in resolution range)	100.0 (52.40-1.87)	Depositor
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.201 , 0.228	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	7575	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: P33, AMP, MG, K, PEG, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	B	0.64	0/4165	0.73	0/5630
2	A	0.63	0/2856	0.69	0/3867
All	All	0.64	0/7021	0.71	0/9497

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	4103	0	4116	24	0
2	A	2798	0	2782	13	0
3	B	22	0	12	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	B	5	0	0	0	0
6	B	3	0	0	0	0
7	A	38	0	50	0	0
8	A	14	0	20	0	0
9	A	265	0	0	0	0
9	B	325	0	0	1	0
All	All	7575	0	6980	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:171:THR:HG22	1:B:199[A]:ILE:HG23	1.59	0.83
1:B:439[B]:ARG:HH11	1:B:439[B]:ARG:HG3	1.55	0.71
1:B:442[A]:VAL:HG21	2:A:127:LYS:HE3	1.77	0.65
1:B:442[B]:VAL:HG11	2:A:127:LYS:HE3	1.79	0.65
1:B:496:GLN:OE1	2:A:396:ARG:NH2	2.29	0.64
1:B:171:THR:HG22	1:B:199[A]:ILE:CG2	2.31	0.60
1:B:209:TYR:OH	1:B:358:GLU:HG3	2.09	0.53
1:B:361:LEU:O	1:B:388:ILE:HD11	2.09	0.51
1:B:414:LEU:C	1:B:414:LEU:HD13	2.31	0.50
1:B:340:LYS:HB2	1:B:341:PRO:HD3	1.93	0.49
1:B:442[B]:VAL:HG11	2:A:127:LYS:CE	2.42	0.48
2:A:287:ILE:HG13	2:A:335[B]:VAL:HG21	1.96	0.48
2:A:253:VAL:HG23	2:A:256:LYS:HB2	1.95	0.47
1:B:439[B]:ARG:HG3	1:B:439[B]:ARG:NH1	2.26	0.47
2:A:323:HIS:CG	2:A:324:PRO:HD2	2.50	0.46
1:B:461:PHE:HB2	1:B:484:LEU:HB3	1.98	0.45
1:B:426:ILE:HD12	1:B:463:ILE:HG23	1.99	0.45
2:A:218:VAL:HG21	2:A:432:ALA:HB1	1.99	0.44
1:B:484:LEU:C	1:B:484:LEU:HD23	2.38	0.44
1:B:442[A]:VAL:HG23	9:B:937:HOH:O	2.17	0.44
2:A:164:THR:O	2:A:168:THR:HG23	2.19	0.43
1:B:154:LYS:HD3	1:B:193:LEU:HD21	2.00	0.43
1:B:36:GLY:HA3	1:B:39:TYR:O	2.19	0.43
2:A:323:HIS:CD2	2:A:324:PRO:HD2	2.54	0.43
2:A:232:ALA:HB1	2:A:393:LYS:HD3	2.01	0.43
1:B:439[B]:ARG:NH1	1:B:439[B]:ARG:CG	2.81	0.42
1:B:443:VAL:HA	1:B:444:PRO:C	2.40	0.42
1:B:207:ILE:HD11	1:B:416:LEU:HD21	2.02	0.41
2:A:285:VAL:O	2:A:335[B]:VAL:HG13	2.19	0.41
2:A:396:ARG:HG3	2:A:400:TYR:CE2	2.55	0.41
1:B:498:GLU:O	1:B:513:ALA:HA	2.21	0.40
1:B:37:THR:HG21	1:B:228:GLY:HA3	2.03	0.40
1:B:171:THR:HA	1:B:199[A]:ILE:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	530/523 (101%)	518 (98%)	12 (2%)	0	100	100
2	A	345/343 (101%)	344 (100%)	1 (0%)	0	100	100
All	All	875/866 (101%)	862 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	447/443 (101%)	446 (100%)	1 (0%)	93	93
2	A	304/301 (101%)	304 (100%)	0	100	100
All	All	751/744 (101%)	750 (100%)	1 (0%)	93	93

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	176	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	302	GLN
2	A	323	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 5 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
8	PEG	A	504	-	6,6,6	0.17	0	5,5,5	0.06	0
5	PO4	B	603	4,6	4,4,4	0.65	0	6,6,6	0.41	0
7	P33	A	502	-	18,18,21	0.23	0	17,17,20	0.15	0
3	AMP	B	601	4,1	18,24,25	0.60	0	18,35,38	0.78	1 (5%)
7	P33	A	501	-	18,18,21	0.18	0	17,17,20	0.20	0
8	PEG	A	503	-	6,6,6	0.14	0	5,5,5	0.11	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	PEG	A	504	-	-	2/4/4/4	-
7	P33	A	502	-	-	9/16/16/19	-
3	AMP	B	601	4,1	-	2/3/25/26	0/3/3/3

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	P33	A	501	-	-	8/16/16/19	-
8	PEG	A	503	-	-	3/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	601	AMP	C5-C6-N6	2.27	123.81	120.35

There are no chirality outliers.


All (24) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	A	502	P33	O13-C14-C15-O16
7	A	501	P33	O10-C11-C12-O13
7	A	501	P33	O4-C5-C6-O7
7	A	502	P33	O10-C11-C12-O13
8	A	503	PEG	O1-C1-C2-O2
7	A	501	P33	O13-C14-C15-O16
7	A	501	P33	O16-C17-C18-O19
7	A	501	P33	O1-C2-C3-O4
7	A	502	P33	O16-C17-C18-O19
8	A	504	PEG	O2-C3-C4-O4
3	B	601	AMP	C3'-C4'-C5'-O5'
8	A	504	PEG	O1-C1-C2-O2
7	A	502	P33	O7-C8-C9-O10
7	A	502	P33	C18-C17-O16-C15
7	A	502	P33	C5-C6-O7-C8
8	A	503	PEG	C1-C2-O2-C3
7	A	501	P33	C6-C5-O4-C3
7	A	501	P33	C11-C12-O13-C14
8	A	503	PEG	C4-C3-O2-C2
7	A	502	P33	C6-C5-O4-C3
7	A	502	P33	O4-C5-C6-O7
7	A	501	P33	C15-C14-O13-C12
3	B	601	AMP	O4'-C4'-C5'-O5'
7	A	502	P33	C9-C8-O7-C6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.