



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 5, 2025 – 10:11 PM EST

PDB ID : 9BA9  
Title : O-GlcNAcase (OGA) inhibitor complex for the Treatment of Alzheimer's Disease  
Authors : Hendle, J.  
Deposited on : 2024-04-03  
Resolution : 2.75 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.21  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.004 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.40


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	1606 (2.78-2.74)
Clashscore	180529	1689 (2.78-2.74)
Ramachandran outliers	177936	1665 (2.78-2.74)
Sidechain outliers	177891	1665 (2.78-2.74)
RSRZ outliers	164620	1606 (2.78-2.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	535	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3710 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein O-GlcNAcase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	461	3657	2359	614	662	22	0	3	0

There are 130 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	53	SER	-	expression tag	UNP O60502
A	54	LEU	-	expression tag	UNP O60502
A	?	-	GLY	deletion	UNP O60502
A	?	-	THR	deletion	UNP O60502
A	?	-	PRO	deletion	UNP O60502
A	?	-	LEU	deletion	UNP O60502
A	?	-	VAL	deletion	UNP O60502
A	?	-	ALA	deletion	UNP O60502
A	?	-	ALA	deletion	UNP O60502
A	?	-	PRO	deletion	UNP O60502
A	?	-	SER	deletion	UNP O60502
A	?	-	LEU	deletion	UNP O60502
A	?	-	ASN	deletion	UNP O60502
A	?	-	ALA	deletion	UNP O60502
A	?	-	THR	deletion	UNP O60502
A	?	-	THR	deletion	UNP O60502
A	?	-	VAL	deletion	UNP O60502
A	?	-	VAL	deletion	UNP O60502
A	?	-	THR	deletion	UNP O60502
A	?	-	THR	deletion	UNP O60502
A	?	-	VAL	deletion	UNP O60502
A	?	-	TYR	deletion	UNP O60502
A	?	-	GLN	deletion	UNP O60502
A	?	-	GLU	deletion	UNP O60502
A	?	-	PRO	deletion	UNP O60502
A	?	-	ILE	deletion	UNP O60502
A	?	-	MET	deletion	UNP O60502

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	SER	deletion	UNP O60502
A	?	-	GLN	deletion	UNP O60502
A	?	-	GLY	deletion	UNP O60502
A	?	-	ALA	deletion	UNP O60502
A	?	-	ALA	deletion	UNP O60502
A	?	-	LEU	deletion	UNP O60502
A	?	-	SER	deletion	UNP O60502
A	?	-	GLY	deletion	UNP O60502
A	?	-	GLU	deletion	UNP O60502
A	?	-	PRO	deletion	UNP O60502
A	?	-	THR	deletion	UNP O60502
A	?	-	THR	deletion	UNP O60502
A	?	-	LEU	deletion	UNP O60502
A	?	-	THR	deletion	UNP O60502
A	?	-	LYS	deletion	UNP O60502
A	?	-	GLU	deletion	UNP O60502
A	?	-	GLU	deletion	UNP O60502
A	?	-	GLU	deletion	UNP O60502
A	?	-	LYS	deletion	UNP O60502
A	?	-	LYS	deletion	UNP O60502
A	?	-	GLN	deletion	UNP O60502
A	?	-	PRO	deletion	UNP O60502
A	?	-	ASP	deletion	UNP O60502
A	?	-	GLU	deletion	UNP O60502
A	?	-	GLU	deletion	UNP O60502
A	?	-	PRO	deletion	UNP O60502
A	?	-	MET	deletion	UNP O60502
A	?	-	ASP	deletion	UNP O60502
A	?	-	MET	deletion	UNP O60502
A	?	-	VAL	deletion	UNP O60502
A	?	-	VAL	deletion	UNP O60502
A	?	-	GLU	deletion	UNP O60502
A	?	-	LYS	deletion	UNP O60502
A	?	-	GLN	deletion	UNP O60502
A	?	-	GLU	deletion	UNP O60502
A	?	-	GLU	deletion	UNP O60502
A	?	-	THR	deletion	UNP O60502
A	?	-	ASP	deletion	UNP O60502
A	?	-	HIS	deletion	UNP O60502
A	?	-	LYS	deletion	UNP O60502
A	?	-	ASN	deletion	UNP O60502
A	?	-	ASP	deletion	UNP O60502

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	ASN	deletion	UNP O60502
A	?	-	GLN	deletion	UNP O60502
A	?	-	ILE	deletion	UNP O60502
A	?	-	LEU	deletion	UNP O60502
A	?	-	SER	deletion	UNP O60502
A	?	-	GLU	deletion	UNP O60502
A	?	-	ILE	deletion	UNP O60502
A	?	-	VAL	deletion	UNP O60502
A	?	-	GLU	deletion	UNP O60502
A	?	-	ALA	deletion	UNP O60502
A	?	-	LYS	deletion	UNP O60502
A	?	-	MET	deletion	UNP O60502
A	?	-	ALA	deletion	UNP O60502
A	?	-	GLU	deletion	UNP O60502
A	?	-	GLU	deletion	UNP O60502
A	?	-	LEU	deletion	UNP O60502
A	?	-	LYS	deletion	UNP O60502
A	?	-	PRO	deletion	UNP O60502
A	?	-	MET	deletion	UNP O60502
A	?	-	ASP	deletion	UNP O60502
A	?	-	THR	deletion	UNP O60502
A	?	-	ASP	deletion	UNP O60502
A	?	-	LYS	deletion	UNP O60502
A	?	-	GLU	deletion	UNP O60502
A	?	-	SER	deletion	UNP O60502
A	?	-	ILE	deletion	UNP O60502
A	?	-	ALA	deletion	UNP O60502
A	?	-	GLU	deletion	UNP O60502
A	?	-	SER	deletion	UNP O60502
A	?	-	LYS	deletion	UNP O60502
A	?	-	SER	deletion	UNP O60502
A	?	-	PRO	deletion	UNP O60502
A	?	-	GLU	deletion	UNP O60502
A	?	-	MET	deletion	UNP O60502
A	?	-	SER	deletion	UNP O60502
A	?	-	MET	deletion	UNP O60502
A	?	-	GLN	deletion	UNP O60502
A	?	-	GLU	deletion	UNP O60502
A	?	-	ASP	deletion	UNP O60502
A	?	-	CYS	deletion	UNP O60502
A	?	-	ILE	deletion	UNP O60502
A	?	-	SER	deletion	UNP O60502

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	ASP	deletion	UNP O60502
A	?	-	ILE	deletion	UNP O60502
A	?	-	ALA	deletion	UNP O60502
A	?	-	PRO	deletion	UNP O60502
A	?	-	MET	deletion	UNP O60502
A	?	-	GLN	deletion	UNP O60502
A	?	-	THR	deletion	UNP O60502
A	?	-	ASP	deletion	UNP O60502
A	?	-	GLU	deletion	UNP O60502
A	?	-	GLN	deletion	UNP O60502
A	?	-	THR	deletion	UNP O60502
A	?	-	ASN	deletion	UNP O60502
A	?	-	LYS	deletion	UNP O60502
A	?	-	GLU	deletion	UNP O60502
A	?	-	GLN	deletion	UNP O60502
A	?	-	PHE	deletion	UNP O60502
A	?	-	VAL	deletion	UNP O60502
A	596	SER	CYS	conflict	UNP O60502
A	663	GLU	CYS	conflict	UNP O60502

- Molecule 2 is N-{5-[(piperidin-1-yl)methyl]-1,3-thiazol-2-yl}acetamide (three-letter code: A1AKL) (formula: C<sub>11</sub>H<sub>17</sub>N<sub>3</sub>OS) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
2	A	1	16	11	3	1	1	0	0


- Molecule 3 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
3	A	37	Total	O	0	0
			37	37		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Protein O-GlcNAcase





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	133.26Å 133.26Å 114.24Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.75 30.00 – 2.75	Depositor EDS
% Data completeness (in resolution range)	99.3 (30.00-2.75) 99.3 (30.00-2.75)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.43 (at 2.77Å)	Xtrriage
Refinement program	REFMAC 5.7.0029	Depositor
R, $R_{free}$	0.217 , 0.260 0.223 , 0.266	Depositor DCC
$R_{free}$ test set	1377 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	70.7	Xtrriage
Anisotropy	0.202	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 51.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3710	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	77.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.16% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: A1AKL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.33	0/3762	0.50	0/5110

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3657	0	3483	29	0
2	A	16	0	0	1	0
3	A	37	0	0	0	0
All	All	3710	0	3483	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:104:ARG:HG3	1:A:138:SER:HB2	1.72	0.71

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:560:LEU:HD23	1:A:624:MET:HG2	1.84	0.60
1:A:267:VAL:O	1:A:271:ILE:HG12	2.02	0.58
1:A:567:LEU:HB3	1:A:568:PRO:HD2	1.85	0.57
1:A:304:ILE:HD12	1:A:335:ASN:HB3	1.86	0.57
1:A:264:ILE:HG13	1:A:303:LEU:HD22	1.87	0.57
1:A:227:ASN:HD21	1:A:230:GLN:HB2	1.70	0.56
1:A:682:ARG:H	1:A:691:ARG:HH12	1.55	0.55
1:A:335:ASN:OD1	1:A:338:GLY:HA3	2.09	0.53
1:A:647:ILE:O	1:A:651:MET:HG2	2.08	0.53
1:A:618:CYS:O	1:A:622:MET:HG2	2.10	0.51
2:A:801:A1AKL:S16	2:A:801:A1AKL:O15	2.69	0.51
1:A:545:LYS:HG3	1:A:546:PRO:HD2	1.95	0.48
1:A:660:TRP:CE2	1:A:671:PHE:HB2	2.49	0.48
1:A:296:TYR:CZ	1:A:299:ARG:HD3	2.50	0.46
1:A:551:GLU:HA	1:A:552:PRO:HD3	1.84	0.45
1:A:60:PHE:CD2	1:A:332:TYR:CE1	3.05	0.45
1:A:104:ARG:HG3	1:A:138:SER:CB	2.46	0.44
1:A:154:LEU:HD21	1:A:171:LEU:HD21	1.98	0.44
1:A:574:LYS:O	1:A:578:MET:HG3	2.17	0.44
1:A:67:GLY:HA2	1:A:96:ALA:O	2.18	0.44
1:A:384:LEU:HD22	1:A:558:LEU:HB3	2.00	0.44
1:A:151:VAL:HG12	1:A:155:LYS:HE3	2.00	0.44
1:A:161:VAL:HG12	1:A:166:CYS:HB3	2.01	0.42
1:A:264:ILE:HD13	1:A:264:ILE:HA	1.91	0.42
1:A:231:SER:HA	1:A:232:PRO:HD2	1.90	0.41
1:A:314:PRO:HB2	1:A:321:ASN:OD1	2.21	0.41
1:A:295:PRO:HB3	1:A:380:LEU:HB2	2.03	0.41
1:A:104:ARG:NH2	1:A:175:ASP:O	2.54	0.40
1:A:138:SER:OG	1:A:172:LEU:HB2	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	456/535 (85%)	427 (94%)	28 (6%)	1 (0%)	44 63

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	287	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	375/472 (79%)	366 (98%)	9 (2%)	44 65

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	65	VAL
1	A	71	ARG
1	A	101	TYR
1	A	278	TRP
1	A	286	TYR
1	A	290[A]	ARG
1	A	290[B]	ARG
1	A	624	MET
1	A	625	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	227	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	A1AKL	A	801	-	15,17,17	0.92	1 (6%)	13,22,22	1.07	2 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A1AKL	A	801	-	-	2/5/16/16	0/2/2/2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	801	A1AKL	C4-N14	2.88	1.41	1.36

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	801	A1AKL	C1-C2-S16	-2.27	109.74	112.00
2	A	801	A1AKL	O15-C4-N14	2.00	125.80	123.06

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	801	A1AKL	C2-C11-N13-C9
2	A	801	A1AKL	C2-C11-N13-C8

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	801	A1AKL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	461/535 (86%)	0.45	37 (8%) <span style="border: 1px solid red; padding: 2px;">20</span> <span style="border: 1px solid red; padding: 2px;">23</span>	38, 74, 120, 161	3 (0%)

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	370	THR	7.1
1	A	601	SER	6.2
1	A	594	VAL	5.9
1	A	393	VAL	5.7
1	A	394	PRO	5.5
1	A	679	TRP	5.4
1	A	287	ASP	5.2
1	A	680	ALA	4.7
1	A	341	LYS	4.6
1	A	59	ARG	4.5
1	A	669	ALA	4.3
1	A	539	ASP	4.2
1	A	697	GLY	4.1
1	A	681	PHE	4.1
1	A	226	PRO	3.9
1	A	670	GLN	3.9
1	A	604	ILE	3.9
1	A	225	TYR	3.9
1	A	590	SER	3.6
1	A	682	ARG	3.5
1	A	678	PRO	3.4
1	A	288	GLN	3.2
1	A	591	VAL	2.9
1	A	603	LYS	2.8
1	A	625	PHE	2.8
1	A	668	SER	2.8
1	A	663	GLU	2.7

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	262	GLU	2.6
1	A	683	GLY	2.6
1	A	607	TRP	2.5
1	A	378	MET	2.5
1	A	615	GLU	2.4
1	A	230	GLN	2.3
1	A	593	SER	2.2
1	A	664	ARG	2.2
1	A	128	GLU	2.1
1	A	661	LEU	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	A1AKL	A	801	16/16	0.97	0.09	57,60,64,68	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.