

# Full wwPDB X-ray Structure Validation Report (i)

Jan 27, 2024 – 12:29 PM EST

PDB ID : 1BD8

Title : STRUCTURE OF CDK INHIBITOR P19INK4D

Authors: Baumgartner, R.; Fernandez-Catalan, C.; Winoto, A.; Huber, R.; Engh, R.;

Holak, T.A.

Deposited on : 1998-05-12

Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Xtriage (Phenix) : NOT EXECUTED EDS : NOT EXECUTED

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\#  ext{Entries},  ext{ resolution range}( ext{Å}))$
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain			
1	A	156	70%	22%	5% •	



## 2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 1363 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called P19INK4D CDK4/6 INHIBITOR.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	A	156	Total 1168	C 721	N 222	O 222	S 3	0	0	0

• Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	195	Total O 195 195	14	0




## 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: P19INK4D CDK4/6 INHIBITOR





# 4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source	
Space group	P 1 21 1	Depositor	
Cell constants	28.47Å 54.56Å 45.59Å	Depositor	
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.16^{\circ}$ $90.00^{\circ}$	Depositor	
Resolution (Å)	(Not available) – 1.80	Depositor	
% Data completeness	(Not available) ((Not available)-1.80)	Depositor	
(in resolution range)	(1100 available) ((1100 available) 1.00)	Веровног	
$R_{merge}$	0.07	Depositor	
$R_{sym}$	(Not available)	Depositor	
Refinement program	REFMAC	Depositor	
$R, R_{free}$	0.190 , 0.260	Depositor	
Estimated twinning fraction	No twinning to report.	Xtriage	
Total number of atoms	1363	wwPDB-VP	
Average B, all atoms (Å <sup>2</sup> )	12.0	wwPDB-VP	



# 5 Model quality (i)

## 5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Mol Chain		nd lengths	Bond angles		
MIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	A	0.85	3/1185 (0.3%)	1.85	21/1605 (1.3%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintenain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

#### All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\mathring{\mathrm{A}})$	Ideal(Å)
1	A	162	MET	C-OXT	-20.50	0.84	1.23
1	A	23	GLU	CD-OE2	5.97	1.32	1.25
1	A	11	ARG	CD-NE	-5.23	1.37	1.46

#### All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
1	A	149	ARG	NE-CZ-NH2	-36.24	102.18	120.30
1	A	149	ARG	NE-CZ-NH1	20.35	130.47	120.30
1	A	11	ARG	CD-NE-CZ	19.21	150.49	123.60
1	A	149	ARG	CD-NE-CZ	14.42	143.79	123.60
1	A	145	LEU	CA-CB-CG	13.86	147.18	115.30
1	A	11	ARG	CG-CD-NE	11.54	136.04	111.80
1	A	146	ALA	CA-C-N	9.66	138.46	117.20
1	A	36	ASP	CB-CG-OD1	9.47	126.83	118.30
1	A	146	ALA	O-C-N	-8.90	108.45	122.70
1	A	80	ASP	CB-CG-OD1	8.00	125.50	118.30
1	A	11	ARG	NE-CZ-NH2	-7.87	116.36	120.30
1	A	147	LEU	N-CA-CB	7.72	125.85	110.40
1	A	36	ASP	CA-CB-CG	7.50	129.90	113.40

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	Atoms Z		$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
1	A	149	ARG	NH1-CZ-NH2	7.21	127.33	119.40
1	A	58	LEU	CB-CG-CD2	-7.01	99.08	111.00
1	A	10	ASP	CB-CG-OD1	6.29	123.96	118.30
1	A	149	ARG	CG-CD-NE	-5.95	99.31	111.80
1	A	11	ARG	NE-CZ-NH1	5.77	123.18	120.30
1	A	71	ASP	CB-CG-OD2	-5.72	113.15	118.30
1	A	26	ARG	NE-CZ-NH1	5.66	123.13	120.30
1	A	36	ASP	OD1-CG-OD2	-5.02	113.75	123.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	128	ALA	Mainchain

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1168	0	1171	37	5
2	A	195	0	0	18	11
All	All	1363	0	1171	37	11

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$egin{array}{ll}  ext{Interatomic} \  ext{distance } ( ext{Å}) \end{array}$	Clash overlap (Å)
1:A:128:ALA:O	1:A:129:GLU:HG3	1.75	0.86
1:A:43:LYS:HE3	2:A:323:HOH:O	1.81	0.79
1:A:84:THR:O	2:A:243:HOH:O	2.06	0.73
1:A:78:VAL:HG21	1:A:100:VAL:HG12	1.75	0.69
1:A:146:ALA:O	1:A:147:LEU:HB2	1.94	0.67
1:A:161:HIS:O	1:A:162:MET:HB2	1.96	0.63

Continued on next page...



 $Continued\ from\ previous\ page...$ 

A Land		Interatomic	Clash	
Atom-1	Atom-2	$\operatorname{distance}\ (\text{\AA})$	overlap (Å)	
1:A:133:HIS:HB3	2:A:252:HOH:O	2.00	0.61	
1:A:7:ARG:HG3	1:A:11:ARG:HD3	1.83	0.61	
1:A:129:GLU:OE1	2:A:331:HOH:O	2.16	0.60	
1:A:146:ALA:O	1:A:147:LEU:CB	2.47	0.59	
1:A:145:LEU:O	1:A:149:ARG:HG3	2.03	0.59	
1:A:11:ARG:NH1	2:A:171:HOH:O	2.36	0.58	
1:A:104:ASP:OD2	2:A:178:HOH:O	2.18	0.56	
1:A:109:LEU:HB3	2:A:163:HOH:O	2.06	0.56	
1:A:140:LEU:HB2	1:A:145:LEU:HD23	1.88	0.55	
1:A:145:LEU:HB3	2:A:349:HOH:O	2.06	0.54	
1:A:78:VAL:HG23	1:A:90:LEU:CD1	2.38	0.54	
1:A:87:LEU:HB2	2:A:332:HOH:O	2.07	0.53	
1:A:128:ALA:HB2	1:A:161:HIS:CE1	2.43	0.53	
1:A:77:PRO:HD3	2:A:172:HOH:O	2.09	0.52	
1:A:112:HIS:HB3	2:A:234:HOH:O	2.10	0.50	
1:A:138:ARG:HH11	1:A:138:ARG:HG2	1.76	0.50	
1:A:143:LEU:O	1:A:146:ALA:O	2.31	0.49	
1:A:149:ARG:HB3	1:A:149:ARG:CZ	2.41	0.47	
1:A:60:LEU:HD13	2:A:166:HOH:O	2.13	0.47	
1:A:71:ASP:OD2	2:A:324:HOH:O	2.20	0.47	
1:A:21:VAL:HG23	2:A:306:HOH:O	2.16	0.44	
1:A:76:SER:HA	2:A:172:HOH:O	2.18	0.44	
1:A:78:VAL:HG23	1:A:90:LEU:HD11	2.00	0.44	
1:A:26:ARG:HD3	2:A:253:HOH:O	2.18	0.43	
1:A:7:ARG:HH11	1:A:7:ARG:HD3	1.72	0.43	
1:A:58:LEU:HD23	1:A:59:GLU:N	2.34	0.42	
1:A:46:LEU:HB3	2:A:169:HOH:O	2.19	0.42	
1:A:34:HIS:CG	1:A:35:PRO:HD2	2.55	0.42	
1:A:161:HIS:O	1:A:162:MET:CB	2.65	0.41	
1:A:12:LEU:HA	1:A:27:LEU:HD23	2.02	0.41	
1:A:79:HIS:CE1	2:A:197:HOH:O	2.73	0.40	

All (11) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
2:A:187:HOH:O	2:A:238:HOH:O[1_455]	1.39	0.81
1:A:149:ARG:NH2	2:A:170:HOH:O[1_455]	1.49	0.71
2:A:188:HOH:O	2:A:274:HOH:O[1_655]	1.56	0.64
1:A:117:GLU:O	2:A:235:HOH:O[1_455]	1.87	0.33

Continued on next page...



Continued from previous page...

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
2:A:304:HOH:O	2:A:356:HOH:O[1_656]	1.95	0.25
1:A:91:LYS:NZ	2:A:248:HOH:O[2_656]	1.96	0.24
2:A:322:HOH:O	2:A:346:HOH:O[1_455]	1.99	0.21
1:A:149:ARG:CZ	2:A:170:HOH:O[1_455]	2.00	0.20
2:A:272:HOH:O	2:A:302:HOH:O[1_554]	2.06	0.14
2:A:207:HOH:O	2:A:250:HOH:O[1_655]	2.08	0.12
1:A:156:ASP:OD1	2:A:222:HOH:O[1_454]	2.15	0.05

## 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	154/156 (99%)	149 (97%)	3 (2%)	2 (1%)	12 3

#### All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	129	GLU
1	A	147	LEU

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	121/121 (100%)	112 (93%)	9 (7%)	13 4

All (9) residues with a non-rotameric sidechain are listed below:



Mol	Chain	Res	Type
1	A	7	ARG
1	A	11	ARG
1	A	27	LEU
1	A	50	MET
1	A	58	LEU
1	A	135	ARG
1	A	148	GLN
1	A	149	ARG
1	A	162	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	70	GLN
1	A	119	HIS
1	A	148	GLN
1	A	159	GLN
1	A	161	HIS

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry (i)

There are no ligands in this entry.

## 5.7 Other polymers (i)

There are no such residues in this entry.



## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

## 6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

## 6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

## 6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

## 6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

