



wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 16, 2023 – 08:50 am GMT

PDB ID : 4BD4
Title : Monomeric Human Cu,Zn Superoxide dismutase, loops IV and VII deleted, apo form, mutant H43F
Authors : Awad, W.; Saraboji, K.; Danielsson, J.; Lang, L.; Kurnik, M.; Marklund, S.L.; Oliveberg, M.; Logan, D.T.
Deposited on : 2012-10-04
Resolution : 2.78 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.78 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4107 (2.80-2.76)
Clashscore	141614	4575 (2.80-2.76)
Ramachandran outliers	138981	4487 (2.80-2.76)
Sidechain outliers	138945	4489 (2.80-2.76)
RSRZ outliers	127900	4027 (2.80-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	110	 70% 26% ..
1	B	110	 55% 41% ..
1	C	110	 56% 39% ..
1	D	110	 65% 25% 5% 3%
1	E	110	 59% 33% 5% 2%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	110	
1	G	110	
1	H	110	
1	I	110	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	H	1111	-	-	-	X

2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 6796 atoms, of which 32 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SUPEROXIDE DISMUTASE [CU-ZN].

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
1	A	109	764	478	136	150	0	0	0
1	B	109	765	478	136	151	0	0	0
1	C	108	760	475	135	150	0	0	0
1	D	104	742	465	131	146	0	0	0
1	E	104	738	464	130	144	0	0	0
1	F	102	732	460	129	143	0	0	0
1	G	107	755	472	134	149	0	0	0
1	H	100	719	452	127	140	0	0	0
1	I	98	712	448	124	140	0	0	0

There are 90 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	6	ALA	CYS	engineered mutation	UNP P00441
A	43	PHE	HIS	engineered mutation	UNP P00441
A	49	GLY	-	linker	PDB ?
A	50	ALA	-	linker	PDB ?
A	51	GLY	-	linker	PDB ?
A	81	SER	CYS	engineered mutation	UNP P00441
A	94	GLY	-	linker	PDB ?
A	95	ALA	-	linker	PDB ?
A	96	GLY	-	linker	PDB ?
A	103	SER	CYS	engineered mutation	UNP P00441
B	6	ALA	CYS	engineered mutation	UNP P00441

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	43	PHE	HIS	engineered mutation	UNP P00441
B	49	GLY	-	linker	PDB ?
B	50	ALA	-	linker	PDB ?
B	51	GLY	-	linker	PDB ?
B	81	SER	CYS	engineered mutation	UNP P00441
B	94	GLY	-	linker	PDB ?
B	95	ALA	-	linker	PDB ?
B	96	GLY	-	linker	PDB ?
B	103	SER	CYS	engineered mutation	UNP P00441
C	6	ALA	CYS	engineered mutation	UNP P00441
C	43	PHE	HIS	engineered mutation	UNP P00441
C	49	GLY	-	linker	PDB ?
C	50	ALA	-	linker	PDB ?
C	51	GLY	-	linker	PDB ?
C	81	SER	CYS	engineered mutation	UNP P00441
C	94	GLY	-	linker	PDB ?
C	95	ALA	-	linker	PDB ?
C	96	GLY	-	linker	PDB ?
C	103	SER	CYS	engineered mutation	UNP P00441
D	6	ALA	CYS	engineered mutation	UNP P00441
D	43	PHE	HIS	engineered mutation	UNP P00441
D	49	GLY	-	linker	PDB ?
D	50	ALA	-	linker	PDB ?
D	51	GLY	-	linker	PDB ?
D	81	SER	CYS	engineered mutation	UNP P00441
D	94	GLY	-	linker	PDB ?
D	95	ALA	-	linker	PDB ?
D	96	GLY	-	linker	PDB ?
D	103	SER	CYS	engineered mutation	UNP P00441
E	6	ALA	CYS	engineered mutation	UNP P00441
E	43	PHE	HIS	engineered mutation	UNP P00441
E	49	GLY	-	linker	PDB ?
E	50	ALA	-	linker	PDB ?
E	51	GLY	-	linker	PDB ?
E	81	SER	CYS	engineered mutation	UNP P00441
E	94	GLY	-	linker	PDB ?
E	95	ALA	-	linker	PDB ?
E	96	GLY	-	linker	PDB ?
E	103	SER	CYS	engineered mutation	UNP P00441
F	6	ALA	CYS	engineered mutation	UNP P00441
F	43	PHE	HIS	engineered mutation	UNP P00441
F	49	GLY	-	linker	PDB ?

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	50	ALA	-	linker	PDB ?
F	51	GLY	-	linker	PDB ?
F	81	SER	CYS	engineered mutation	UNP P00441
F	94	GLY	-	linker	PDB ?
F	95	ALA	-	linker	PDB ?
F	96	GLY	-	linker	PDB ?
F	103	SER	CYS	engineered mutation	UNP P00441
G	6	ALA	CYS	engineered mutation	UNP P00441
G	43	PHE	HIS	engineered mutation	UNP P00441
G	49	GLY	-	linker	PDB ?
G	50	ALA	-	linker	PDB ?
G	51	GLY	-	linker	PDB ?
G	81	SER	CYS	engineered mutation	UNP P00441
G	94	GLY	-	linker	PDB ?
G	95	ALA	-	linker	PDB ?
G	96	GLY	-	linker	PDB ?
G	103	SER	CYS	engineered mutation	UNP P00441
H	6	ALA	CYS	engineered mutation	UNP P00441
H	43	PHE	HIS	engineered mutation	UNP P00441
H	49	GLY	-	linker	PDB ?
H	50	ALA	-	linker	PDB ?
H	51	GLY	-	linker	PDB ?
H	81	SER	CYS	engineered mutation	UNP P00441
H	94	GLY	-	linker	PDB ?
H	95	ALA	-	linker	PDB ?
H	96	GLY	-	linker	PDB ?
H	103	SER	CYS	engineered mutation	UNP P00441
I	6	ALA	CYS	engineered mutation	UNP P00441
I	43	PHE	HIS	engineered mutation	UNP P00441
I	49	GLY	-	linker	PDB ?
I	50	ALA	-	linker	PDB ?
I	51	GLY	-	linker	PDB ?
I	81	SER	CYS	engineered mutation	UNP P00441
I	94	GLY	-	linker	PDB ?
I	95	ALA	-	linker	PDB ?
I	96	GLY	-	linker	PDB ?
I	103	SER	CYS	engineered mutation	UNP P00441

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	O	0	0
			14	3	8	3		
2	B	1	Total	C	H	O	0	0
			14	3	8	3		
2	G	1	Total	C	H	O	0	0
			14	3	8	3		
2	H	1	Total	C	H	O	0	0
			14	3	8	3		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	15	Total	O	0	0
			15	15		
3	B	17	Total	O	0	0
			17	17		
3	C	4	Total	O	0	0
			4	4		
3	D	6	Total	O	0	0
			6	6		
3	E	3	Total	O	0	0
			3	3		
3	F	1	Total	O	0	0
			1	1		
3	G	2	Total	O	0	0
			2	2		
3	H	2	Total	O	0	0
			2	2		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	I	3	Total	O	0	0
			3	3		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: SUPEROXIDE DISMUTASE [CU-ZN]

Chain A: 




- Molecule 1: SUPEROXIDE DISMUTASE [CU-ZN]

Chain B: 




- Molecule 1: SUPEROXIDE DISMUTASE [CU-ZN]

Chain C: 




- Molecule 1: SUPEROXIDE DISMUTASE [CU-ZN]

Chain D: 





- Molecule 1: SUPEROXIDE DISMUTASE [CU-ZN]

Chain E: 2% 59% 33% 5%




- Molecule 1: SUPEROXIDE DISMUTASE [CU-ZN]

Chain F: 4% 53% 35% 5% 7%



- Molecule 1: SUPEROXIDE DISMUTASE [CU-ZN]

Chain G: 2% 50% 45% 2% 1%




- Molecule 1: SUPEROXIDE DISMUTASE [CU-ZN]

Chain H: 7% 62% 25% 9%



- Molecule 1: SUPEROXIDE DISMUTASE [CU-ZN]

Chain I: 42% 43% 5% 11%



I74	S75	I82	I83	L87	V88	V89	H90	E91	LYS	ALA	ALA	GLY	ALA	GLY	ALA	GLY	ALA	GLY	ALA	GLY	S99	A102	S103	I106	G107	I108	A109	GLN
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	------	------	------	------	------	------	-----

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	235.03Å 49.89Å 97.94Å 90.00° 97.19° 90.00°	Depositor
Resolution (Å)	29.15 – 2.78 29.15 – 2.78	Depositor EDS
% Data completeness (in resolution range)	98.1 (29.15-2.78) 98.2 (29.15-2.78)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.99 (at 2.76Å)	Xtrriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.180 , 0.246 0.192 , 0.252	Depositor DCC
R_{free} test set	1437 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	69.6	Xtrriage
Anisotropy	0.510	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 50.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6796	wwPDB-VP
Average B, all atoms (Å ²)	85.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.37% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.52	0/774	0.70	0/1045
1	B	0.55	0/775	0.72	0/1046
1	C	0.46	0/770	0.67	0/1039
1	D	0.50	0/751	0.60	0/1012
1	E	0.42	0/746	0.63	0/1004
1	F	0.38	0/741	0.61	0/999
1	G	0.39	0/765	0.64	0/1032
1	H	0.39	0/727	0.60	0/980
1	I	0.38	0/720	0.57	0/971
All	All	0.45	0/6769	0.64	0/9128

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	764	0	771	31	0
1	B	765	0	771	63	0
1	C	760	0	763	52	1
1	D	742	0	749	36	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	738	0	747	36	0
1	F	732	0	741	42	0
1	G	755	0	758	54	1
1	H	719	0	726	28	0
1	I	712	0	718	67	0
2	A	6	8	8	1	0
2	B	6	8	8	0	0
2	G	6	8	8	0	0
2	H	6	8	8	0	0
3	A	15	0	0	0	0
3	B	17	0	0	5	0
3	C	4	0	0	0	0
3	D	6	0	0	0	0
3	E	3	0	0	0	0
3	F	1	0	0	0	0
3	G	2	0	0	0	0
3	H	2	0	0	0	0
3	I	3	0	0	1	0
All	All	6764	32	6776	385	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 29.

The worst 5 of 385 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:42:LEU:HD11	1:F:56:ASN:HB3	1.36	1.07
1:G:60:ASP:HB3	1:G:62:ASP:H	1.18	1.07
1:I:22:GLN:HB2	1:I:29:VAL:HG22	1.33	1.05
1:I:74:ILE:HB	1:I:82:ILE:HD11	1.42	1.02
1:F:42:LEU:HD13	1:F:58:THR:HG22	1.40	1.01

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:24:GLU:OE2	1:G:26:ASN:ND2[2_556]	1.99	0.21

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	107/110 (97%)	104 (97%)	3 (3%)	0	100	100
1	B	107/110 (97%)	103 (96%)	4 (4%)	0	100	100
1	C	106/110 (96%)	103 (97%)	3 (3%)	0	100	100
1	D	100/110 (91%)	96 (96%)	4 (4%)	0	100	100
1	E	98/110 (89%)	96 (98%)	2 (2%)	0	100	100
1	F	98/110 (89%)	98 (100%)	0	0	100	100
1	G	105/110 (96%)	100 (95%)	5 (5%)	0	100	100
1	H	94/110 (86%)	92 (98%)	2 (2%)	0	100	100
1	I	92/110 (84%)	91 (99%)	1 (1%)	0	100	100
All	All	907/990 (92%)	883 (97%)	24 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	78/79 (99%)	74 (95%)	4 (5%)	24	53
1	B	78/79 (99%)	74 (95%)	4 (5%)	24	53
1	C	78/79 (99%)	75 (96%)	3 (4%)	33	64
1	D	78/79 (99%)	73 (94%)	5 (6%)	17	42
1	E	77/79 (98%)	69 (90%)	8 (10%)	7	19

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	F	77/79 (98%)	69 (90%)	8 (10%)	7	19
1	G	78/79 (99%)	74 (95%)	4 (5%)	24	53
1	H	76/79 (96%)	71 (93%)	5 (7%)	16	41
1	I	77/79 (98%)	70 (91%)	7 (9%)	9	25
All	All	697/711 (98%)	649 (93%)	48 (7%)	15	38

5 of 48 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	89	VAL
1	G	99	SER
1	F	100	ARG
1	G	17	ILE
1	H	35	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	80	HIS
1	E	80	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	A	1110	-	5,5,5	0.37	0	5,5,5	0.21	0
2	GOL	G	1109	-	5,5,5	0.36	0	5,5,5	0.28	0
2	GOL	H	1111	-	5,5,5	0.37	0	5,5,5	0.33	0
2	GOL	B	1110	-	5,5,5	0.37	0	5,5,5	0.33	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	1110	-	-	3/4/4/4	-
2	GOL	G	1109	-	-	2/4/4/4	-
2	GOL	H	1111	-	-	2/4/4/4	-
2	GOL	B	1110	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 9 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1110	GOL	C1-C2-C3-O3
2	A	1110	GOL	C1-C2-C3-O3
2	H	1111	GOL	C1-C2-C3-O3
2	A	1110	GOL	O2-C2-C3-O3
2	B	1110	GOL	O2-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1110	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	109/110 (99%)	-0.34	0 100 100	43, 62, 101, 108	0
1	B	109/110 (99%)	-0.33	0 100 100	37, 58, 93, 141	0
1	C	108/110 (98%)	-0.12	1 (0%) 84 82	51, 83, 121, 162	1 (0%)
1	D	104/110 (94%)	-0.12	3 (2%) 51 46	60, 77, 112, 133	0
1	E	104/110 (94%)	-0.05	2 (1%) 66 63	47, 79, 125, 167	0
1	F	102/110 (92%)	0.19	4 (3%) 39 34	57, 93, 149, 188	0
1	G	107/110 (97%)	-0.03	2 (1%) 66 63	58, 94, 130, 177	0
1	H	100/110 (90%)	0.17	8 (8%) 12 8	54, 84, 150, 190	0
1	I	98/110 (89%)	0.04	1 (1%) 82 80	54, 97, 144, 159	0
All	All	941/990 (95%)	-0.07	21 (2%) 62 57	37, 80, 133, 190	1 (0%)

The worst 5 of 21 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	101	LEU	4.6
1	D	1	ALA	4.6
1	E	97	ALA	4.2
1	I	1	ALA	4.1
1	H	39	THR	3.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GOL	H	1111	6/6	0.59	0.66	113,135,159,159	0
2	GOL	G	1109	6/6	0.65	0.39	143,172,178,179	0
2	GOL	A	1110	6/6	0.85	0.33	104,126,151,151	0
2	GOL	B	1110	6/6	0.95	0.36	105,127,133,134	0

6.5 Other polymers [i](#)

There are no such residues in this entry.