



wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 22, 2022 – 06:32 pm GMT

PDB ID : 8BET
Title : Structure of D188A-fructofuranosidase from *Rhodotorula dairenensis* in complex with sucrose
Authors : Jimenez-Ortega, E.; Sanz-Aparicio, J.
Deposited on : 2022-10-21
Resolution : 2.38 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.31.3
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.3


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.38 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.













Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5509 (2.40-2.36)
Clashscore	141614	6082 (2.40-2.36)
Ramachandran outliers	138981	5973 (2.40-2.36)
Sidechain outliers	138945	5975 (2.40-2.36)
RSRZ outliers	127900	5397 (2.40-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	675	 2% 73% 5% 22%
1	B	675	 5% 72% 5% 22%
1	C	675	 2% 73% 5% 22%
1	D	675	 7% 72% 5% 22%
2	E	2	 100%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	H	2	 100%
2	K	2	 100%
3	F	3	 100%
3	I	3	 67% 33%
3	J	3	 100%
4	G	6	 33% 67%
5	L	2	 100%
5	M	2	 100%
5	N	2	 50% 50%
5	O	2	 50% 50%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	MAN	B	702	-	-	-	X
6	MAN	C	703	-	-	-	X
6	MAN	D	701	-	-	-	X
7	NAG	B	705	-	-	-	X

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 17723 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-fructofuranosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	525	4032	2569	656	796	11	0	0	0
1	B	526	4038	2572	657	798	11	0	0	0
1	C	528	4050	2578	659	802	11	0	0	0
1	D	524	4026	2566	655	794	11	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	188	ALA	ASP	engineered mutation	UNP A0A856TAI5
B	188	ALA	ASP	engineered mutation	UNP A0A856TAI5
C	188	ALA	ASP	engineered mutation	UNP A0A856TAI5
D	188	ALA	ASP	engineered mutation	UNP A0A856TAI5

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	E	2	28	16	2	10	0	0	0
2	H	2	28	16	2	10	0	0	0
2	K	2	28	16	2	10	0	0	0

- Molecule 3 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	F	3	39	22	2	15	0	0	0
3	I	3	39	22	2	15	0	0	0
3	J	3	39	22	2	15	0	0	0

- Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	G	6	72	40	2	30	0	0	0

- Molecule 5 is an oligosaccharide called beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
5	L	2	23	12	11	0	0	0
5	M	2	23	12	11	0	0	0
5	N	2	23	12	11	0	0	0
5	O	2	23	12	11	0	0	0

- Molecule 6 is alpha-D-mannopyranose (three-letter code: MAN) (formula: C₆H₁₂O₆).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf		
6	A	1	Total	C	O	0	0
			11	6	5		
6	A	1	Total	C	O	0	0
			11	6	5		
6	A	1	Total	C	O	0	0
			11	6	5		
6	A	1	Total	C	O	0	0
			11	6	5		
6	B	1	Total	C	O	0	0
			11	6	5		
6	B	1	Total	C	O	0	0
			11	6	5		
6	B	1	Total	C	O	0	0
			11	6	5		
6	C	1	Total	C	O	0	0
			11	6	5		
6	C	1	Total	C	O	0	0
			11	6	5		
6	C	1	Total	C	O	0	0
			11	6	5		
6	C	1	Total	C	O	0	0
			11	6	5		
6	D	1	Total	C	O	0	0
			11	6	5		
6	D	1	Total	C	O	0	0
			11	6	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	D	1	Total	C	O	0	0
			11	6	5		

- Molecule 7 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	C	N	O	0	0
			14	8	1	5		
7	A	1	Total	C	N	O	0	0
			14	8	1	5		
7	A	1	Total	C	N	O	0	0
			14	8	1	5		
7	A	1	Total	C	N	O	0	0
			14	8	1	5		
7	A	1	Total	C	N	O	0	0
			14	8	1	5		
7	A	1	Total	C	N	O	0	0
			14	8	1	5		
7	B	1	Total	C	N	O	0	0
			14	8	1	5		
7	B	1	Total	C	N	O	0	0
			14	8	1	5		
7	B	1	Total	C	N	O	0	0
			14	8	1	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	B	1	Total	C	N	O	0	0
			14	8	1	5		
7	B	1	Total	C	N	O	0	0
			14	8	1	5		
7	B	1	Total	C	N	O	0	0
			14	8	1	5		
7	B	1	Total	C	N	O	0	0
			14	8	1	5		
7	C	1	Total	C	N	O	0	0
			14	8	1	5		
7	C	1	Total	C	N	O	0	0
			14	8	1	5		
7	C	1	Total	C	N	O	0	0
			14	8	1	5		
7	C	1	Total	C	N	O	0	0
			14	8	1	5		
7	C	1	Total	C	N	O	0	0
			14	8	1	5		
7	D	1	Total	C	N	O	0	0
			14	8	1	5		
7	D	1	Total	C	N	O	0	0
			14	8	1	5		
7	D	1	Total	C	N	O	0	0
			14	8	1	5		
7	D	1	Total	C	N	O	0	0
			14	8	1	5		
7	D	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	200	Total	O	0	0
			200	200		
8	B	176	Total	O	0	0
			176	176		
8	C	166	Total	O	0	0
			166	166		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	D	152	Total 152	O 152	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Beta-fructofuranosidase




- Molecule 1: Beta-fructofuranosidase





- Molecule 1: Beta-fructofuranosidase





• Molecule 1: Beta-fructofuranosidase




• Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose




MAG1
MAG2

• Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose




MAG1
MAG2

• Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain K:  100%


NAG1
NAG2

- Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  100%


NAG1
NAG2
BMA3

- Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I:  67% 33%

NAG1
NAG2
BMA3

- Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain J:  100%

NAG1
NAG2
BMA3

- Molecule 4: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:  33% 67%


NAG3
NAG2
BMA3
MAN4
MAN5
MAN6

- Molecule 5: beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose

Chain L:  100%

GLC1
FRU2

- Molecule 5: beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose

Chain M:  100%

GLC1
FRU2

- Molecule 5: beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose

Chain N:  50% 50%

 GLC1
FRU2

- Molecule 5: beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose

Chain O:  50% 50%

 GLC1
FRU2

4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	80.34Å 112.81Å 139.08Å 90.00° 104.74° 90.00°	Depositor
Resolution (Å)	45.69 – 2.38 45.65 – 2.38	Depositor EDS
% Data completeness (in resolution range)	99.8 (45.69-2.38) 99.8 (45.65-2.38)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.31 (at 2.37Å)	Xtrriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.184 , 0.222 0.186 , 0.224	Depositor DCC
R_{free} test set	4728 reflections (4.92%)	wwPDB-VP
Wilson B-factor (Å ²)	36.8	Xtrriage
Anisotropy	0.184	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 27.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.021 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	17723	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, GLC, FRU, BMA, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.65	0/4160	0.75	0/5721
1	B	0.65	0/4166	0.75	0/5729
1	C	0.66	0/4178	0.75	0/5745
1	D	0.64	0/4154	0.74	0/5713
All	All	0.65	0/16658	0.75	0/22908

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4032	0	3751	21	0
1	B	4038	0	3757	29	0
1	C	4050	0	3765	25	0
1	D	4026	0	3747	27	0
2	E	28	0	25	0	0
2	H	28	0	25	0	0
2	K	28	0	25	0	0
3	F	39	0	34	0	0
3	I	39	0	34	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	J	39	0	34	0	0
4	G	72	0	61	0	0
5	L	23	0	21	1	0
5	M	23	0	21	0	0
5	N	23	0	21	1	0
5	O	23	0	21	1	0
6	A	44	0	40	0	0
6	B	33	0	30	0	0
6	C	44	0	40	2	0
6	D	33	0	30	2	0
7	A	98	0	91	2	0
7	B	98	0	91	3	0
7	C	84	0	78	1	0
7	D	84	0	78	4	0
8	A	200	0	0	5	0
8	B	176	0	0	0	0
8	C	166	0	0	0	0
8	D	152	0	0	1	0
All	All	17723	0	15820	102	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 102 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:487:PRO:HB2	1:B:490:LEU:HD11	1.14	1.12
1:B:487:PRO:HB2	1:B:490:LEU:CD1	1.96	0.95
1:C:467:LEU:HD12	1:C:625:GLU:OE1	1.80	0.80
1:B:467:LEU:HD12	1:B:625:GLU:OE1	1.81	0.80
1:D:467:LEU:HD12	1:D:625:GLU:OE1	1.81	0.79

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	521/675 (77%)	501 (96%)	18 (4%)	2 (0%)	34	46
1	B	522/675 (77%)	501 (96%)	19 (4%)	2 (0%)	34	46
1	C	524/675 (78%)	502 (96%)	20 (4%)	2 (0%)	34	46
1	D	520/675 (77%)	503 (97%)	15 (3%)	2 (0%)	34	46
All	All	2087/2700 (77%)	2007 (96%)	72 (3%)	8 (0%)	34	46

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	463	SER
1	C	463	SER
1	A	463	SER
1	A	489	ASP
1	B	489	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	441/545 (81%)	441 (100%)	0	100	100
1	B	442/545 (81%)	442 (100%)	0	100	100
1	C	444/545 (82%)	444 (100%)	0	100	100
1	D	440/545 (81%)	440 (100%)	0	100	100
All	All	1767/2180 (81%)	1767 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	518	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	664	GLN
1	D	666	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

29 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	E	1	2,1	14,14,15	0.79	0	17,19,21	0.95	1 (5%)
2	NAG	E	2	2	14,14,15	0.60	0	17,19,21	1.19	2 (11%)
3	NAG	F	1	1,3	14,14,15	0.64	0	17,19,21	1.11	1 (5%)
3	NAG	F	2	3	14,14,15	0.64	0	17,19,21	1.03	1 (5%)
3	BMA	F	3	3	11,11,12	0.71	0	15,15,17	1.09	1 (6%)
4	NAG	G	1	4,1	14,14,15	0.51	0	17,19,21	1.00	0
4	NAG	G	2	4	14,14,15	0.36	0	17,19,21	0.94	0
4	BMA	G	3	4	11,11,12	0.61	0	15,15,17	0.99	1 (6%)
4	MAN	G	4	4	11,11,12	0.50	0	15,15,17	1.09	1 (6%)
4	MAN	G	5	4	11,11,12	0.58	0	15,15,17	1.24	2 (13%)
4	MAN	G	6	4	11,11,12	0.84	0	15,15,17	1.55	3 (20%)
2	NAG	H	1	2,1	14,14,15	0.76	0	17,19,21	1.09	2 (11%)
2	NAG	H	2	2	14,14,15	0.81	0	17,19,21	1.40	1 (5%)
3	NAG	I	1	1,3	14,14,15	0.51	0	17,19,21	1.04	2 (11%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	I	2	3	14,14,15	0.56	0	17,19,21	1.29	2 (11%)
3	BMA	I	3	3	11,11,12	0.62	0	15,15,17	1.12	1 (6%)
3	NAG	J	1	1,3	14,14,15	0.49	0	17,19,21	1.28	2 (11%)
3	NAG	J	2	3	14,14,15	0.54	0	17,19,21	1.12	2 (11%)
3	BMA	J	3	3	11,11,12	0.64	0	15,15,17	0.91	1 (6%)
2	NAG	K	1	2,1	14,14,15	0.57	0	17,19,21	1.20	1 (5%)
2	NAG	K	2	2	14,14,15	0.49	0	17,19,21	1.01	1 (5%)
5	GLC	L	1	5	11,11,12	0.73	0	15,15,17	1.04	0
5	FRU	L	2	5	11,12,12	0.53	0	10,18,18	0.89	1 (10%)
5	GLC	M	1	5	11,11,12	0.69	0	15,15,17	1.57	5 (33%)
5	FRU	M	2	5	11,12,12	0.75	1 (9%)	10,18,18	0.77	0
5	GLC	N	1	5	11,11,12	0.81	1 (9%)	15,15,17	0.99	2 (13%)
5	FRU	N	2	5	11,12,12	0.81	0	10,18,18	0.70	0
5	GLC	O	1	5	11,11,12	0.54	0	15,15,17	1.93	6 (40%)
5	FRU	O	2	5	11,12,12	0.62	0	10,18,18	0.97	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	E	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	E	2	2	-	0/6/23/26	0/1/1/1
3	NAG	F	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	F	2	3	-	2/6/23/26	0/1/1/1
3	BMA	F	3	3	-	2/2/19/22	0/1/1/1
4	NAG	G	1	4,1	-	0/6/23/26	0/1/1/1
4	NAG	G	2	4	-	1/6/23/26	0/1/1/1
4	BMA	G	3	4	-	2/2/19/22	0/1/1/1
4	MAN	G	4	4	-	1/2/19/22	0/1/1/1
4	MAN	G	5	4	-	2/2/19/22	0/1/1/1
4	MAN	G	6	4	-	0/2/19/22	0/1/1/1
2	NAG	H	1	2,1	-	2/6/23/26	0/1/1/1
2	NAG	H	2	2	-	0/6/23/26	0/1/1/1
3	NAG	I	1	1,3	-	4/6/23/26	0/1/1/1
3	NAG	I	2	3	-	0/6/23/26	0/1/1/1
3	BMA	I	3	3	-	2/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	J	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	J	2	3	-	2/6/23/26	0/1/1/1
3	BMA	J	3	3	-	2/2/19/22	0/1/1/1
2	NAG	K	1	2,1	-	2/6/23/26	0/1/1/1
2	NAG	K	2	2	-	0/6/23/26	0/1/1/1
5	GLC	L	1	5	-	0/2/19/22	0/1/1/1
5	FRU	L	2	5	-	1/5/24/24	0/1/1/1
5	GLC	M	1	5	-	2/2/19/22	0/1/1/1
5	FRU	M	2	5	-	3/5/24/24	0/1/1/1
5	GLC	N	1	5	-	0/2/19/22	0/1/1/1
5	FRU	N	2	5	-	3/5/24/24	0/1/1/1
5	GLC	O	1	5	-	2/2/19/22	0/1/1/1
5	FRU	O	2	5	-	1/5/24/24	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	N	1	GLC	O5-C1	-2.07	1.40	1.43
5	M	2	FRU	O5-C2	-2.05	1.40	1.43

The worst 5 of 42 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	2	NAG	C1-O5-C5	4.33	118.05	112.19
5	O	1	GLC	O3-C3-C4	-3.47	102.32	110.35
3	J	1	NAG	O5-C1-C2	-3.34	106.01	111.29
2	K	2	NAG	C1-O5-C5	3.26	116.61	112.19
3	I	2	NAG	C1-O5-C5	3.14	116.44	112.19

There are no chirality outliers.

5 of 36 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	M	2	FRU	O1-C1-C2-C3
5	M	2	FRU	O1-C1-C2-O2
5	M	2	FRU	O1-C1-C2-O5
5	N	2	FRU	O1-C1-C2-O2
5	N	2	FRU	O1-C1-C2-O5


There are no ring outliers.


4 monomers are involved in 5 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	I	1	NAG	2	0
5	L	1	GLC	1	0
5	O	1	GLC	1	0
5	N	1	GLC	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.




















5.6 Ligand geometry

40 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	MAN	C	703	1	11,11,12	2.60	6 (54%)	15,15,17	1.77	3 (20%)
7	NAG	D	708	1	14,14,15	0.62	0	17,19,21	1.31	2 (11%)
6	MAN	B	702	1	11,11,12	2.28	4 (36%)	15,15,17	1.68	3 (20%)
7	NAG	A	705	1	14,14,15	0.52	0	17,19,21	1.07	1 (5%)
7	NAG	D	707	1	14,14,15	0.39	0	17,19,21	1.41	3 (17%)
7	NAG	C	705	1	14,14,15	0.64	0	17,19,21	1.18	2 (11%)
6	MAN	D	702	1	11,11,12	0.50	0	15,15,17	1.16	1 (6%)
6	MAN	C	702	1	11,11,12	0.52	0	15,15,17	1.91	4 (26%)
7	NAG	B	707	1	14,14,15	0.55	0	17,19,21	1.19	1 (5%)
6	MAN	C	704	1	11,11,12	0.61	0	15,15,17	0.74	0
7	NAG	A	706	1	14,14,15	0.41	0	17,19,21	1.68	2 (11%)
7	NAG	B	705	1	14,14,15	0.95	1 (7%)	17,19,21	1.43	3 (17%)
6	MAN	B	703	1	11,11,12	0.56	0	15,15,17	1.29	1 (6%)
6	MAN	A	702	1	11,11,12	0.75	0	15,15,17	1.30	3 (20%)
7	NAG	B	709	1	14,14,15	0.35	0	17,19,21	0.76	0
7	NAG	A	711	1	14,14,15	0.78	0	17,19,21	1.48	4 (23%)
7	NAG	C	706	1	14,14,15	0.37	0	17,19,21	1.23	1 (5%)
6	MAN	A	703	1	11,11,12	0.87	0	15,15,17	1.48	1 (6%)
7	NAG	A	709	1	14,14,15	0.74	0	17,19,21	1.19	2 (11%)
7	NAG	B	706	1	14,14,15	0.60	0	17,19,21	1.65	4 (23%)
6	MAN	D	701	1	11,11,12	0.74	0	15,15,17	1.43	2 (13%)
7	NAG	B	704	1	14,14,15	0.70	0	17,19,21	1.46	1 (5%)
7	NAG	C	708	1	14,14,15	0.48	0	17,19,21	0.98	0
7	NAG	C	710	1	14,14,15	0.48	0	17,19,21	1.50	3 (17%)
7	NAG	D	704	1	14,14,15	0.60	0	17,19,21	1.34	1 (5%)
7	NAG	C	707	1	14,14,15	0.73	1 (7%)	17,19,21	0.87	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	MAN	B	701	1	11,11,12	0.63	0	15,15,17	1.65	2 (13%)
7	NAG	A	708	1	14,14,15	0.90	1 (7%)	17,19,21	0.94	1 (5%)
7	NAG	B	710	1	14,14,15	0.64	0	17,19,21	1.33	2 (11%)
7	NAG	D	705	1	14,14,15	0.72	0	17,19,21	2.43	6 (35%)
7	NAG	B	708	1	14,14,15	0.83	0	17,19,21	0.97	0
7	NAG	D	706	1	14,14,15	0.48	0	17,19,21	1.09	2 (11%)
6	MAN	A	704	1	11,11,12	0.64	0	15,15,17	1.62	2 (13%)
6	MAN	D	703	1	11,11,12	0.95	0	15,15,17	1.38	2 (13%)
7	NAG	C	709	1	14,14,15	0.67	0	17,19,21	1.25	1 (5%)
7	NAG	A	707	1	14,14,15	0.67	1 (7%)	17,19,21	0.79	0
6	MAN	C	701	1	11,11,12	0.69	0	15,15,17	1.52	3 (20%)
6	MAN	A	701	1	11,11,12	0.63	0	15,15,17	1.02	0
7	NAG	A	710	1	14,14,15	0.64	0	17,19,21	1.12	0
7	NAG	D	709	1	14,14,15	0.52	0	17,19,21	0.94	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	MAN	C	703	1	-	1/2/19/22	0/1/1/1
7	NAG	D	708	1	-	2/6/23/26	0/1/1/1
6	MAN	B	702	1	-	2/2/19/22	0/1/1/1
7	NAG	A	705	1	-	2/6/23/26	0/1/1/1
7	NAG	D	707	1	-	2/6/23/26	0/1/1/1
7	NAG	C	705	1	-	4/6/23/26	0/1/1/1
6	MAN	D	702	1	-	1/2/19/22	0/1/1/1
6	MAN	C	702	1	-	2/2/19/22	0/1/1/1
7	NAG	B	707	1	-	2/6/23/26	0/1/1/1
6	MAN	C	704	1	-	0/2/19/22	0/1/1/1
7	NAG	A	706	1	-	2/6/23/26	0/1/1/1
7	NAG	B	705	1	-	4/6/23/26	0/1/1/1
6	MAN	B	703	1	-	2/2/19/22	0/1/1/1
6	MAN	A	702	1	-	2/2/19/22	0/1/1/1
7	NAG	B	709	1	-	0/6/23/26	0/1/1/1
7	NAG	A	711	1	-	2/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	NAG	C	706	1	-	2/6/23/26	0/1/1/1
6	MAN	A	703	1	-	1/2/19/22	0/1/1/1
7	NAG	A	709	1	-	2/6/23/26	0/1/1/1
7	NAG	B	706	1	-	4/6/23/26	0/1/1/1
6	MAN	D	701	1	-	0/2/19/22	0/1/1/1
7	NAG	B	704	1	-	5/6/23/26	0/1/1/1
7	NAG	C	708	1	-	2/6/23/26	0/1/1/1
7	NAG	C	710	1	-	2/6/23/26	0/1/1/1
7	NAG	D	704	1	-	2/6/23/26	0/1/1/1
7	NAG	C	707	1	-	2/6/23/26	0/1/1/1
6	MAN	B	701	1	-	0/2/19/22	0/1/1/1
7	NAG	A	708	1	-	0/6/23/26	0/1/1/1
7	NAG	B	710	1	-	2/6/23/26	0/1/1/1
7	NAG	D	705	1	-	5/6/23/26	0/1/1/1
7	NAG	B	708	1	-	0/6/23/26	0/1/1/1
7	NAG	D	706	1	-	2/6/23/26	0/1/1/1
6	MAN	A	704	1	-	2/2/19/22	0/1/1/1
6	MAN	D	703	1	-	2/2/19/22	0/1/1/1
7	NAG	C	709	1	-	3/6/23/26	0/1/1/1
7	NAG	A	707	1	-	1/6/23/26	0/1/1/1
6	MAN	C	701	1	-	2/2/19/22	0/1/1/1
6	MAN	A	701	1	-	0/2/19/22	0/1/1/1
7	NAG	A	710	1	-	1/6/23/26	0/1/1/1
7	NAG	D	709	1	-	2/6/23/26	0/1/1/1

The worst 5 of 14 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	C	703	MAN	O3-C3	4.44	1.53	1.43
6	B	702	MAN	O4-C4	4.41	1.53	1.43
6	C	703	MAN	O2-C2	3.90	1.51	1.43
6	B	702	MAN	O5-C1	3.81	1.49	1.43
6	C	703	MAN	O5-C1	3.60	1.49	1.43

The worst 5 of 70 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	D	705	NAG	C2-N2-C7	6.66	132.39	122.90

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	706	NAG	O5-C1-C2	-5.33	102.87	111.29
6	C	702	MAN	O5-C1-C2	-5.19	102.77	110.77
7	B	704	NAG	C2-N2-C7	-4.89	115.93	122.90
6	A	704	MAN	O5-C1-C2	-4.72	103.49	110.77

There are no chirality outliers.

5 of 74 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	B	704	NAG	C3-C2-N2-C7
7	B	704	NAG	C8-C7-N2-C2
7	B	704	NAG	O7-C7-N2-C2
7	B	705	NAG	C3-C2-N2-C7
7	B	705	NAG	C8-C7-N2-C2

There are no ring outliers.

12 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	C	705	NAG	1	0
6	D	702	MAN	1	0
6	C	704	MAN	1	0
7	A	706	NAG	1	0
7	A	709	NAG	1	0
7	B	706	NAG	2	0
6	D	701	MAN	1	0
7	D	704	NAG	1	0
7	B	710	NAG	1	0
7	D	705	NAG	2	0
6	C	701	MAN	1	0
7	D	709	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	525/675 (77%)	0.06	13 (2%) 57 59	22, 32, 56, 87	0
1	B	526/675 (77%)	0.28	33 (6%) 20 21	22, 35, 72, 105	0
1	C	528/675 (78%)	0.16	16 (3%) 50 53	22, 36, 62, 109	0
1	D	524/675 (77%)	0.46	44 (8%) 11 12	22, 38, 78, 103	0
All	All	2103/2700 (77%)	0.24	106 (5%) 28 31	22, 35, 68, 109	0

The worst 5 of 106 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	555	CYS	6.2
1	C	143	CYS	4.9
1	D	554	LYS	4.8
1	B	513	THR	4.7
1	D	436	ASN	4.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	MAN	G	5	11/12	0.68	0.40	92,102,108,109	0
3	BMA	I	3	11/12	0.69	0.36	89,103,108,111	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	MAN	G	6	11/12	0.71	0.25	27,28,29,31	11
3	BMA	F	3	11/12	0.78	0.29	70,74,80,81	0
2	NAG	H	2	14/15	0.78	0.28	71,78,91,91	0
3	NAG	I	2	14/15	0.80	0.25	69,89,95,97	0
3	NAG	J	2	14/15	0.80	0.21	59,72,80,85	0
2	NAG	E	2	14/15	0.82	0.39	60,67,72,73	0
3	BMA	J	3	11/12	0.82	0.28	85,95,101,101	0
2	NAG	K	2	14/15	0.83	0.32	79,92,102,103	0
3	NAG	F	2	14/15	0.85	0.31	54,67,79,80	0
4	BMA	G	3	11/12	0.86	0.16	66,78,82,84	0
3	NAG	I	1	14/15	0.91	0.20	56,68,72,77	0
4	NAG	G	2	14/15	0.92	0.24	51,61,72,75	0
4	MAN	G	4	11/12	0.92	0.23	84,86,90,96	0
2	NAG	E	1	14/15	0.94	0.14	42,47,50,57	0
5	GLC	N	1	11/12	0.94	0.13	44,48,52,53	0
5	GLC	O	1	11/12	0.94	0.12	34,37,41,43	0
4	NAG	G	1	14/15	0.95	0.18	46,51,55,56	0
3	NAG	J	1	14/15	0.95	0.15	34,38,41,51	0
3	NAG	F	1	14/15	0.95	0.13	35,40,45,47	0
2	NAG	K	1	14/15	0.95	0.11	44,48,60,66	0
2	NAG	H	1	14/15	0.96	0.14	37,42,46,57	0
5	FRU	M	2	12/12	0.96	0.17	27,30,34,44	0
5	GLC	M	1	11/12	0.97	0.12	35,37,41,43	0
5	FRU	N	2	12/12	0.97	0.15	29,35,40,44	0
5	FRU	L	2	12/12	0.97	0.11	28,32,34,37	0
5	FRU	O	2	12/12	0.97	0.12	26,29,33,38	0
5	GLC	L	1	11/12	0.98	0.11	37,39,41,44	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain H:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain K:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain F:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain I:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain J:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain G:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain L:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain M:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain N:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	MAN	D	701	11/12	0.69	0.43	88,106,121,121	0
6	MAN	C	703	11/12	0.71	0.43	27,29,31,32	0
6	MAN	B	702	11/12	0.75	0.43	27,29,31,31	0
6	MAN	A	702	11/12	0.76	0.31	82,90,95,96	0
7	NAG	B	705	14/15	0.76	0.45	72,80,83,84	0
6	MAN	D	703	11/12	0.77	0.34	74,79,84,86	0
7	NAG	B	704	14/15	0.80	0.31	59,67,82,84	0
7	NAG	D	708	14/15	0.80	0.35	96,112,121,129	0
6	MAN	C	702	11/12	0.81	0.26	76,85,88,89	0
7	NAG	A	706	14/15	0.82	0.33	66,74,77,78	0
6	MAN	A	704	11/12	0.84	0.25	64,71,78,78	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	NAG	D	705	14/15	0.84	0.19	57,63,69,69	0
6	MAN	C	704	11/12	0.84	0.21	56,64,69,71	0
6	MAN	B	703	11/12	0.85	0.21	77,80,89,90	0
7	NAG	B	706	14/15	0.85	0.34	63,74,84,85	0
6	MAN	C	701	11/12	0.86	0.21	71,79,84,85	0
6	MAN	A	703	11/12	0.86	0.23	63,68,75,76	0
7	NAG	D	706	14/15	0.86	0.49	59,70,77,78	0
6	MAN	B	701	11/12	0.86	0.25	79,86,89,89	0
7	NAG	D	709	14/15	0.86	0.22	69,72,78,85	0
7	NAG	A	705	14/15	0.87	0.25	58,65,79,80	0
7	NAG	B	710	14/15	0.88	0.26	66,75,80,80	0
7	NAG	C	709	14/15	0.88	0.18	54,63,71,76	0
6	MAN	A	701	11/12	0.88	0.34	72,79,82,83	0
7	NAG	C	705	14/15	0.89	0.26	62,71,81,84	0
6	MAN	D	702	11/12	0.90	0.45	81,92,99,110	0
7	NAG	B	709	14/15	0.91	0.21	64,71,80,82	0
7	NAG	A	710	14/15	0.91	0.15	51,65,70,75	0
7	NAG	A	707	14/15	0.92	0.15	46,51,55,56	0
7	NAG	D	704	14/15	0.92	0.14	55,59,66,67	0
7	NAG	C	707	14/15	0.93	0.22	51,58,61,65	0
7	NAG	A	711	14/15	0.93	0.16	46,56,64,67	0
7	NAG	C	710	14/15	0.93	0.15	50,52,62,68	0
7	NAG	C	706	14/15	0.93	0.15	55,63,72,72	0
7	NAG	A	709	14/15	0.94	0.16	46,53,56,60	0
7	NAG	A	708	14/15	0.94	0.13	37,40,46,47	0
7	NAG	D	707	14/15	0.95	0.16	43,48,56,58	0
7	NAG	C	708	14/15	0.95	0.17	48,52,57,58	0
7	NAG	B	708	14/15	0.95	0.14	40,43,49,50	0
7	NAG	B	707	14/15	0.97	0.10	30,33,41,42	0

6.5 Other polymers [i](#)

There are no such residues in this entry.