



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 27, 2024 – 02:42 PM EST

PDB ID : 1BFT  
Title : STRUCTURE OF NF-KB P50 HOMODIMER BOUND TO A KB SITE  
Authors : Huang, D.B.; Huxford, T.; Chen, Y.Q.; Ghosh, G.  
Deposited on : 1997-09-12  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtrriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	101	
1	B	101	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2241 atoms, of which 534 are hydrogens and 8 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NUCLEAR FACTOR NF-KAPPA-B P65.

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
			Total	C	D	H	N	O	S			
1	A	101	987	504	4	178	142	154	5	0	0	0
1	B	101	987	504	4	178	142	154	5	0	0	0

- Molecule 2 is water.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	H	O		
2	A	55	165	110	55	0	0
2	B	34	102	68	34	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: NUCLEAR FACTOR NF-KAPPA-B P65

Chain A: 



- Molecule 1: NUCLEAR FACTOR NF-KAPPA-B P65

Chain B: 



## 4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	55.50Å 74.49Å 105.61Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 2.00	Depositor
% Data completeness (in resolution range)	(Not available) (8.00-2.00)	Depositor
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, $R_{free}$	0.204 , 0.294	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	2241	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	15.0	wwPDB-VP

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.67	0/823	0.90	1/1113 (0.1%)
1	B	0.68	0/823	0.91	0/1113
All	All	0.68	0/1646	0.91	1/2226 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	280	LEU	CA-CB-CG	5.66	128.31	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	809	178	782	37	0
1	B	809	178	782	41	1
2	A	55	110	0	1	1
2	B	34	68	0	1	1
All	All	1707	534	1564	75	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 24.

All (75) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:195:LYS:NZ	1:B:197:CYS:SG	2.39	0.94
1:A:206:CYS:HB3	1:A:262:LEU:HD22	1.54	0.89
1:B:271:GLN:HG3	1:B:280:LEU:HD22	1.60	0.82
1:B:233:TRP:CD1	1:B:258:ALA:HB2	2.15	0.82
1:B:191:THR:N	1:B:281:SER:HA	1.94	0.81
1:A:194:LEU:HB3	1:A:281:SER:HB3	1.66	0.76
1:A:195:LYS:NZ	1:A:197:CYS:SG	2.57	0.76
1:B:193:GLU:HG3	1:B:274:ARG:NH2	2.03	0.73
1:A:221:LYS:HG3	1:A:241:GLN:HB3	1.70	0.72
1:B:273:ARG:HG2	1:B:280:LEU:CD2	2.20	0.71
1:A:236:ARG:HH11	1:A:236:ARG:HG3	1.59	0.68
1:B:199:VAL:HA	1:B:213:PHE:O	1.95	0.66
1:A:228:PHE:O	1:A:234:GLU:HA	1.96	0.66
1:A:229:THR:HG22	1:A:269:SER:HB3	1.80	0.64
1:B:195:LYS:CE	1:B:197:CYS:SG	2.86	0.63
1:B:226:VAL:HG11	1:B:252:PHE:CZ	2.35	0.62
1:B:273:ARG:HG2	1:B:280:LEU:HD21	1.81	0.61
1:A:193:GLU:HG3	1:A:274:ARG:HH22	1.67	0.60
1:B:206:CYS:HB3	1:B:262:LEU:HD22	1.82	0.60
1:A:195:LYS:CE	1:A:197:CYS:SG	2.92	0.58
1:B:273:ARG:HG2	1:B:280:LEU:HD23	1.86	0.57
1:A:225:GLU:OE2	1:A:273:ARG:HD3	2.05	0.57
1:A:206:CYS:CB	1:A:262:LEU:HD22	2.33	0.56
1:B:245:HIS:HB3	1:B:249:ALA:HB3	1.87	0.56
1:A:195:LYS:HE2	1:A:197:CYS:SG	2.45	0.56
1:A:198:ARG:HB3	1:A:215:LEU:HD12	1.87	0.56
1:B:204:GLY:O	1:B:288:TYR:HA	2.05	0.56
1:A:222:GLU:H	1:A:222:GLU:CD	2.08	0.55
1:B:259:ASP:HB3	1:B:262:LEU:HD13	1.88	0.55
1:A:245:HIS:HB3	1:A:249:ALA:HB3	1.88	0.54
1:A:246:ARG:O	1:A:247:GLN:HB2	2.07	0.54
1:B:195:LYS:HE2	1:B:197:CYS:SG	2.48	0.54
1:B:201:ARG:NH1	1:B:203:SER:O	2.41	0.54
1:B:274:ARG:NH2	1:B:279:GLU:HG2	2.23	0.53
1:A:273:ARG:HG3	1:A:280:LEU:CD1	2.39	0.53
1:A:233:TRP:CD1	1:A:258:ALA:HB2	2.43	0.53
1:B:191:THR:N	1:B:280:LEU:O	2.42	0.52
1:B:227:TYR:HE2	1:B:234:GLU:CD	2.13	0.52
1:A:206:CYS:HA	1:A:288:TYR:CD1	2.45	0.52
1:B:246:ARG:HB3	2:B:625:HOH:O	2.10	0.51
1:A:248:VAL:HG21	1:B:246:ARG:HB2	1.92	0.51
1:B:193:GLU:CD	1:B:274:ARG:HH22	2.13	0.51

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:239:PHE:HB3	1:B:252:PHE:HB3	1.92	0.51
1:A:273:ARG:HG3	1:A:280:LEU:HD13	1.91	0.51
1:B:207:LEU:HD12	1:B:291:ASP:OD2	2.11	0.50
1:B:239:PHE:CD1	1:B:252:PHE:HB3	2.47	0.50
1:A:219:VAL:HG22	1:A:247:GLN:O	2.12	0.50
1:A:225:GLU:HG3	1:A:273:ARG:HB3	1.94	0.49
1:A:229:THR:HB	2:A:699:HOH:O	2.12	0.49
1:A:262:LEU:HD21	1:A:266:VAL:HG12	1.94	0.49
1:A:236:ARG:HG3	1:A:236:ARG:NH1	2.28	0.48
1:B:193:GLU:HG3	1:B:274:ARG:HH21	1.75	0.48
1:B:245:HIS:CD2	1:B:246:ARG:HG2	2.49	0.47
1:B:194:LEU:HB3	1:B:281:SER:HB3	1.96	0.47
1:B:193:GLU:CG	1:B:274:ARG:NH2	2.76	0.46
1:B:246:ARG:O	1:B:247:GLN:HB2	2.15	0.46
1:A:206:CYS:HB3	1:A:262:LEU:CD2	2.37	0.46
1:A:218:LYS:HG3	1:A:247:GLN:HB3	1.98	0.45
1:B:201:ARG:HD3	1:B:203:SER:O	2.16	0.45
1:B:227:TYR:O	1:B:270:MET:HA	2.16	0.45
1:A:220:GLN:HB2	1:A:223:ASP:HB3	1.99	0.44
1:A:213:PHE:CZ	1:B:198:ARG:HD2	2.51	0.44
1:B:268:VAL:HG23	1:B:286:PHE:HB3	2.00	0.44
1:A:194:LEU:HB3	1:A:281:SER:CB	2.41	0.44
1:B:228:PHE:O	1:B:234:GLU:HA	2.18	0.43
1:B:288:TYR:C	1:B:289:LEU:HD13	2.39	0.43
1:B:239:PHE:HB2	1:B:243:ASP:CB	2.48	0.43
1:A:193:GLU:CG	1:A:274:ARG:HH22	2.31	0.42
1:A:200:ASN:HA	1:B:213:PHE:CE1	2.55	0.42
1:A:267:ARG:HG2	1:A:287:GLN:HG2	2.01	0.42
1:B:226:VAL:HG21	1:B:252:PHE:CE1	2.55	0.41
1:B:267:ARG:HG3	1:B:285:GLU:OE2	2.21	0.41
1:A:225:GLU:HG2	1:A:275:PRO:HD3	2.02	0.41
1:A:218:LYS:HB2	1:A:218:LYS:NZ	2.36	0.40
1:A:212:ILE:O	1:A:251:VAL:HA	2.21	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:253:ARG:HH21	2:B:662:HOH:H2[8_556]	1.33	0.27
2:A:687:HOH:O	2:A:725:HOH:H1[3_655]	1.54	0.06



## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	99/101 (98%)	94 (95%)	5 (5%)	0	100	100
1	B	99/101 (98%)	93 (94%)	5 (5%)	1 (1%)	15	9
All	All	198/202 (98%)	187 (94%)	10 (5%)	1 (0%)	29	23

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	192	ALA

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	89/89 (100%)	80 (90%)	9 (10%)	7	4
1	B	89/89 (100%)	80 (90%)	9 (10%)	7	4
All	All	178/178 (100%)	160 (90%)	18 (10%)	7	4

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	194	LEU
1	A	198	ARG
1	A	222	GLU
1	A	229	THR
1	A	246	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	262	LEU
1	A	279	GLU
1	A	280	LEU
1	A	289	LEU
1	B	193	GLU
1	B	194	LEU
1	B	201	ARG
1	B	203	SER
1	B	240	SER
1	B	262	LEU
1	B	269	SER
1	B	273	ARG
1	B	289	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	220	GLN
1	B	245	HIS
1	B	287	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

### 6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.