



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 13, 2023 – 12:28 am GMT

PDB ID : 8BJY
Title : Engineered Fructosyl Peptide Oxidase - X02B mutant
Authors : Estiri, H.; Bhattacharya, S.; Rodriguez-Buitrago, J.A.; Parisini, E.
Deposited on : 2022-11-08
Resolution : 1.48 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.48 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4690 (1.50-1.46)
Clashscore	141614	4955 (1.50-1.46)
Ramachandran outliers	138981	4846 (1.50-1.46)
Sidechain outliers	138945	4844 (1.50-1.46)
RSRZ outliers	127900	4614 (1.50-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	432	 2% 90% 7% ..

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	503	-	-	-	X

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 7127 atoms, of which 3365 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fructosyl Peptide Oxidase mutant (X02B).

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	427	6682	2150	3318	586	609	19	184	5	0

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	9	LYS	SER	conflict	UNP Q0UIL6
A	48	CYS	SER	conflict	UNP Q0UIL6
A	63	ALA	LEU	conflict	UNP Q0UIL6
A	64	ASP	ARG	conflict	UNP Q0UIL6
A	65	PRO	ASN	conflict	UNP Q0UIL6
A	66	ALA	PRO	conflict	UNP Q0UIL6
A	67	ALA	VAL	conflict	UNP Q0UIL6
A	?	-	LEU	deletion	UNP Q0UIL6
A	?	-	GLN	deletion	UNP Q0UIL6
A	?	-	LEU	deletion	UNP Q0UIL6
A	?	-	ALA	deletion	UNP Q0UIL6
A	?	-	LEU	deletion	UNP Q0UIL6
A	69	ALA	GLU	conflict	UNP Q0UIL6
A	104	LYS	SER	conflict	UNP Q0UIL6
A	105	LEU	GLY	conflict	UNP Q0UIL6
A	108	ASN	ALA	conflict	UNP Q0UIL6
A	112	TRP	ALA	conflict	UNP Q0UIL6
A	118	VAL	ASN	conflict	UNP Q0UIL6
A	133	GLN	LEU	conflict	UNP Q0UIL6
A	135	THR	SER	conflict	UNP Q0UIL6
A	160	LYS	ASN	conflict	UNP Q0UIL6
A	162	ILE	VAL	conflict	UNP Q0UIL6
A	183	GLN	ALA	conflict	UNP Q0UIL6
A	222	GLN	HIS	conflict	UNP Q0UIL6
A	267	ILE	VAL	conflict	UNP Q0UIL6
A	301	CYS	ALA	conflict	UNP Q0UIL6
A	313	ILE	ASP	conflict	UNP Q0UIL6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	315	ARG	SER	conflict	UNP Q0UIL6
A	331	PRO	LYS	conflict	UNP Q0UIL6
A	332	LEU	MET	conflict	UNP Q0UIL6
A	346	GLY	ALA	conflict	UNP Q0UIL6
A	347	HIS	ALA	conflict	UNP Q0UIL6
A	360	TYR	VAL	conflict	UNP Q0UIL6
A	375	ILE	ASN	conflict	UNP Q0UIL6
A	379	TYR	HIS	conflict	UNP Q0UIL6
A	391	GLU	ASP	conflict	UNP Q0UIL6
A	395	LYS	ALA	conflict	UNP Q0UIL6
A	411	GLU	SER	conflict	UNP Q0UIL6
A	427	HIS	-	expression tag	UNP Q0UIL6
A	428	HIS	-	expression tag	UNP Q0UIL6
A	429	HIS	-	expression tag	UNP Q0UIL6
A	430	HIS	-	expression tag	UNP Q0UIL6
A	431	HIS	-	expression tag	UNP Q0UIL6
A	432	HIS	-	expression tag	UNP Q0UIL6

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	O	2	0
			14	3	8	3		
2	A	1	Total	C	H	O	2	0
			14	3	8	3		

- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula:

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Fructosyl Peptide Oxidase mutant (X02B)



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	63.11Å 88.49Å 90.94Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	63.42 – 1.48 63.42 – 1.48	Depositor EDS
% Data completeness (in resolution range)	93.7 (63.42-1.48) 93.7 (63.42-1.48)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.33 (at 1.48Å)	Xtrriage
Refinement program	REFMAC 5.8.0352	Depositor
R, R_{free}	0.189 , 0.215 0.188 , 0.215	Depositor DCC
R_{free} test set	4009 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	27.7	Xtrriage
Anisotropy	0.279	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.43 , 44.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.004 for -h,l,k	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7127	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FAD, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.52	0/3472	0.88	5/4710 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	6

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	293	ARG	NE-CZ-NH2	-13.03	113.78	120.30
1	A	293	ARG	NE-CZ-NH1	9.59	125.09	120.30
1	A	399	ARG	NE-CZ-NH2	-7.27	116.66	120.30
1	A	279	ARG	NE-CZ-NH1	6.39	123.50	120.30
1	A	399	ARG	NE-CZ-NH1	5.89	123.24	120.30

There are no chirality outliers.

All (6) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	172	ARG	Sidechain
1	A	2	ALA	Peptide
1	A	201	ARG	Sidechain
1	A	293	ARG	Sidechain
1	A	317	ARG	Sidechain

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	A	409	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3364	3318	3308	35	0
2	A	12	16	16	5	0
3	A	53	31	31	7	0
4	A	333	0	0	11	0
All	All	3762	3365	3355	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:338:CYS:SG	3:A:502:FAD:HM82	1.55	1.47
1:A:338:CYS:SG	3:A:502:FAD:C8M	2.30	1.18
2:A:501:GOL:H32	4:A:801:HOH:O	1.41	1.18
1:A:71:ARG:NH1	4:A:601:HOH:O	1.97	0.97
1:A:293:ARG:HD2	4:A:735:HOH:O	1.80	0.81
1:A:61:VAL:HA	2:A:503:GOL:H31	1.66	0.77
1:A:104:LYS:HE3	4:A:803:HOH:O	1.88	0.74
1:A:338:CYS:CB	3:A:502:FAD:HM82	2.19	0.72
1:A:419:ASP:OD2	2:A:501:GOL:H31	1.96	0.66
1:A:61:VAL:HA	2:A:503:GOL:C3	2.26	0.64
1:A:60:GLY:CA	4:A:605:HOH:O	2.45	0.64
1:A:280:PHE:CG	1:A:293:ARG:HD3	2.34	0.62
3:A:502:FAD:N6A	4:A:603:HOH:O	2.32	0.57
1:A:68:ASP:OD1	2:A:503:GOL:O1	2.22	0.56
1:A:338:CYS:SG	3:A:502:FAD:HM81	2.40	0.56
1:A:280:PHE:CD1	1:A:293:ARG:HD3	2.41	0.56
1:A:60:GLY:HA2	4:A:605:HOH:O	2.06	0.55
1:A:315:ARG:HG3	4:A:624:HOH:O	2.06	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:167:ARG:HH11	1:A:167:ARG:HG2	1.75	0.52
1:A:2:ALA:HB3	1:A:3:PRO:CD	2.39	0.51
1:A:61:VAL:HG22	1:A:71:ARG:HH21	1.76	0.49
1:A:61:VAL:N	4:A:605:HOH:O	2.45	0.49
1:A:15:GLY:HA3	1:A:39:VAL:HG13	1.95	0.48
1:A:71:ARG:HH22	1:A:150:GLY:N	2.10	0.48
1:A:338:CYS:SG	3:A:502:FAD:C8	3.03	0.47
1:A:74:TRP:O	1:A:81:LYS:HA	2.16	0.45
1:A:231:VAL:HG21	1:A:312:SER:HB3	1.98	0.45
1:A:104:LYS:CE	4:A:803:HOH:O	2.55	0.43
1:A:58:ILE:HD11	1:A:258:PHE:CD2	2.54	0.43
1:A:230:TRP:HB2	3:A:502:FAD:HM83	2.01	0.42
1:A:60:GLY:C	4:A:605:HOH:O	2.58	0.42
1:A:129:LYS:HA	1:A:129:LYS:HE3	2.02	0.42
1:A:213:ALA:HA	1:A:341:THR:HB	2.02	0.42
1:A:56:ASN:HA	1:A:153:LEU:O	2.19	0.42
1:A:130:ARG:HD3	1:A:130:ARG:HA	1.93	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	430/432 (100%)	416 (97%)	13 (3%)	1 (0%)	47 23

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	97	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	353/353 (100%)	350 (99%)	3 (1%)	81 64

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	97	LYS
1	A	129	LYS
1	A	338	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	86	ASN
1	A	347	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	A	503	-	5,5,5	0.18	0	5,5,5	0.50	0
3	FAD	A	502	-	53,58,58	0.84	1 (1%)	68,89,89	0.91	2 (2%)
2	GOL	A	501	-	5,5,5	0.14	0	5,5,5	0.42	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	503	-	-	4/4/4/4	-
3	FAD	A	502	-	-	2/30/50/50	0/6/6/6
2	GOL	A	501	-	-	0/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	502	FAD	C1'-C2'	-3.21	1.48	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	502	FAD	O4B-C1B-C2B	-2.40	103.42	106.93
3	A	502	FAD	O4-C4-C4X	-2.07	121.11	126.60

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	503	GOL	O1-C1-C2-C3
2	A	503	GOL	C1-C2-C3-O3
2	A	503	GOL	O2-C2-C3-O3
3	A	502	FAD	O4B-C4B-C5B-O5B

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	A	503	GOL	O1-C1-C2-O2
3	A	502	FAD	C2'-C3'-C4'-O4'

There are no ring outliers.

3 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	503	GOL	3	0
3	A	502	FAD	7	0
2	A	501	GOL	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	427/432 (98%)	0.31	9 (2%) 63 67	17, 27, 47, 103	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1	MET	12.7
1	A	61	VAL	11.6
1	A	2	ALA	8.8
1	A	96	GLU	4.4
1	A	149	ASP	3.2
1	A	409	ARG	3.1
1	A	55	LEU	2.6
1	A	51	ALA	2.1
1	A	100	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	GOL	A	503	6/6	0.57	0.49	55,72,78,78	2
2	GOL	A	501	6/6	0.86	0.24	33,52,55,58	2
3	FAD	A	502	53/53	0.96	0.10	18,22,35,38	5

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.