



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 8, 2020 – 06:02 AM BST

PDB ID : 1BTE
Title : CRYSTAL STRUCTURE OF THE EXTRACELLULAR DOMAIN OF THE
TYPE II ACTIVIN RECEPTOR
Authors : Greenwald, J.; Fischer, W.; Vale, W.; Choe, S.
Deposited on : 1998-09-01
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	3144 (1.50-1.50)
Ramachandran outliers	138981	3066 (1.50-1.50)
Sidechain outliers	138945	3064 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	97	93% 5%
1	B	97	89% 7%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 1745 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEIN (ACTIVIN RECEPTOR TYPE II).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	92	Total 740	C 460	N 121	O 147	S 12	0	1	0
1	B	94	Total 745	C 465	N 124	O 145	S 11	0	0	0

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	Total 15	C 8	N 1	O 6	0	1
2	A	1	Total 14	C 8	N 1	O 5	0	0
2	B	1	Total 15	C 8	N 1	O 6	0	1
2	B	1	Total 14	C 8	N 1	O 5	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	116	Total 116	O 116	0	0
3	B	86	Total 86	O 86	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.

- Molecule 1: PROTEIN (ACTIVIN RECEPTOR TYPE II)

Chain A:  93% 5%



- Molecule 1: PROTEIN (ACTIVIN RECEPTOR TYPE II)

Chain B:  89% 7% ..



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 3	Depositor
Cell constants a, b, c, α , β , γ	71.63Å 71.63Å 37.38Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 1.50	Depositor
% Data completeness (in resolution range)	83.9 (20.00-1.50)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.03	Depositor
Refinement program	REFMAC	Depositor
R, R_{free}	0.181 , 0.222	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	1745	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.69	0/765	1.19	2/1028 (0.2%)
1	B	0.64	0/764	1.18	0/1034
All	All	0.66	0/1529	1.18	2/2062 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	21	ASP	CB-CG-OD1	5.46	123.22	118.30
1	A	77	ASP	CB-CG-OD1	5.46	123.21	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	68	ASP	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	740	0	628	0	0
1	B	745	0	645	4	1
2	A	29	0	19	0	0
2	B	29	0	19	0	0
3	A	116	0	0	0	1
3	B	86	0	0	3	3
All	All	1745	0	1311	4	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (4) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:75:LYS:HD2	3:B:226:HOH:O	2.05	0.56
1:B:40:HIS:HD2	3:B:151:HOH:O	1.94	0.49
1:B:48:ILE:O	1:B:48:ILE:HG13	2.13	0.48
1:B:76:LYS:HE2	3:B:229:HOH:O	2.22	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:149:HOH:O	3:B:149:HOH:O[2_555]	0.59	1.61
3:A:214:HOH:O	3:B:222:HOH:O[2_555]	2.09	0.11
1:B:93:GLU:OE2	3:B:149:HOH:O[2_555]	2.10	0.10

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	88/97 (91%)	87 (99%)	1 (1%)	0	100	100
1	B	92/97 (95%)	90 (98%)	2 (2%)	0	100	100
All	All	180/194 (93%)	177 (98%)	3 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	81/90 (90%)	81 (100%)	0	100	100
1	B	80/90 (89%)	77 (96%)	3 (4%)	33	7
All	All	161/180 (89%)	158 (98%)	3 (2%)	57	27

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	48	ILE
1	B	52	ILE
1	B	78	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	40	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	A	124[B]	-	14,14,15	1.16	1 (7%)	17,19,21	1.41	3 (17%)
2	NAG	B	147	1	14,14,15	1.43	3 (21%)	17,19,21	1.65	4 (23%)
2	NAG	A	124[A]	-	14,14,15	1.14	1 (7%)	17,19,21	1.41	3 (17%)
2	NAG	B	124[A]	-	14,14,15	1.19	1 (7%)	17,19,21	1.64	3 (17%)
2	NAG	A	147	1	14,14,15	1.01	1 (7%)	17,19,21	1.10	0
2	NAG	B	124[B]	-	14,14,15	1.17	1 (7%)	17,19,21	1.63	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	124[B]	-	-	0/6/23/26	0/1/1/1
2	NAG	B	147	1	-	1/6/23/26	0/1/1/1
2	NAG	A	124[A]	-	-	0/6/23/26	0/1/1/1
2	NAG	B	124[A]	-	-	0/6/23/26	0/1/1/1
2	NAG	A	147	1	-	0/6/23/26	0/1/1/1
2	NAG	B	124[B]	-	-	0/6/23/26	0/1/1/1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	147	NAG	O7-C7	-3.42	1.15	1.23
2	B	124[A]	NAG	O7-C7	-3.18	1.16	1.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	124[B]	NAG	O7-C7	-3.18	1.16	1.23
2	A	124[B]	NAG	O7-C7	-2.79	1.16	1.23
2	A	124[A]	NAG	O7-C7	-2.79	1.16	1.23
2	B	147	NAG	O5-C1	2.74	1.48	1.43
2	A	147	NAG	O7-C7	-2.67	1.17	1.23
2	B	147	NAG	C2-N2	2.44	1.50	1.46

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	124[A]	NAG	O5-C1-C2	-3.77	105.33	111.29
2	B	124[B]	NAG	O5-C1-C2	-3.77	105.33	111.29
2	B	147	NAG	O5-C5-C6	-3.33	101.98	107.20
2	A	124[B]	NAG	O5-C1-C2	-3.30	106.07	111.29
2	A	124[A]	NAG	O5-C1-C2	-3.30	106.07	111.29
2	B	147	NAG	C1-O5-C5	-3.18	107.89	112.19
2	B	124[A]	NAG	C4-C3-C2	-2.61	107.20	111.02
2	B	124[B]	NAG	C4-C3-C2	-2.61	107.20	111.02
2	B	147	NAG	O5-C1-C2	-2.46	107.40	111.29
2	A	124[B]	NAG	C1-O5-C5	-2.38	108.97	112.19
2	A	124[A]	NAG	C1-O5-C5	-2.38	108.97	112.19
2	B	124[A]	NAG	O3-C3-C4	-2.16	105.36	110.35
2	B	124[B]	NAG	O3-C3-C4	-2.16	105.36	110.35
2	A	124[B]	NAG	O3-C3-C2	2.11	113.84	109.47
2	A	124[A]	NAG	O3-C3-C2	2.11	113.84	109.47
2	B	147	NAG	C8-C7-N2	-2.01	112.70	116.10

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	147	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.