

Full wwPDB EM Validation Report (i)

Mar 3, 2024 – 10:06 AM EST

PDB ID : 6BUZ

EMDB ID : EMD-7293

Title: Cryo-EM structure of CENP-A nucleosome in complex with kinetochore pro-

tein CENP-N

Authors : Chittori, S.; Hong, J.; Kelly, A.E.; Bai, Y.; Subramaniam, S.

Deposited on : 2017-12-11

Resolution : 3.92 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev70

MolProbity : 4.02b-467

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $MapQ \quad : \quad 1.9.13$

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 3.92 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	${ m EM\ structures} \ (\#{ m Entries})$
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion <40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality	of chair	1	
1	A	160	52%	•	43%	
1	Е	160	<u>•</u> 53%	·	43%	
2	В	103	67%		11%	22%
2	F	103	69%		9%	22%
3	С	130	75%		8%	18%
3	G	130	75%		7%	18%
4	D	126	• 67%		8%	25%
4	Н	126	67%		8%	25%



Continued from previous page...

Mol	Chain	Length	Quality of chain		
5	I	147	75%	24%	-
6	J	147	71%	27%	•
7	N	668	20% 80%		-



2 Entry composition (i)

There are 7 unique types of molecules in this entry. The entry contains 12509 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Histone H3-like centromeric protein A.

Mol	Chain	Residues		At	oms	AltConf	Trace		
1	А	91	Total				S	0	0
1	1 11	01	749	488	139	121	1		
1	E	01	Total	С	N	О	S	0	0
1	$1 \mid \mathbf{E}$	91	749	488	139	121	1		

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	expression tag	UNP P49450
A	-18	GLY	-	expression tag	UNP P49450
A	-17	SER	-	expression tag	UNP P49450
A	-16	SER	-	expression tag	UNP P49450
A	-15	HIS	-	expression tag	UNP P49450
A	-14	HIS	-	expression tag	UNP P49450
A	-13	HIS	-	expression tag	UNP P49450
A	-12	HIS	-	expression tag	UNP P49450
A	-11	HIS	-	expression tag	UNP P49450
A	-10	HIS	-	expression tag	UNP P49450
A	-9	SER	-	expression tag	UNP P49450
A	-8	SER	-	expression tag	UNP P49450
A	-7	GLY	-	expression tag	UNP P49450
A	-6	LEU	-	expression tag	UNP P49450
A	-5	VAL	-	expression tag	UNP P49450
A	-4	PRO	-	expression tag	UNP P49450
A	-3	ARG	-	expression tag	UNP P49450
A	-2	GLY	-	expression tag	UNP P49450
A	-1	SER	-	expression tag	UNP P49450
A	0	HIS	-	expression tag	UNP P49450
Е	-19	MET	-	expression tag	UNP P49450
Е	-18	GLY	-	expression tag	UNP P49450
Е	-17	SER		expression tag	UNP P49450
Е	-16	SER	-	expression tag	UNP P49450
Е	-15	HIS	-	expression tag	UNP P49450
Е	-14	HIS	-	expression tag	UNP P49450



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled	Actual	Comment	Reference
Е	-13	HIS	=	expression tag	UNP P49450
Е	-12	HIS	-	expression tag	UNP P49450
E	-11	HIS	-	expression tag	UNP P49450
Е	-10	HIS	-	expression tag	UNP P49450
Е	-9	SER	-	expression tag	UNP P49450
Е	-8	SER	-	expression tag	UNP P49450
Е	-7	GLY	-	expression tag	UNP P49450
E	-6	LEU	-	expression tag	UNP P49450
Е	-5	VAL	-	expression tag	UNP P49450
E	-4	PRO	-	expression tag	UNP P49450
Е	-3	ARG	-	expression tag	UNP P49450
Е	-2	GLY	-	expression tag	UNP P49450
Е	-1	SER	-	expression tag	UNP P49450
Е	0	HIS	-	expression tag	UNP P49450

• Molecule 2 is a protein called Histone H4.

Mol	Chain	Residues		At	oms	AltConf	Trace	
2	В	80	Total 638		N 125	S 1	0	0
2	F	80	Total 638		N 125	S 1	0	0

• Molecule 3 is a protein called Histone H2A.

Mol	Chain	Residues	Atoms		AltConf	Trace
3	С	107	Total C N 826 520 163	O 143	0	0
3	G	107	Total C N 826 520 163	O 143	0	0

• Molecule 4 is a protein called Histone H2B.

Mol	Chain	Residues		At	oms	AltConf	Trace		
4	D	94	Total	С	N	О	S	0	0
4	4 D	94	736	462	134	138	2	0	
1	П	94	Total	С	N	О	S	0	0
4	4 H	94	736	462	134	138	2	0	U

• Molecule 5 is a DNA chain called DNA (147-MER).



Mol	Chain	Residues		A	AltConf	Trace			
5	I	145	Total 2952	C 1403	N 538	O 867	P 144	0	0

• Molecule 6 is a DNA chain called DNA (147-MER).

Mol	Chain	Residues		\mathbf{A}^{1}	AltConf	Trace			
6	J	145	Total 2987	C 1415	N 559	O 869	P 144	0	0

• Molecule 7 is a protein called Maltose-binding periplasmic protein, Centromere protein N chimera.

Mol	Chain	Residues		Ato	ms	AltConf	Trace	
7	N	134	Total 672		N 135	O 135	0	0

There are 18 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
N	-381	MET	-	expression tag	UNP P0AEY0
N	-380	LYS	-	expression tag	UNP P0AEY0
N	-379	GLU	-	expression tag	UNP P0AEY0
N	-378	HIS	-	expression tag	UNP P0AEY0
N	-377	HIS	-	expression tag	UNP P0AEY0
N	-376	HIS	-	expression tag	UNP P0AEY0
N	-375	HIS	-	expression tag	UNP P0AEY0
N	-374	HIS	-	expression tag	UNP P0AEY0
N	-373	HIS	-	expression tag	UNP P0AEY0
N	-372	HIS	-	expression tag	UNP P0AEY0
N	-371	HIS	-	expression tag	UNP P0AEY0
N	-11	ALA	GLU	engineered mutation	UNP P0AEY0
N	-8	ALA	LYS	engineered mutation	UNP P0AEY0
N	-7	ALA	ASP	engineered mutation	UNP P0AEY0
N	-3	ASN	ARG	engineered mutation	UNP P0AEY0
N	-2	ALA	ILE	engineered mutation	UNP P0AEY0
N	-1	ALA	-	linker	UNP P0AEY0
N	0	ALA	-	linker	UNP P0AEY0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Histone H3-like centromeric protein A





• Molecule 7: Maltose-binding periplasmic protein, Centromere protein N chimera







4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	61749	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE	Depositor
	CORRECTION	
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{Å}^2)$	40	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.006	Depositor
Minimum map value	-0.002	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.000	Depositor
Recommended contour level	0.0014	Depositor
Map size (Å)	235.2, 235.2, 235.2	wwPDB
Map dimensions	560, 560, 560	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.42, 0.42, 0.42	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond	angles
WIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.41	0/764	0.56	0/1029
1	Е	0.42	0/764	0.56	0/1029
2	В	0.42	0/645	0.54	0/862
2	F	0.42	0/645	0.54	0/862
3	С	0.38	0/836	0.55	0/1128
3	G	0.38	0/836	0.54	0/1128
4	D	0.39	0/747	0.49	0/1004
4	Н	0.41	0/747	0.50	0/1004
5	I	0.92	0/3308	0.99	0/5099
6	J	0.93	0/3354	1.00	0/5180
7	N	0.26	0/664	0.45	0/915
All	All	0.71	0/13310	0.81	0/19240

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	749	0	789	5	0
1	Е	749	0	789	4	0
2	В	638	0	676	10	0
2	F	638	0	676	7	0
3	С	826	0	884	9	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	G	826	0	884	8	0
4	D	736	0	758	8	0
4	Н	736	0	758	8	0
5	I	2952	0	1628	32	0
6	J	2987	0	1629	38	0
7	N	672	0	291	2	0
All	All	12509	0	9762	84	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (84) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
	7100H1 2	${ m distance} ({ m \AA})$	overlap (Å)
5:I:-27:DC:N3	6:J:26:DA:N6	2.48	0.61
3:G:78:ARG:NH1	6:J:-53:DA:OP1	2.34	0.60
5:I:-61:DG:H1	6:J:61:DC:H42	1.48	0.60
5:I:8:DC:H42	6:J:-8:DG:H1	1.49	0.60
5:I:7:DC:H42	6:J:-7:DG:H1	1.50	0.58
5:I:52:DG:O6	6:J:-53:DA:N6	2.35	0.58
1:A:69:ARG:NH1	6:J:17:DA:OP1	2.37	0.58
5:I:-49:DG:H1	6:J:49:DC:H42	1.51	0.57
1:E:62:ILE:O	1:E:95:GLN:NE2	2.38	0.57
5:I:5:DC:H42	6:J:-5:DG:H1	1.51	0.57
3:C:30:ARG:NH1	4:D:37:SER:O	2.38	0.56
1:A:62:ILE:O	1:A:95:GLN:NE2	2.37	0.56
4:H:37:SER:OG	4:H:38:TYR:N	2.39	0.56
3:C:45:GLY:HA2	6:J:38:DG:H5"	1.87	0.55
1:A:108:ASP:OD2	1:A:133:ARG:NH2	2.38	0.55
1:E:108:ASP:OD2	1:E:133:ARG:NH2	2.38	0.55
5:I:27:DG:O6	6:J:-28:DC:N4	2.40	0.55
3:C:78:ARG:HH21	6:J:57:DG:H4'	1.73	0.54
3:G:30:ARG:NH1	4:H:37:SER:O	2.40	0.54
5:I:17:DA:H61	6:J:-17:DT:H3	1.55	0.53
5:I:45:DC:N4	6:J:-46:DT:O4	2.41	0.53
3:C:30:ARG:NH2	6:J:49:DC:OP1	2.42	0.53
2:B:92:LYS:NZ	4:H:69:ASP:OD1	2.42	0.52
2:B:52:TYR:O	2:B:56:ARG:NH1	2.43	0.52
2:F:52:TYR:O	2:F:56:ARG:NH1	2.43	0.51
3:G:86:LEU:O	3:G:90:ASN:ND2	2.45	0.50
2:B:76:HIS:HE2	4:D:94:GLU:HG3	1.76	0.50



Continued from previous page...

Atom-1	1 A 1 🙃 '	Interatomic	Clash
	Atom-2	${\rm distance}\ (\mathring{\rm A})$	overlap (Å)
3:C:86:LEU:O	3:C:90:ASN:ND2	2.45	0.50
1:A:61:LEU:HD12	2:B:38:LEU:HD23	1.93	0.50
5:I:69:DC:O2	5:I:70:DC:N4	2.45	0.50
6:J:-15:DA:H1'	6:J:-14:DA:H5'	1.93	0.50
1:A:120:THR:OG1	2:B:46:ARG:NH1	2.44	0.49
2:F:93:ARG:HH21	4:H:102:LEU:HG	1.78	0.49
4:D:80:ARG:O	4:D:84:TYR:N	2.45	0.49
7:N:1:MET:O	7:N:5:VAL:N	2.44	0.48
4:D:105:GLY:O	4:D:109:LYS:NZ	2.39	0.48
3:G:43:ARG:HH21	5:I:38:DT:H4'	1.79	0.47
5:I:-17:DT:O4	6:J:16:DA:N6	2.47	0.47
5:I:-60:DG:O6	6:J:59:DA:N6	2.48	0.47
1:E:120:THR:OG1	2:F:46:ARG:NH1	2.39	0.46
5:I:-33:DA:N6	6:J:32:DG:O6	2.47	0.46
6:J:5:DT:H2"	6:J:6:DA:C8	2.50	0.46
2:B:93:ARG:HH21	4:D:102:LEU:HG	1.80	0.46
5:I:-15:DA:H1'	5:I:-14:DA:H5'	1.97	0.46
5:I:-16:DT:H4'	5:I:-15:DA:H5'	1.97	0.46
1:E:49:LYS:NZ	5:I:-65:DT:OP1	2.49	0.46
6:J:-20:DC:H2"	6:J:-19:DG:H5'	1.98	0.45
2:B:97:THR:HB	3:G:101:VAL:HG22	1.98	0.45
5:I:-62:DC:H2"	5:I:-61:DG:C8	2.52	0.45
3:C:40:TYR:O	4:D:79:SER:OG	2.31	0.45
5:I:12:DT:O4	6:J:-13:DA:N6	2.50	0.45
3:C:17:THR:HG22	3:C:19:SER:H	1.82	0.44
2:B:40:ARG:NH1	2:B:45:LYS:O	2.50	0.44
5:I:6:DC:O2	6:J:-5:DG:N2	2.50	0.44
3:G:17:THR:HG22	3:G:19:SER:H	1.82	0.44
4:D:37:SER:OG	4:D:38:TYR:N	2.49	0.44
2:F:40:ARG:NH1	2:F:45:LYS:O	2.50	0.44
5:I:-45:DA:N6	6:J:44:DT:O4	2.51	0.44
5:I:36:DC:H42	6:J:-36:DG:H1	1.67	0.43
5:I:-64:DC:H42	6:J:64:DG:H1	1.67	0.43
2:B:76:HIS:HD2	4:D:97:THR:HG21	1.84	0.42
3:G:65:GLU:HA	4:H:50:HIS:CE1	2.54	0.42
2:F:36:ARG:HH21	2:F:40:ARG:HH21	1.66	0.42
5:I:31:DT:H2"	5:I:32:DA:C8	2.53	0.42
3:G:65:GLU:HA	4:H:50:HIS:HE1	1.84	0.42
5:I:19:DC:H42	6:J:-19:DG:H1	1.65	0.42
6:J:-18:DG:H4'	6:J:-17:DT:H5'	2.02	0.42
2:F:37:ARG:NH2	6:J:-13:DA:OP2	2.42	0.42



Continued from previous page...

Atom-1	Atom-2	Interatomic	Clash
7100111-1	1100111-2	distance (Å)	overlap (Å)
5:I:-68:DG:H22	6:J:68:DC:H2'	1.85	0.42
6:J:-57:DT:H2"	6:J:-56:DG:C8	2.55	0.42
6:J:-63:DT:H1'	6:J:-62:DA:H5'	2.02	0.41
2:B:36:ARG:HH21	2:B:40:ARG:HH21	1.66	0.41
4:H:80:ARG:O	4:H:84:TYR:N	2.46	0.41
5:I:-21:DC:OP1	7:N:170:ARG:N	2.53	0.41
6:J:-2:DC:H2''	6:J:-1:DA:C8	2.55	0.41
5:I:-53:DG:N2	6:J:54:DT:O2	2.54	0.41
4:H:62:ILE:O	4:H:66:PHE:N	2.53	0.41
5:I:-8:DC:H2'	5:I:-7:DG:C4	2.56	0.41
3:C:21:ARG:NH2	5:I:-42:DT:OP1	2.39	0.41
6:J:38:DG:H2"	6:J:39:DA:H8	1.86	0.40
3:C:101:VAL:HG22	2:F:97:THR:HB	2.03	0.40
5:I:36:DC:N4	6:J:-37:DG:O6	2.54	0.40
6:J:-32:DT:H2"	6:J:-31:DA:H8	1.85	0.40
6:J:43:DA:H2'	6:J:44:DT:H71	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	89/160~(56%)	85 (96%)	4 (4%)	0	100	100
1	E	89/160 (56%)	85 (96%)	4 (4%)	0	100	100
2	В	78/103 (76%)	75 (96%)	3 (4%)	0	100	100
2	F	78/103 (76%)	75 (96%)	3 (4%)	0	100	100
3	С	105/130~(81%)	98 (93%)	7 (7%)	0	100	100
3	G	105/130 (81%)	98 (93%)	7 (7%)	0	100	100
4	D	92/126 (73%)	90 (98%)	2 (2%)	0	100	100



Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
4	Н	92/126 (73%)	88 (96%)	4 (4%)	0	100	100
7	N	118/668 (18%)	105 (89%)	13 (11%)	0	100	100
All	All	846/1706 (50%)	799 (94%)	47 (6%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percer	$_{ m tiles}$
1	A	77/135~(57%)	77 (100%)	0	100	100
1	E	77/135 (57%)	77 (100%)	0	100	100
2	В	65/79~(82%)	65 (100%)	0	100	100
2	F	65/79~(82%)	65 (100%)	0	100	100
3	\mathbf{C}	84/100 (84%)	84 (100%)	0	100	100
3	G	84/100 (84%)	84 (100%)	0	100	100
4	D	80/105 (76%)	80 (100%)	0	100	100
4	Η	80/105 (76%)	79 (99%)	1 (1%)	69	82
7	N	1/566~(0%)	1 (100%)	0	100	100
All	All	613/1404 (44%)	612 (100%)	1 (0%)	93	96

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	Н	32	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
4	D	85	ASN



5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.




6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-7293. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections (i)


6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

6.2 Central slices (i)

6.2.1 Primary map






The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)


6.3.1 Primary map



The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) (i)

6.4.1 Primary map




The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.



6.5 Orthogonal surface views (i)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.0014. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.6 Mask visualisation (i)


This section was not generated. No masks/segmentation were deposited.



7 Map analysis (i)

This section contains the results of statistical analysis of the map.


7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



7.2 Volume estimate (i)




The volume at the recommended contour level is $114~\mathrm{nm}^3$; this corresponds to an approximate mass of $103~\mathrm{kDa}$.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



7.3 Rotationally averaged power spectrum (i)



*Reported resolution corresponds to spatial frequency of 0.255 $\rm \mathring{A}^{-1}$



8 Fourier-Shell correlation (i)


This section was not generated. No FSC curve or half-maps provided.



9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-7293 and PDB model 6BUZ. Per-residue inclusion information can be found in section 3 on page 7.


9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.0014 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.




9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.0014).



9.4 Atom inclusion (i)



At the recommended contour level, 94% of all backbone atoms, 84% of all non-hydrogen atoms, are inside the map.



9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.0014) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8360	0.3720
A	0.7460	0.3910
В	0.7120	0.3840
С	0.7160	0.3920
D	0.7340	0.3920
E	0.7660	0.3880
F	0.7730	0.4040
G	0.7220	0.3830
Н	0.7340	0.3920
I	0.9280	0.3510
J	0.9240	0.3440
N	0.8690	0.4250



