



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 19, 2023 – 09:45 PM JST

PDB ID : 7BUG
Title : Reduced oxygenase of carbazole 1,9a-dioxygenase
Authors : Matsuzawa, J.; Wang, Y.X.; Suzuki-Minakuchi, C.; Nojiri, H.
Deposited on : 2020-04-06
Resolution : 1.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	392	 2% 85% 12% .
1	B	392	 2% 88% 10% .
1	C	392	 2% 88% 9% ..

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
11	MXE	A	411	-	-	X	-

2 Entry composition [i](#)

There are 13 unique types of molecules in this entry. The entry contains 10529 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Terminal oxygenase component of carbazole.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	383	3125	2000	528	584	13	0	7	0
1	B	383	3164	2028	534	588	14	0	13	0
1	C	383	3134	2005	532	584	13	0	8	0

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	385	LEU	-	expression tag	UNP Q84II6
A	386	GLU	-	expression tag	UNP Q84II6
A	387	HIS	-	expression tag	UNP Q84II6
A	388	HIS	-	expression tag	UNP Q84II6
A	389	HIS	-	expression tag	UNP Q84II6
A	390	HIS	-	expression tag	UNP Q84II6
A	391	HIS	-	expression tag	UNP Q84II6
A	392	HIS	-	expression tag	UNP Q84II6
B	385	LEU	-	expression tag	UNP Q84II6
B	386	GLU	-	expression tag	UNP Q84II6
B	387	HIS	-	expression tag	UNP Q84II6
B	388	HIS	-	expression tag	UNP Q84II6
B	389	HIS	-	expression tag	UNP Q84II6
B	390	HIS	-	expression tag	UNP Q84II6
B	391	HIS	-	expression tag	UNP Q84II6
B	392	HIS	-	expression tag	UNP Q84II6
C	385	LEU	-	expression tag	UNP Q84II6
C	386	GLU	-	expression tag	UNP Q84II6
C	387	HIS	-	expression tag	UNP Q84II6
C	388	HIS	-	expression tag	UNP Q84II6
C	389	HIS	-	expression tag	UNP Q84II6
C	390	HIS	-	expression tag	UNP Q84II6
C	391	HIS	-	expression tag	UNP Q84II6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	392	HIS	-	expression tag	UNP Q84II6

- Molecule 2 is FE2/S2 (INORGANIC) CLUSTER (three-letter code: FES) (formula: Fe₂S₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	Fe	S		
2	A	1	4	2	2	0	0
2	B	1	4	2	2	0	0
2	C	1	4	2	2	0	0


- Molecule 3 is FE (II) ION (three-letter code: FE2) (formula: Fe) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Fe		
3	A	1	1	1	0	0
3	B	1	1	1	0	0
3	C	1	1	1	0	0

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Mg 1 1	0	0
4	B	3	Total Mg 3 3	0	0
4	C	1	Total Mg 1 1	0	0

- Molecule 5 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 7 4 3	0	0
5	B	1	Total C O 7 4 3	0	0
5	B	1	Total C O 7 4 3	0	0
5	B	1	Total C O 7 4 3	0	0
5	C	1	Total C O 7 4 3	0	0
5	C	1	Total C O 7 4 3	0	0
5	C	1	Total C O 7 4 3	0	0
5	C	1	Total C O 7 4 3	0	0

- Molecule 6 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: C₆H₁₄O₄).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			10	6	4		
6	A	1	Total	C	O	0	0
			10	6	4		
6	B	1	Total	C	O	0	0
			10	6	4		
6	C	1	Total	C	O	0	0
			10	6	4		

- Molecule 7 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: $C_8H_{18}O_5$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			13	8	5		

- Molecule 8 is HEXAETHYLENE GLYCOL (three-letter code: P6G) (formula: $C_{12}H_{26}O_7$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			19	12	7		

- Molecule 9 is PENTAETHYLENE GLYCOL (three-letter code: 1PE) (formula: $C_{10}H_{22}O_6$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	A	1	Total	C	O	0	0
			16	10	6		
9	B	1	Total	C	O	0	0
			16	10	6		
9	C	1	Total	C	O	0	0
			16	10	6		

- Molecule 10 is CARBONATE ION (three-letter code: CO3) (formula: CO₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	A	1	Total	C	O	0	0
			4	1	3		
10	B	1	Total	C	O	0	0
			4	1	3		

- Molecule 11 is 2-METHOXYETHANOL (three-letter code: MXE) (formula: C₃H₈O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
11	A	1	Total	C	O	0	0
			5	3	2		
11	C	1	Total	C	O	0	0
			5	3	2		

- Molecule 12 is 2-(2-METHOXYETHOXY)ETHANOL (three-letter code: PG0) (formula: $C_5H_{12}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
12	A	1	Total	C	O	0	0
			8	5	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
12	B	1	Total	C	O	0	0
			8	5	3		
12	B	1	Total	C	O	0	0
			8	5	3		
12	C	1	Total	C	O	0	0
			8	5	3		


- Molecule 13 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
13	A	302	Total	O	0	0
			302	302		
13	B	299	Total	O	0	0
			299	299		
13	C	259	Total	O	0	0
			259	259		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Terminal oxygenase component of carbazole



- Molecule 1: Terminal oxygenase component of carbazole



- Molecule 1: Terminal oxygenase component of carbazole



4 Data and refinement statistics i

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, α , β , γ	91.94Å 91.94Å 243.46Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.01 – 1.60 45.97 – 1.60	Depositor EDS
% Data completeness (in resolution range)	100.0 (46.01-1.60) 100.0 (45.97-1.60)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.14 (at 1.60Å)	Xtrriage
Refinement program	REFMAC 5.8.0238	Depositor
R, R_{free}	0.154 , 0.188 0.165 , 0.197	Depositor DCC
R_{free} test set	7679 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	16.7	Xtrriage
Anisotropy	0.005	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 53.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.047 for h,-h-k,-l	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	10529	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, FES, PG0, P6G, PGE, MXE, CO3, FE2, 1PE, PG4, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.76	0/3219	0.90	2/4368 (0.0%)
1	B	0.76	1/3264 (0.0%)	0.93	2/4425 (0.0%)
1	C	0.73	0/3231	0.90	2/4382 (0.0%)
All	All	0.75	1/9714 (0.0%)	0.91	6/13175 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	291	GLU	CD-OE2	6.40	1.32	1.25

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	340	MET	CG-SD-CE	-6.20	90.28	100.20
1	A	340	MET	CG-SD-CE	-6.12	90.41	100.20
1	B	57	ARG	NE-CZ-NH1	5.40	123.00	120.30
1	A	224	ARG	CG-CD-NE	-5.38	100.49	111.80
1	C	295	TYR	CB-CG-CD1	-5.36	117.79	121.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3125	0	3045	36	0
1	B	3164	0	3102	23	0
1	C	3134	0	3061	28	0
2	A	4	0	0	0	0
2	B	4	0	0	0	0
2	C	4	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
4	A	1	0	0	0	0
4	B	3	0	0	0	0
4	C	1	0	0	0	0
5	A	7	0	10	0	0
5	B	21	0	30	2	0
5	C	28	0	40	5	0
6	A	20	0	28	1	0
6	B	10	0	14	0	0
6	C	10	0	14	3	0
7	A	13	0	18	0	0
8	A	19	0	26	5	0
9	A	16	0	22	0	0
9	B	16	0	22	0	0
9	C	16	0	22	3	0
10	A	4	0	0	0	0
10	B	4	0	0	0	0
11	A	5	0	8	4	0
11	C	5	0	8	0	0
12	A	8	0	12	1	0
12	B	16	0	24	4	0
12	C	8	0	12	1	0
13	A	302	0	0	5	0
13	B	299	0	0	9	0
13	C	259	0	0	5	0
All	All	10529	0	9518	93	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 93 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:330:ASN:HD21	12:B:412:PG0:H12	1.23	1.02

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:145:LEU:H	11:A:411:MXE:H33	1.25	0.97
1:A:279:ASP:OD2	1:A:302:LYS:HE2	1.84	0.77
1:A:145:LEU:N	11:A:411:MXE:H33	2.01	0.76
1:A:192:LEU:HA	1:A:324:MET:CE	2.16	0.75

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	388/392 (99%)	371 (96%)	15 (4%)	2 (0%)	29	11
1	B	394/392 (100%)	377 (96%)	16 (4%)	1 (0%)	41	21
1	C	389/392 (99%)	373 (96%)	15 (4%)	1 (0%)	41	21
All	All	1171/1176 (100%)	1121 (96%)	46 (4%)	4 (0%)	41	21

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	268	GLY
1	B	268	GLY
1	C	268	GLY
1	A	15	TRP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	336/339 (99%)	330 (98%)	6 (2%)	59	36
1	B	342/339 (101%)	333 (97%)	9 (3%)	46	21
1	C	337/339 (99%)	330 (98%)	7 (2%)	53	29
All	All	1015/1017 (100%)	993 (98%)	22 (2%)	52	27

5 of 22 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	383	SER
1	C	220	ASP
1	C	137	LEU
1	C	221	VAL
1	B	46	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	177	ASN
1	A	352	ASN
1	B	292	ASN
1	B	352	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 36 ligands modelled in this entry, 8 are monoatomic - leaving 28 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
9	1PE	A	409	-	15,15,15	0.74	0	14,14,14	0.75	0
9	1PE	B	408	-	15,15,15	0.59	0	14,14,14	0.84	0
5	PEG	B	409	-	6,6,6	0.24	0	5,5,5	0.11	0
5	PEG	A	404	-	6,6,6	0.44	0	5,5,5	0.36	0
12	PG0	A	412	-	7,7,7	0.25	0	6,6,6	0.53	0
11	MXE	C	410	-	4,4,4	0.34	0	3,3,3	0.27	0
6	PGE	C	405	-	9,9,9	0.22	0	8,8,8	0.14	0
2	FES	A	401	1	0,4,4	-	-	-	-	-
6	PGE	B	406	-	9,9,9	0.20	0	8,8,8	0.17	0
12	PG0	C	411	-	7,7,7	0.26	0	6,6,6	0.18	0
2	FES	C	401	1	0,4,4	-	-	-	-	-
9	1PE	C	407	-	15,15,15	0.76	0	14,14,14	1.03	2 (14%)
10	CO3	B	411	-	2,3,3	0.56	0	2,3,3	0.18	0
7	PG4	A	406	-	12,12,12	0.24	0	11,11,11	0.22	0
5	PEG	C	409	-	6,6,6	0.13	0	5,5,5	0.31	0
2	FES	B	401	1	0,4,4	-	-	-	-	-
12	PG0	B	412	-	7,7,7	0.37	0	6,6,6	0.30	0
12	PG0	B	413	-	7,7,7	0.35	0	6,6,6	0.43	0
6	PGE	A	405	-	9,9,9	0.35	0	8,8,8	0.22	0
5	PEG	B	407	-	6,6,6	0.23	0	5,5,5	0.23	0
8	P6G	A	408	-	18,18,18	0.75	0	17,17,17	0.86	0
5	PEG	C	406	-	6,6,6	0.15	0	5,5,5	0.27	0
10	CO3	A	410	-	2,3,3	0.12	0	2,3,3	0.42	0
11	MXE	A	411	-	4,4,4	0.47	0	3,3,3	1.12	0
5	PEG	C	408	-	6,6,6	0.17	0	5,5,5	0.23	0
6	PGE	A	407	-	9,9,9	0.30	0	8,8,8	0.29	0
5	PEG	B	410	-	6,6,6	0.27	0	5,5,5	0.25	0
5	PEG	C	404	-	6,6,6	0.36	0	5,5,5	0.37	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	1PE	A	409	-	-	6/13/13/13	-
9	1PE	B	408	-	-	5/13/13/13	-
5	PEG	B	409	-	-	2/4/4/4	-
5	PEG	A	404	-	-	1/4/4/4	-
12	PG0	A	412	-	-	2/5/5/5	-
11	MXE	C	410	-	-	1/2/2/2	-
6	PGE	C	405	-	-	4/7/7/7	-
2	FES	A	401	1	-	-	0/1/1/1
6	PGE	B	406	-	-	1/7/7/7	-
12	PG0	C	411	-	-	1/5/5/5	-
2	FES	C	401	1	-	-	0/1/1/1
9	1PE	C	407	-	-	8/13/13/13	-
7	PG4	A	406	-	-	3/10/10/10	-
5	PEG	C	409	-	-	1/4/4/4	-
12	PG0	B	412	-	-	2/5/5/5	-
2	FES	B	401	1	-	-	0/1/1/1
12	PG0	B	413	-	-	3/5/5/5	-
6	PGE	A	405	-	-	2/7/7/7	-
5	PEG	B	407	-	-	3/4/4/4	-
8	P6G	A	408	-	-	11/16/16/16	-
5	PEG	C	406	-	-	2/4/4/4	-
11	MXE	A	411	-	-	2/2/2/2	-
5	PEG	C	408	-	-	3/4/4/4	-
6	PGE	A	407	-	-	4/7/7/7	-
5	PEG	B	410	-	-	3/4/4/4	-
5	PEG	C	404	-	-	0/4/4/4	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	C	407	1PE	OH3-C23-C13	2.49	121.62	110.39
9	C	407	1PE	OH3-C22-C12	2.09	119.26	110.07

There are no chirality outliers.

5 of 70 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	A	407	PGE	C3-C4-O3-C5
9	C	407	1PE	OH7-C16-C26-OH6
5	B	410	PEG	C1-C2-O2-C3
8	A	408	P6G	O7-C8-C9-O10
8	A	408	P6G	O13-C14-C15-O16

There are no ring outliers.

13 monomers are involved in 29 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
12	A	412	PG0	1	0
6	C	405	PGE	3	0
12	C	411	PG0	1	0
9	C	407	1PE	3	0
5	C	409	PEG	1	0
12	B	412	PG0	4	0
5	B	407	PEG	1	0
8	A	408	P6G	5	0
5	C	406	PEG	3	0
11	A	411	MXE	4	0
5	C	408	PEG	1	0
6	A	407	PGE	1	0
5	B	410	PEG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	383/392 (97%)	-0.35	3 (0%) 86 86	9, 19, 43, 58	0
1	B	383/392 (97%)	-0.35	6 (1%) 72 71	10, 18, 38, 67	0
1	C	383/392 (97%)	-0.37	7 (1%) 68 67	13, 21, 40, 61	0
All	All	1149/1176 (97%)	-0.36	16 (1%) 75 75	9, 19, 42, 67	0

The worst 5 of 16 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	221	VAL	4.6
1	B	384	GLY	4.2
1	C	234	HIS	3.2
1	A	253	TYR	3.0
1	C	221	VAL	2.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
6	PGE	A	405	10/10	0.71	0.21	41,45,47,53	0
5	PEG	C	406	7/7	0.72	0.13	49,51,54,57	0
5	PEG	A	404	7/7	0.72	0.18	33,39,43,45	0
9	1PE	A	409	16/16	0.72	0.13	41,47,53,53	0
5	PEG	B	410	7/7	0.74	0.14	46,47,49,49	0
12	PG0	B	413	8/8	0.77	0.12	44,46,48,50	0
5	PEG	B	407	7/7	0.78	0.13	44,52,54,55	0
5	PEG	B	409	7/7	0.79	0.13	41,49,51,56	0
6	PGE	A	407	10/10	0.79	0.22	45,49,53,56	0
5	PEG	C	409	7/7	0.80	0.11	46,47,48,53	0
6	PGE	C	405	10/10	0.80	0.14	43,48,54,56	0
12	PG0	A	412	8/8	0.81	0.13	36,40,43,50	0
8	P6G	A	408	19/19	0.82	0.22	44,47,61,62	0
9	1PE	C	407	16/16	0.83	0.18	37,41,48,49	0
5	PEG	C	408	7/7	0.83	0.23	49,52,54,59	0
9	1PE	B	408	16/16	0.83	0.13	43,50,60,60	0
11	MXE	C	410	5/5	0.87	0.09	43,45,49,55	0
7	PG4	A	406	13/13	0.88	0.18	25,36,50,54	0
12	PG0	B	412	8/8	0.89	0.10	19,26,35,40	0
11	MXE	A	411	5/5	0.89	0.14	25,26,36,37	0
10	CO3	B	411	4/4	0.90	0.17	34,36,38,39	0
4	MG	B	403	1/1	0.90	0.21	39,39,39,39	0
10	CO3	A	410	4/4	0.90	0.10	21,25,29,36	0
6	PGE	B	406	10/10	0.91	0.13	35,40,48,48	0
5	PEG	C	404	7/7	0.91	0.11	25,26,39,50	0
12	PG0	C	411	8/8	0.92	0.10	25,27,42,45	0
4	MG	B	405	1/1	0.95	0.15	29,29,29,29	0
4	MG	A	403	1/1	0.99	0.10	18,18,18,18	0
2	FES	B	401	4/4	0.99	0.07	13,13,13,14	0
4	MG	B	404	1/1	0.99	0.14	20,20,20,20	0
2	FES	C	401	4/4	0.99	0.06	13,15,15,15	0
4	MG	C	403	1/1	0.99	0.08	21,21,21,21	0
3	FE2	A	402	1/1	0.99	0.05	19,19,19,19	0
2	FES	A	401	4/4	1.00	0.07	9,9,9,9	0
3	FE2	B	402	1/1	1.00	0.04	13,13,13,13	0
3	FE2	C	402	1/1	1.00	0.04	16,16,16,16	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around P6G A 408:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FE2 A 402:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around FE2 B 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.