



Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 06:12 pm BST

PDB ID : 3BWJ
Title : Complex of PKA with the bisubstrate protein kinase inhibitor lead compound Arc-1034
Authors : Lavogina, D.; Koenig, N.; Uri, A.; Bossemeyer, D.
Deposited on : 2008-01-09
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	350	

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 3098 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called cAMP-dependent protein kinase, alpha-catalytic subunit.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
1	A	341	2823	1824	472	515	3	9	0	0	0

- Molecule 2 is (2S,3S,4R,5R)-5-(6-amino-9H-purin-9-yl)-N-(6-[[[(1R)-4-carbamimidamido-1-[[[(1R)-4-carbamimidamido-1-carbamoylbutyl]carbamoyl]butyl]amino]-6-oxohexyl]-3,4-dihydroxytetrahydrofuran-2-carboxamide (three-letter code: ARX) (formula: C₂₈H₄₇N₁₅O₇).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	50	28	15	7	0	0
2	A	1	50	28	15	7	0	0


- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	175	Total 175	O 175	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: cAMP-dependent protein kinase, alpha-catalytic subunit

Chain A: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	71.19Å 72.63Å 77.22Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	52.93 – 2.30 24.80 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.9 (52.93-2.30) 100.0 (24.80-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.02 (at 2.31Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.199 , 0.249 0.194 , 0.244	Depositor DCC
R_{free} test set	939 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å ²)	26.0	Xtrriage
Anisotropy	0.223	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 41.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.020 for k,h,-l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3098	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, ARX, SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.79	0/2862	0.76	0/3853

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2823	0	2790	28	1
2	A	100	0	90	9	0
3	A	175	0	0	8	2
All	All	3098	0	2880	36	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:352:ARX:HN20	2:A:352:ARX:H8	1.27	0.99

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:96:GLN:HG2	3:A:592:HOH:O	1.67	0.93
2:A:352:ARX:H8	2:A:352:ARX:N20	1.94	0.82
1:A:92:LYS:NZ	1:A:350:PHE:OXT	2.18	0.73
1:A:104:VAL:HG23	3:A:546:HOH:O	1.91	0.69
1:A:336:ARG:HH11	1:A:336:ARG:CG	2.08	0.67
1:A:158:HIS:HE1	1:A:220:ASP:OD2	1.80	0.65
1:A:346:GLU:OE1	1:A:346:GLU:N	2.31	0.62
1:A:173:LEU:HD11	2:A:351:ARX:H2'	1.83	0.60
1:A:295:LYS:HE2	3:A:569:HOH:O	2.01	0.59
1:A:104:VAL:CG2	3:A:546:HOH:O	2.49	0.58
2:A:352:ARX:HN20	2:A:352:ARX:C8	2.11	0.56
1:A:103:LEU:HD22	1:A:185:PHE:HZ	1.69	0.55
1:A:56:ARG:NH1	1:A:333:GLU:OE2	2.44	0.51
1:A:336:ARG:HG2	1:A:336:ARG:HH11	1.76	0.51
1:A:103:LEU:HD22	1:A:185:PHE:CZ	2.46	0.51
2:A:352:ARX:H2	3:A:417:HOH:O	2.11	0.49
1:A:62:HIS:HD2	1:A:65:THR:H	1.61	0.49
1:A:42:GLN:O	1:A:63:MET:HG2	2.14	0.48
1:A:226:VAL:HG13	1:A:237:PRO:HD2	1.97	0.47
2:A:351:ARX:HN76	2:A:351:ARX:H79A	1.80	0.47
1:A:28:LYS:HE3	3:A:597:HOH:O	2.14	0.46
1:A:308:ARG:NH2	3:A:562:HOH:O	2.47	0.46
1:A:46:ILE:HD12	1:A:59:LEU:HD13	1.96	0.46
1:A:336:ARG:HG2	1:A:336:ARG:NH1	2.31	0.45
1:A:45:ARG:HD3	1:A:335:ILE:HG13	1.98	0.44
1:A:196:TRP:HB2	3:A:549:HOH:O	2.16	0.44
1:A:238:PHE:CE1	1:A:255:VAL:HG22	2.52	0.43
2:A:351:ARX:O5'	2:A:351:ARX:H8	2.18	0.43
1:A:80:VAL:HG22	1:A:85:ILE:HD11	2.01	0.43
2:A:352:ARX:H22	2:A:352:ARX:H8	2.01	0.43
2:A:351:ARX:H25	2:A:351:ARX:H22	1.66	0.42
1:A:28:LYS:HE2	1:A:28:LYS:HB2	1.75	0.42
1:A:238:PHE:CZ	1:A:255:VAL:HG22	2.55	0.41
1:A:336:ARG:NH1	1:A:336:ARG:CG	2.73	0.41
1:A:101:PRO:O	1:A:308:ARG:NH2	2.55	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:549:HOH:O	3:A:556:HOH:O 3_645]	1.51	0.69

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:133:ARG:O	3:A:549:HOH:O[3_655]	1.89	0.31

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	336/350 (96%)	325 (97%)	10 (3%)	1 (0%)	41 50

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	33	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	299/302 (99%)	280 (94%)	19 (6%)	17 23

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	GLN
1	A	14	SER
1	A	21	LYS
1	A	40	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	44	GLU
1	A	59	LEU
1	A	130	SER
1	A	160	LEU
1	A	173	LEU
1	A	191	VAL
1	A	217	LYS
1	A	242	GLN
1	A	268	LEU
1	A	285	LYS
1	A	317	LYS
1	A	336	ARG
1	A	339	ILE
1	A	348	SER
1	A	349	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	42	GLN
1	A	62	HIS
1	A	68	HIS
1	A	113	ASN
1	A	158	HIS
1	A	181	GLN
1	A	286	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	SEP	A	338	1	8,9,10	1.63	1 (12%)	8,12,14	2.26	2 (25%)
1	SEP	A	139	1	8,9,10	1.68	3 (37%)	8,12,14	1.04	0
1	TPO	A	197	1	8,10,11	1.14	1 (12%)	10,14,16	1.13	1 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SEP	A	338	1	-	5/5/8/10	-
1	SEP	A	139	1	-	0/5/8/10	-
1	TPO	A	197	1	-	2/9/11/13	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	338	SEP	P-O1P	3.48	1.61	1.50
1	A	139	SEP	P-O1P	3.14	1.60	1.50
1	A	197	TPO	P-O1P	2.28	1.57	1.50
1	A	139	SEP	P-O2P	2.13	1.63	1.54
1	A	139	SEP	P-O3P	2.10	1.62	1.54

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	338	SEP	OG-CB-CA	5.12	113.13	108.14
1	A	338	SEP	O2P-P-OG	3.32	115.56	106.73
1	A	197	TPO	O2P-P-O1P	-2.10	102.45	110.68

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	338	SEP	N-CA-CB-OG
1	A	338	SEP	CA-CB-OG-P
1	A	338	SEP	CB-OG-P-O1P

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
1	A	197	TPO	O-C-CA-CB
1	A	197	TPO	CB-OG1-P-O3P
1	A	338	SEP	CB-OG-P-O3P
1	A	338	SEP	CB-OG-P-O2P

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	ARX	A	351	-	49,52,52	0.92	3 (6%)	55,70,70	1.46	10 (18%)
2	ARX	A	352	-	49,52,52	0.97	3 (6%)	55,70,70	1.40	7 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ARX	A	351	-	-	14/43/63/63	0/3/3/3
2	ARX	A	352	-	-	6/43/63/63	0/3/3/3

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	352	ARX	C5-C4	3.37	1.49	1.40
2	A	352	ARX	C2-N3	2.64	1.36	1.32
2	A	352	ARX	O4'-C1'	2.42	1.44	1.41
2	A	351	ARX	O4'-C1'	2.10	1.44	1.41
2	A	351	ARX	C2-N3	2.08	1.35	1.32
2	A	351	ARX	C5-C4	2.03	1.46	1.40

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	351	ARX	C32-N31-C30	4.48	131.26	121.67
2	A	352	ARX	C4'-C5'-N20	4.05	120.50	115.47
2	A	351	ARX	C3'-C2'-C1'	3.75	106.63	100.98
2	A	352	ARX	C2'-C3'-C4'	3.36	105.55	101.64
2	A	352	ARX	C3'-C2'-C1'	3.12	105.67	100.98
2	A	351	ARX	C32-C75-N76	2.79	121.46	116.69
2	A	351	ARX	C4-C5-N7	-2.70	106.58	109.40
2	A	352	ARX	C32-N31-C30	2.65	127.36	121.67
2	A	352	ARX	C25-C26-N27	2.62	120.37	115.83
2	A	351	ARX	N3-C2-N1	-2.58	124.64	128.68
2	A	352	ARX	O4'-C4'-C5'	2.52	115.31	111.11
2	A	351	ARX	C75-C32-N31	2.46	116.14	110.18
2	A	351	ARX	C78-C32-C75	2.44	115.87	110.21
2	A	351	ARX	N6-C6-N1	2.40	123.56	118.57
2	A	351	ARX	C21-N20-C5'	-2.37	118.37	122.59
2	A	352	ARX	C79-C78-C32	-2.23	107.05	113.92
2	A	351	ARX	O28-C26-C25	-2.12	118.14	122.02

There are no chirality outliers.

All (20) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	351	ARX	N43-C41-N40-C39
2	A	351	ARX	N42-C41-N40-C39
2	A	351	ARX	C78-C32-C75-O77
2	A	351	ARX	C78-C32-C75-N76
2	A	351	ARX	N93-C86-N77-C85
2	A	351	ARX	N94-C86-N77-C85
2	A	351	ARX	O35-C30-N31-C32
2	A	351	ARX	C29-C30-N31-C32
2	A	352	ARX	N20-C21-C22-C23
2	A	352	ARX	C23-C24-C25-C26

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	A	352	ARX	C21-C22-C23-C24
2	A	352	ARX	C78-C79-C85-N77
2	A	351	ARX	C37-C38-C39-N40
2	A	351	ARX	C29-C37-C38-C39
2	A	351	ARX	C78-C79-C85-N77
2	A	351	ARX	C22-C23-C24-C25
2	A	352	ARX	C24-C25-C26-O28
2	A	352	ARX	C24-C25-C26-N27
2	A	351	ARX	C79-C85-N77-C86
2	A	351	ARX	O4'-C4'-C5'-N20

There are no ring outliers.

2 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	351	ARX	4	0
2	A	352	ARX	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	338/350 (96%)	-0.24	13 (3%) 40 47	9, 21, 47, 67	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	34	ALA	5.0
1	A	35	GLN	4.2
1	A	12	GLN	3.8
1	A	11	GLU	3.8
1	A	318	PHE	3.7
1	A	331	GLU	2.5
1	A	10	SER	2.5
1	A	36	ASN	2.5
1	A	13	GLU	2.4
1	A	345	LYS	2.3
1	A	336	ARG	2.2
1	A	242	GLN	2.2
1	A	337	VAL	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	SEP	A	338	10/11	0.88	0.13	46,47,50,51	0
1	SEP	A	139	10/11	0.90	0.14	18,21,32,33	0
1	TPO	A	197	11/12	0.99	0.09	12,14,17,18	0

6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ARX	A	351	50/50	0.88	0.17	12,21,60,62	0
2	ARX	A	352	50/50	0.90	0.13	8,27,45,48	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.