



wwPDB X-ray Structure Validation Summary Report ⓘ

May 13, 2020 – 12:08 am BST

PDB ID : 5BY9
Title : The crystal structure of polyglycylated 14-3-3 protein from *Giardia intestinalis*
Authors : Fiorillo, A.; Ilari, A.; Lalle, M.
Deposited on : 2015-06-10
Resolution : 4.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 4.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1087 (4.30-3.70)
Clashscore	141614	1148 (4.30-3.70)
Ramachandran outliers	138981	1108 (4.30-3.70)
Sidechain outliers	138945	1099 (4.30-3.70)
RSRZ outliers	127900	1028 (4.34-3.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	256	89% 8%
1	B	256	87% 5% 8%
1	C	256	83% 7% 9%
1	D	256	86% 5% 9%

2 Entry composition i

There is only 1 type of molecule in this entry. The entry contains 7633 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 14-3-3 protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	236	Total 1921	C 1213	N 330	O 370	S 8	0	0	0
1	B	235	Total 1915	C 1209	N 329	O 369	S 8	0	0	0
1	C	232	Total 1888	C 1190	N 325	O 365	S 8	0	0	0
1	D	234	Total 1909	C 1206	N 328	O 367	S 8	0	0	0

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	247	GLY	-	expression tag	UNP Q2QBT8
A	248	GLY	-	expression tag	UNP Q2QBT8
A	249	GLY	-	expression tag	UNP Q2QBT8
A	250	GLY	-	expression tag	UNP Q2QBT8
A	251	GLY	-	expression tag	UNP Q2QBT8
A	252	GLY	-	expression tag	UNP Q2QBT8
A	253	GLY	-	expression tag	UNP Q2QBT8
A	254	GLY	-	expression tag	UNP Q2QBT8
A	255	GLY	-	expression tag	UNP Q2QBT8
A	256	GLY	-	expression tag	UNP Q2QBT8
B	247	GLY	-	expression tag	UNP Q2QBT8
B	248	GLY	-	expression tag	UNP Q2QBT8
B	249	GLY	-	expression tag	UNP Q2QBT8
B	250	GLY	-	expression tag	UNP Q2QBT8
B	251	GLY	-	expression tag	UNP Q2QBT8
B	252	GLY	-	expression tag	UNP Q2QBT8
B	253	GLY	-	expression tag	UNP Q2QBT8
B	254	GLY	-	expression tag	UNP Q2QBT8
B	255	GLY	-	expression tag	UNP Q2QBT8
B	256	GLY	-	expression tag	UNP Q2QBT8
C	247	GLY	-	expression tag	UNP Q2QBT8

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	248	GLY	-	expression tag	UNP Q2QBT8
C	249	GLY	-	expression tag	UNP Q2QBT8
C	250	GLY	-	expression tag	UNP Q2QBT8
C	251	GLY	-	expression tag	UNP Q2QBT8
C	252	GLY	-	expression tag	UNP Q2QBT8
C	253	GLY	-	expression tag	UNP Q2QBT8
C	254	GLY	-	expression tag	UNP Q2QBT8
C	255	GLY	-	expression tag	UNP Q2QBT8
C	256	GLY	-	expression tag	UNP Q2QBT8
D	247	GLY	-	expression tag	UNP Q2QBT8
D	248	GLY	-	expression tag	UNP Q2QBT8
D	249	GLY	-	expression tag	UNP Q2QBT8
D	250	GLY	-	expression tag	UNP Q2QBT8
D	251	GLY	-	expression tag	UNP Q2QBT8
D	252	GLY	-	expression tag	UNP Q2QBT8
D	253	GLY	-	expression tag	UNP Q2QBT8
D	254	GLY	-	expression tag	UNP Q2QBT8
D	255	GLY	-	expression tag	UNP Q2QBT8
D	256	GLY	-	expression tag	UNP Q2QBT8

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 14-3-3 protein

Chain A: 





- Molecule 1: 14-3-3 protein

Chain B: 




- Molecule 1: 14-3-3 protein

Chain C: 



- Molecule 1: 14-3-3 protein

Chain D: 



4 Data and refinement statistics i

Property	Value	Source
Space group	P 32	Depositor
Cell constants a, b, c, α , β , γ	100.21Å 100.21Å 140.96Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	36.95 – 4.00 36.95 – 4.00	Depositor EDS
% Data completeness (in resolution range)	92.6 (36.95-4.00) 92.8 (36.95-4.00)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.09 (at 3.99Å)	Xtriage
Refinement program	REFMAC 5.8.0107	Depositor
R, R_{free}	0.233 , 0.233 0.232 , 0.233	Depositor DCC
R_{free} test set	611 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	95.9	Xtriage
Anisotropy	0.686	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 64.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.39$, $\langle L^2 \rangle = 0.21$	Xtriage
Estimated twinning fraction	0.125 for -h,-k,l 0.307 for h,-h-k,-l 0.126 for -k,-h,-l	Xtriage
Reported twinning fraction	0.571 for H, K, L 0.429 for K, H, -L	Depositor
Outliers	0 of 12369 reflections	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	7633	wwPDB-VP
Average B, all atoms (Å ²)	117.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.39	0/1954	0.54	0/2632
1	B	0.40	0/1948	0.55	0/2624
1	C	0.39	0/1919	0.54	0/2582
1	D	0.40	0/1942	0.55	0/2615
All	All	0.39	0/7763	0.55	0/10453

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1921	0	1908	3	0
1	B	1915	0	1901	4	0
1	C	1888	0	1877	5	0
1	D	1909	0	1896	2	0
All	All	7633	0	7582	14	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 14 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:223:LEU:HA	1:C:226:GLN:HE21	1.72	0.53
1:B:80:ASN:HD22	1:B:83:ARG:H	1.57	0.52
1:B:49:SER:HB2	1:B:125:PHE:HZ	1.80	0.46
1:B:210:LEU:HD12	1:B:213:LEU:HD12	2.00	0.44
1:A:171:HIS:HE1	1:A:173:ILE:HD12	1.82	0.44

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	234/256 (91%)	232 (99%)	2 (1%)	0	100	100
1	B	233/256 (91%)	233 (100%)	0	0	100	100
1	C	230/256 (90%)	230 (100%)	0	0	100	100
1	D	232/256 (91%)	231 (100%)	1 (0%)	0	100	100
All	All	929/1024 (91%)	926 (100%)	3 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	205/212 (97%)	202 (98%)	3 (2%)	65	80
1	B	205/212 (97%)	200 (98%)	5 (2%)	49	69

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	202/212 (95%)	193 (96%)	9 (4%)	27	55
1	D	204/212 (96%)	192 (94%)	12 (6%)	19	48
All	All	816/848 (96%)	787 (96%)	29 (4%)	35	61

5 of 29 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	181	PHE
1	C	229	ARG
1	D	194	ARG
1	C	204	ASP
1	D	16	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	171	HIS
1	C	226	GLN
1	B	86	GLN
1	A	130	GLN
1	B	80	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	236/256 (92%)	0.04	0 100 100	103, 119, 120, 120	0
1	B	235/256 (91%)	-0.01	0 100 100	105, 120, 120, 120	0
1	C	232/256 (90%)	0.04	0 100 100	105, 119, 120, 120	0
1	D	234/256 (91%)	-0.01	0 100 100	100, 120, 120, 120	0
All	All	937/1024 (91%)	0.01	0 100 100	100, 120, 120, 120	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.