



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 25, 2024 – 03:05 AM EST

PDB ID : 5BYL  
Title : Aminoglycoside Phosphotransferase (2'')-Ia (CTD of AAC(6')-Ie/APH(2'')-Ia)  
 ) in complex with GMPPCP and Magnesium  
Authors : Caldwell, S.J.; Berghuis, A.M.  
Deposited on : 2015-06-10  
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1479 (2.16-2.16)
Clashscore	141614	1585 (2.16-2.16)
Ramachandran outliers	138981	1560 (2.16-2.16)
Sidechain outliers	138945	1559 (2.16-2.16)
RSRZ outliers	127900	1456 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	305	 90% 7% •
1	B	305	 90% 7% •
1	C	305	 90% 7% •
1	D	305	 91% 6% •

## 2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 11080 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bifunctional AAC/APH.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	295	Total 2452	C 1566	N 378	O 498	S 10	0	5	0
1	B	298	Total 2486	C 1588	N 381	O 507	S 10	0	8	0
1	C	296	Total 2442	C 1558	N 381	O 493	S 10	0	2	0
1	D	296	Total 2447	C 1563	N 379	O 495	S 10	0	2	0

- Molecule 2 is PHOSPHOMETHYLPHOSPHONIC ACID GUANYLATE ESTER (three-letter code: GCP) (formula:  $C_{11}H_{18}N_5O_{13}P_3$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
2	A	1	Total 41	C 12	N 5	O 19	P 5	0	1
2	B	1	Total 32	C 11	N 5	O 13	P 3	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	C	1	Total	C	N	O	P	0	1
			41	12	5	19	5		
2	D	1	Total	C	N	O	P	0	0
			32	11	5	13	3		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Mg	0	0
			2	2		
3	B	2	Total	Mg	0	0
			2	2		
3	C	3	Total	Mg	0	0
			3	3		
3	D	2	Total	Mg	0	0
			2	2		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cl	0	0
			1	1		
4	B	1	Total	Cl	0	0
			1	1		
4	C	1	Total	Cl	0	0
			1	1		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	B	1	Total C O 6 3 3	0	0
5	C	1	Total C O 6 3 3	0	0
5	C	1	Total C O 6 3 3	0	0
5	D	1	Total C O 6 3 3	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	275	Total O 275 275	0	0
6	B	326	Total O 326 326	0	0
6	C	255	Total O 255 255	0	0
6	D	203	Total O 203 203	0	0



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Bifunctional AAC/APH



- Molecule 1: Bifunctional AAC/APH



- Molecule 1: Bifunctional AAC/APH



- Molecule 1: Bifunctional AAC/APH



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.29Å 99.90Å 93.62Å 90.00° 105.09° 90.00°	Depositor
Resolution (Å)	58.92 – 2.15 49.95 – 2.15	Depositor EDS
% Data completeness (in resolution range)	94.9 (58.92-2.15) 94.9 (49.95-2.15)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.62 (at 2.16Å)	Xtrriage
Refinement program	REFMAC 5.8.0073	Depositor
R, $R_{free}$	0.156 , 0.198 0.164 , 0.204	Depositor DCC
$R_{free}$ test set	4159 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	39.6	Xtrriage
Anisotropy	0.060	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 44.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.015 for l,-k,h	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	11080	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	48.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.74% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CL, GCP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.85	1/2510 (0.0%)	0.81	2/3386 (0.1%)
1	B	0.91	3/2554 (0.1%)	0.81	1/3446 (0.0%)
1	C	0.85	1/2491 (0.0%)	0.82	2/3359 (0.1%)
1	D	0.75	0/2497	0.78	0/3367
All	All	0.85	5/10052 (0.0%)	0.81	5/13558 (0.0%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	377	CYS	CB-SG	-5.93	1.72	1.81
1	B	191	TYR	CE2-CZ	-5.90	1.30	1.38
1	A	377	CYS	CB-SG	-5.22	1.73	1.81
1	B	236	GLY	C-O	5.04	1.31	1.23
1	B	445	GLU	CD-OE1	5.02	1.31	1.25

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	357	ARG	NE-CZ-NH2	-6.54	117.03	120.30
1	A	357	ARG	NE-CZ-NH2	-6.38	117.11	120.30
1	A	404	CYS	CA-CB-SG	-6.00	103.20	114.00
1	C	404	CYS	CA-CB-SG	-5.32	104.42	114.00
1	C	357	ARG	NE-CZ-NH2	-5.20	117.70	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2452	0	2365	10	0
1	B	2486	0	2383	9	0
1	C	2442	0	2342	12	0
1	D	2447	0	2346	10	0
2	A	41	0	4	2	0
2	B	32	0	14	0	0
2	C	41	0	4	2	0
2	D	32	0	14	0	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
3	C	3	0	0	0	0
3	D	2	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
5	A	12	0	16	0	0
5	B	6	0	8	0	0
5	C	12	0	16	0	0
5	D	6	0	8	0	0
6	A	275	0	0	3	1
6	B	326	0	0	3	0
6	C	255	0	0	6	0
6	D	203	0	0	2	1
All	All	11080	0	9520	44	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:500[A]:GCP:O2G	6:C:901:HOH:O	1.92	0.86
2:A:500[B]:GCP:O1G	6:A:1038:HOH:O	1.94	0.86
2:C:500[A]:GCP:PG	6:C:901:HOH:O	2.51	0.68
1:A:367:LYS:HE2	6:A:1238:HOH:O	1.98	0.62

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:297:LEU:HD23	1:C:387:ARG:HD3	1.81	0.62
1:D:326:GLN:NE2	6:D:1013:HOH:O	2.31	0.55
1:B:189:MET:HG2	1:B:217:TYR:CE1	2.41	0.55
1:C:189:MET:HG2	1:C:217:TYR:CE1	2.42	0.54
1:D:189:MET:HG2	1:D:217:TYR:CE1	2.41	0.54
1:C:186:VAL:HG13	1:C:205:ILE:HG23	1.90	0.53
1:A:214:SER:OG	2:A:500[B]:GCP:H3B2	2.08	0.52
1:B:436:LYS:NZ	6:B:1113:HOH:O	2.41	0.52
1:A:186:VAL:HG13	1:A:205:ILE:HG23	1.92	0.52
1:D:290:MET:HE3	1:D:294:GLU:HB3	1.91	0.51
1:A:189:MET:HG2	1:A:217:TYR:CE1	2.45	0.51
1:B:186:VAL:HG13	1:B:205:ILE:HG23	1.92	0.50
1:A:208:ILE:HD11	1:A:224:ILE:HD12	1.94	0.49
1:C:436:LYS:NZ	6:C:1113:HOH:O	2.44	0.49
1:C:300:ARG:NH2	6:C:1359:HOH:O	2.46	0.48
1:C:198:ASP:O	6:C:1027:HOH:O	2.20	0.48
1:D:208:ILE:HD11	1:D:224:ILE:HD12	1.94	0.48
1:B:290:MET:HE3	1:B:294:GLU:HB3	1.95	0.47
1:C:471:GLU:CG	1:C:475:ARG:HH11	2.28	0.46
1:C:240:GLU:HG2	1:C:244:TYR:CE2	2.51	0.46
1:D:240:GLU:HG2	1:D:244:TYR:CE2	2.51	0.46
1:A:290:MET:HE3	1:A:294:GLU:HB3	1.98	0.46
1:B:240:GLU:HG2	1:B:244:TYR:CE2	2.51	0.45
1:B:189:MET:SD	1:B:227:THR:HG21	2.57	0.44
1:C:290:MET:HE3	1:C:294:GLU:HB3	1.99	0.44
1:A:189:MET:SD	1:A:227:THR:HG21	2.59	0.42
1:D:471:GLU:CG	1:D:475:ARG:HH11	2.33	0.42
1:D:208:ILE:CD1	1:D:224:ILE:HD12	2.50	0.42
1:B:208:ILE:HD11	1:B:224:ILE:HD12	2.00	0.42
1:B:298:LEU:HA	6:B:1268:HOH:O	2.20	0.42
1:D:189:MET:SD	1:D:227:THR:HG21	2.60	0.41
1:A:240:GLU:HG2	1:A:244:TYR:CE2	2.55	0.41
1:C:414:GLU:HG2	6:C:1234:HOH:O	2.21	0.41
1:D:212:TYR:HB3	1:D:455:TYR:OH	2.21	0.41
1:B:301:ASP:OD1	6:B:1273:HOH:O	2.22	0.41
1:C:189:MET:SD	1:C:227:THR:HG21	2.61	0.41
1:C:471:GLU:HG3	1:C:475:ARG:HH11	1.85	0.41
1:A:359:ASN:ND2	6:A:1050:HOH:O	2.49	0.40
1:D:182:ASN:HB3	6:D:1229:HOH:O	2.20	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:1534:HOH:O	6:D:1312:HOH:O[2_646]	2.07	0.13

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	296/305 (97%)	285 (96%)	10 (3%)	1 (0%)	41 37
1	B	302/305 (99%)	289 (96%)	12 (4%)	1 (0%)	41 37
1	C	294/305 (96%)	284 (97%)	9 (3%)	1 (0%)	41 37
1	D	294/305 (96%)	284 (97%)	9 (3%)	1 (0%)	41 37
All	All	1186/1220 (97%)	1142 (96%)	40 (3%)	4 (0%)	41 37

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	400	ILE
1	B	400	ILE
1	A	400	ILE
1	C	400	ILE

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	274/281 (98%)	272 (99%)	2 (1%)	84 89
1	B	277/281 (99%)	274 (99%)	3 (1%)	73 78

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	269/281 (96%)	266 (99%)	3 (1%)	73	78
1	D	270/281 (96%)	269 (100%)	1 (0%)	91	93
All	All	1090/1124 (97%)	1081 (99%)	9 (1%)	81	86

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	221	ASN
1	A	230	SER
1	B	221	ASN
1	B	228	LYS
1	B	285	GLU
1	C	208	ILE
1	C	221	ASN
1	C	297	LEU
1	D	221	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	467	ASN
1	B	467	ASN
1	D	420	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 24 ligands modelled in this entry, 12 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	GOL	A	806	-	5,5,5	0.69	0	5,5,5	0.34	0
2	GCP	A	500[A]	3	27,34,34	1.13	2 (7%)	34,54,54	2.10	7 (20%)
5	GOL	C	804	-	5,5,5	0.74	0	5,5,5	0.49	0
5	GOL	A	804	-	5,5,5	0.62	0	5,5,5	0.46	0
2	GCP	D	500	3	27,34,34	1.02	2 (7%)	34,54,54	2.39	8 (23%)
5	GOL	D	804	-	5,5,5	0.59	0	5,5,5	0.24	0
2	GCP	C	500[B]	3	27,34,34	0.91	2 (7%)	34,54,54	2.30	6 (17%)
2	GCP	A	500[B]	3	27,34,34	1.05	3 (11%)	34,54,54	2.17	5 (14%)
5	GOL	B	804	-	5,5,5	0.69	0	5,5,5	0.43	0
5	GOL	C	805	-	5,5,5	0.44	0	5,5,5	0.31	0
2	GCP	C	500[A]	3	27,34,34	0.95	2 (7%)	34,54,54	2.19	7 (20%)
2	GCP	B	500	3	27,34,34	1.27	3 (11%)	34,54,54	2.26	8 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	806	-	-	0/4/4/4	-
2	GCP	A	500[A]	3	-	2/15/38/38	0/3/3/3
5	GOL	C	804	-	-	4/4/4/4	-
5	GOL	A	804	-	-	2/4/4/4	-
2	GCP	D	500	3	-	1/15/38/38	0/3/3/3
5	GOL	D	804	-	-	0/4/4/4	-
2	GCP	C	500[B]	3	-	1/15/38/38	0/3/3/3
2	GCP	A	500[B]	3	-	3/15/38/38	0/3/3/3
5	GOL	B	804	-	-	0/4/4/4	-
5	GOL	C	805	-	-	1/4/4/4	-

*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GCP	C	500[A]	3	-	3/15/38/38	0/3/3/3
2	GCP	B	500	3	-	2/15/38/38	0/3/3/3

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	500[A]	GCP	PB-O3A	3.20	1.61	1.58
2	D	500	GCP	C6-N1	3.07	1.38	1.33
2	A	500[A]	GCP	C6-N1	2.89	1.38	1.33
2	A	500[B]	GCP	C6-N1	2.89	1.38	1.33
2	C	500[B]	GCP	PB-O3A	-2.82	1.55	1.58
2	B	500	GCP	C6-N1	2.71	1.37	1.33
2	B	500	GCP	PB-O1B	2.63	1.57	1.51
2	C	500[A]	GCP	C6-N1	2.34	1.37	1.33
2	C	500[B]	GCP	C6-N1	2.34	1.37	1.33
2	A	500[A]	GCP	C5-C6	2.31	1.45	1.41
2	A	500[B]	GCP	C5-C6	2.31	1.45	1.41
2	B	500	GCP	PB-O3A	-2.27	1.55	1.58
2	A	500[B]	GCP	PB-O3A	2.17	1.60	1.58
2	D	500	GCP	PB-O3A	2.16	1.60	1.58

All (41) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	500	GCP	C5-C6-N1	-8.49	111.82	123.43
2	C	500[A]	GCP	C5-C6-N1	-8.35	112.02	123.43
2	C	500[B]	GCP	C5-C6-N1	-8.35	112.02	123.43
2	A	500[A]	GCP	C5-C6-N1	-8.15	112.28	123.43
2	A	500[B]	GCP	C5-C6-N1	-8.15	112.28	123.43
2	B	500	GCP	C5-C6-N1	-8.08	112.38	123.43
2	C	500[A]	GCP	C2-N1-C6	6.21	125.80	115.93
2	C	500[B]	GCP	C2-N1-C6	6.21	125.80	115.93
2	D	500	GCP	C2-N1-C6	6.09	125.60	115.93
2	B	500	GCP	C2-N1-C6	5.85	125.22	115.93
2	A	500[A]	GCP	C2-N1-C6	5.82	125.17	115.93
2	A	500[B]	GCP	C2-N1-C6	5.82	125.17	115.93
2	C	500[B]	GCP	O1B-PB-C3B	5.63	123.96	109.07
2	D	500	GCP	O1G-PG-C3B	-5.36	99.69	111.24
2	A	500[B]	GCP	O1B-PB-C3B	4.92	122.07	109.07
2	B	500	GCP	O1G-PG-C3B	-4.32	101.94	111.24
2	D	500	GCP	O3G-PG-C3B	3.77	115.55	106.40
2	D	500	GCP	O1B-PB-C3B	3.38	118.02	109.07

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	500	GCP	O2B-PB-O1B	-3.31	99.02	110.07
2	C	500[A]	GCP	PB-O3A-PA	3.27	142.93	132.56
2	A	500[A]	GCP	C4-C5-C6	-3.12	117.82	120.80
2	A	500[B]	GCP	C4-C5-C6	-3.12	117.82	120.80
2	A	500[A]	GCP	N3-C2-N1	-2.99	123.24	127.22
2	A	500[B]	GCP	N3-C2-N1	-2.99	123.24	127.22
2	C	500[A]	GCP	C2-N3-C4	-2.97	111.96	115.36
2	C	500[B]	GCP	C2-N3-C4	-2.97	111.96	115.36
2	B	500	GCP	N3-C2-N1	-2.84	123.43	127.22
2	C	500[A]	GCP	N3-C2-N1	-2.80	123.49	127.22
2	C	500[B]	GCP	N3-C2-N1	-2.80	123.49	127.22
2	D	500	GCP	N3-C2-N1	-2.78	123.52	127.22
2	B	500	GCP	C4-C5-C6	-2.68	118.24	120.80
2	D	500	GCP	C4-C5-C6	-2.63	118.29	120.80
2	D	500	GCP	C2-N3-C4	-2.62	112.36	115.36
2	B	500	GCP	PB-O3A-PA	2.53	140.58	132.56
2	B	500	GCP	C2-N3-C4	-2.51	112.49	115.36
2	A	500[A]	GCP	O1G-PG-C3B	-2.42	106.02	111.24
2	C	500[A]	GCP	C4-C5-C6	-2.29	118.61	120.80
2	C	500[B]	GCP	C4-C5-C6	-2.29	118.61	120.80
2	C	500[A]	GCP	O1B-PB-C3B	2.14	114.74	109.07
2	A	500[A]	GCP	O2G-PG-O1G	2.06	117.84	112.39
2	A	500[A]	GCP	PB-O3A-PA	2.05	139.06	132.56

There are no chirality outliers.

All (19) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	500[B]	GCP	PG-C3B-PB-O1B
5	C	804	GOL	O1-C1-C2-C3
5	C	804	GOL	C1-C2-C3-O3
5	C	804	GOL	O2-C2-C3-O3
5	A	804	GOL	O1-C1-C2-C3
5	A	804	GOL	O1-C1-C2-O2
5	C	804	GOL	O1-C1-C2-O2
2	A	500[A]	GCP	C5'-O5'-PA-O3A
2	A	500[B]	GCP	PG-C3B-PB-O2B
2	A	500[B]	GCP	PB-O3A-PA-O2A
2	C	500[B]	GCP	PB-O3A-PA-O2A
2	B	500	GCP	PG-C3B-PB-O1B
2	C	500[A]	GCP	PG-C3B-PB-O1B
5	C	805	GOL	C1-C2-C3-O3

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	A	500[A]	GCP	PB-C3B-PG-O1G
2	B	500	GCP	PB-C3B-PG-O1G
2	C	500[A]	GCP	PB-C3B-PG-O1G
2	D	500	GCP	PB-C3B-PG-O1G
2	C	500[A]	GCP	C5'-O5'-PA-O3A


There are no ring outliers.


2 monomers are involved in 4 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	500[B]	GCP	2	0
2	C	500[A]	GCP	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	295/305 (96%)	-0.09	2 (0%) 87 91	28, 43, 66, 94	0
1	B	298/305 (97%)	-0.07	8 (2%) 54 63	27, 42, 71, 112	0
1	C	296/305 (97%)	-0.11	7 (2%) 59 67	26, 43, 80, 115	0
1	D	296/305 (97%)	0.16	19 (6%) 19 26	36, 53, 93, 122	0
All	All	1185/1220 (97%)	-0.03	36 (3%) 50 59	26, 45, 78, 122	0

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	179	TYR	7.7
1	D	212	TYR	7.2
1	D	473	TYR	5.7
1	B	179	TYR	5.0
1	C	212	TYR	4.7
1	D	464	PHE	4.6
1	C	182	ASN	3.8
1	B	182	ASN	3.7
1	D	182	ASN	3.7
1	B	181	ASP	3.5
1	A	212	TYR	3.5
1	C	479	ASP	3.4
1	D	455	TYR	3.3
1	C	185	ASN	3.0
1	B	212	TYR	2.9
1	C	184	THR	2.9
1	D	183	ALA	2.8
1	D	479	ASP	2.7
1	C	183	ALA	2.7
1	D	477	TYR	2.7
1	D	180	ASP	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	D	181	ASP	2.7
1	B	208	ILE	2.6
1	D	470	LYS	2.5
1	C	199	ASN	2.5
1	A	231	THR	2.5
1	D	463	GLU	2.5
1	D	210	SER	2.4
1	B	231	THR	2.4
1	D	211	GLY	2.3
1	D	254	ASN	2.3
1	D	460	ILE	2.2
1	B	479	ASP	2.2
1	D	474	LYS	2.1
1	B	180	ASP	2.1
1	D	462	GLN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	GOL	A	806	6/6	0.52	0.38	78,83,106,109	0
5	GOL	B	804	6/6	0.77	0.16	67,75,78,79	0
5	GOL	C	804	6/6	0.78	0.23	65,85,89,90	0
5	GOL	A	804	6/6	0.81	0.21	57,71,78,81	0
5	GOL	C	805	6/6	0.82	0.17	90,96,100,101	0
5	GOL	D	804	6/6	0.83	0.15	69,70,73,74	0
3	MG	C	800	1/1	0.87	0.11	74,74,74,74	0
3	MG	A	702	1/1	0.94	0.09	46,46,46,46	0

*Continued on next page...*




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	CL	B	802	1/1	0.95	0.22	72,72,72,72	0
4	CL	C	802	1/1	0.95	0.05	59,59,59,59	0
4	CL	A	802	1/1	0.96	0.13	70,70,70,70	0
3	MG	B	700	1/1	0.98	0.10	26,26,26,26	0
3	MG	B	702	1/1	0.98	0.06	40,40,40,40	0
2	GCP	C	500[A]	32/32	0.98	0.10	33,38,43,45	9
2	GCP	C	500[B]	32/32	0.98	0.10	33,37,41,43	9
3	MG	A	700	1/1	0.98	0.12	30,30,30,30	0
2	GCP	B	500	32/32	0.98	0.10	31,37,44,46	0
3	MG	C	700	1/1	0.99	0.07	35,35,35,35	0
3	MG	C	702	1/1	0.99	0.05	42,42,42,42	0
2	GCP	A	500[A]	32/32	0.99	0.11	32,36,40,43	9
3	MG	D	700	1/1	0.99	0.05	36,36,36,36	0
3	MG	D	702	1/1	0.99	0.03	41,41,41,41	0
2	GCP	D	500	32/32	0.99	0.09	36,44,49,50	0
2	GCP	A	500[B]	32/32	0.99	0.11	26,35,39,43	9

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




#### Electron density around GCP C 500 (A):

2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray  
 mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative)  
 and green (positive)







**Electron density around GCP C 500 (B):**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around GCP B 500:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.5 Other polymers [i](#)

There are no such residues in this entry.