

wwPDB NMR Structure Validation Summary Report (i)

Feb 8, 2022 – 06:28 PM EST

PDB ID : 1BZU
Title : STABILIZATION OF THE ANTICODON STEM-LOOP OF TRNALYS, 3
BY AN A+C BASE PAIR AND BY PSEUDOURIDINE, NMR, 1 STRUC-TURE
Authors : Durant, P.C.; Davis, D.R.
Deposited on : 1998-11-05

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/NMRValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	2.26
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.26

RNA backbone

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $SOLUTION\ NMR$

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.

Metric	Percentile	Ranks	Value				
RNA backbone			0.46				
Worse							
Percent	ile relative to all structures						
Percent	ile relative to all NMR structures						
Metric	Whole archive	NMR archive					
wietric	$(\# {\rm Entries})$	$(\# {\rm Entries})$					

4643

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

676

Mol	Chain	Length	Quality of chain
1	А	17	100%



2 Ensemble composition and analysis (i)

This entry contains 1 models. Identification of well-defined residues and clustering analysis are not possible.



3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 535 atoms, of which 182 are hydrogens and 0 are deuteriums.

• Molecule 1 is a RNA chain called TRNA.

Mol	Chain	Residues		Atoms					Trace
1	Λ	17	Total	С	Η	Ν	0	Р	0
1	A	A 17	535	160	182	58	119	16	0



4 Residue-property plots (i)

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: TRNA

Chain A: 100%



5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *molecular dynamics*.

Of the 32 calculated structures, 1 were deposited, based on the following criterion: *LEAST RE-STRAINT VIOLATION*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
Discover	refinement	95.0
VNMR	structure solution	
Felix	structure solution	
Discover	structure solution	

No chemical shift data was provided.



6 Model quality (i)

6.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: PSU

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	А	0	0	0	0
All	All	0	0	0	-

The all-atom clash score is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clash score for this structure is -.

There are no clashes.

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

There are no protein molecules in this entry.

6.3.2 Protein sidechains (i)

There are no protein molecules in this entry.

6.3.3 RNA (i)



Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers	Suiteness
1	А	16/17~(94%)	3~(19%)	2(12%)	0.46
All	All	16/17~(94%)	3~(19%)	2(12%)	0.46

The overall RNA backbone suiteness is 0.46.

All RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	А	34	U
1	А	35	U
1	А	36	U

All RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	А	34	U
1	А	35	U

6.4 Non-standard residues in protein, DNA, RNA chains (i)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with |Z| > 2 is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mal	Type	Chain	Res Link			nd lengt	
	туре	Chain	nes		Counts	RMSZ	#Z>2
1	PSU	А	39	1	17,21,22	1.32	1 (5%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with |Z| > 2 is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.



Mol	Type	Chain	Dec	Tink	В	ond ang	les
10101			nes		Counts	RMSZ	#Z>2
1	PSU	А	39	1	20,30,33	5.17	6 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PSU	А	39	1	-	0,7,25,26	0,2,2,2

All bond outliers are listed below.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	А	39	PSU	C4-N3	3.55	1.39	1.33

5 of 6 angle outliers are listed below. They are sorted according to the Z-score.

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	А	39	PSU	N3-C2-N1	15.87	115.81	128.43
1	А	39	PSU	C2-N3-C4	13.02	126.13	115.14
1	А	39	PSU	C5-C4-N3	8.39	114.55	125.36
1	А	39	PSU	C6-N1-C2	3.66	121.39	115.36
1	А	39	PSU	C5-C1'-C2'	3.34	109.36	115.32

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.



6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



7 Chemical shift validation (i)

No chemical shift data were provided

