



# wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 11, 2021 – 02:13 AM EDT

PDB ID : 3C0B  
Title : Crystal structure of the conserved archaeal protein Q6M145. Northeast Structural Genomics Consortium target MrR63  
Authors : Kuzin, A.P.; Su, M.; Seetharaman, J.; Wang, D.; Fang, Y.; Cunningham, K.; Ma, L.-C.; Xiao, R.; Liu, J.; Baran, M.C.; Acton, T.B.; Rost, B.; Montelione, G.T.; Hunt, J.F.; Tong, L.; Northeast Structural Genomics Consortium (NESG)  
Deposited on : 2008-01-19  
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.23.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	334	 9% 61% 30% • 5%
1	B	334	 5% 63% 28% • 5%
1	C	334	 4% 64% 27% • 5%
1	D	334	 6% 70% 21% • 6%

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 9912 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Conserved archaeal protein Q6M145.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	316	2438	1537	397	496	4	4	0	0	0
1	B	318	2455	1544	402	501	4	4	0	0	0
1	C	317	2444	1540	397	498	4	5	0	0	0
1	D	313	2413	1522	391	491	4	5	0	0	0

There are 84 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	48	ASN	LYS	engineered mutation	UNP Q6M145
A	50	VAL	ILE	engineered mutation	UNP Q6M145
A	93	ASN	ASP	engineered mutation	UNP Q6M145
A	101	SER	GLY	engineered mutation	UNP Q6M145
A	106	ASN	TYR	engineered mutation	UNP Q6M145
A	146	GLU	ASP	engineered mutation	UNP Q6M145
A	182	SER	LEU	engineered mutation	UNP Q6M145
A	187	ASP	ASN	engineered mutation	UNP Q6M145
A	254	ASN	THR	engineered mutation	UNP Q6M145
A	268	GLU	LYS	engineered mutation	UNP Q6M145
A	269	ASN	LYS	engineered mutation	UNP Q6M145
A	284	LEU	ILE	engineered mutation	UNP Q6M145
A	294	ALA	GLY	engineered mutation	UNP Q6M145
A	327	LEU	-	expression tag	UNP Q6M145
A	328	GLU	-	expression tag	UNP Q6M145
A	329	HIS	-	expression tag	UNP Q6M145
A	330	HIS	-	expression tag	UNP Q6M145
A	331	HIS	-	expression tag	UNP Q6M145
A	332	HIS	-	expression tag	UNP Q6M145
A	333	HIS	-	expression tag	UNP Q6M145
A	334	HIS	-	expression tag	UNP Q6M145

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	48	ASN	LYS	engineered mutation	UNP Q6M145
B	50	VAL	ILE	engineered mutation	UNP Q6M145
B	93	ASN	ASP	engineered mutation	UNP Q6M145
B	101	SER	GLY	engineered mutation	UNP Q6M145
B	106	ASN	TYR	engineered mutation	UNP Q6M145
B	146	GLU	ASP	engineered mutation	UNP Q6M145
B	182	SER	LEU	engineered mutation	UNP Q6M145
B	187	ASP	ASN	engineered mutation	UNP Q6M145
B	254	ASN	THR	engineered mutation	UNP Q6M145
B	268	GLU	LYS	engineered mutation	UNP Q6M145
B	269	ASN	LYS	engineered mutation	UNP Q6M145
B	284	LEU	ILE	engineered mutation	UNP Q6M145
B	294	ALA	GLY	engineered mutation	UNP Q6M145
B	327	LEU	-	expression tag	UNP Q6M145
B	328	GLU	-	expression tag	UNP Q6M145
B	329	HIS	-	expression tag	UNP Q6M145
B	330	HIS	-	expression tag	UNP Q6M145
B	331	HIS	-	expression tag	UNP Q6M145
B	332	HIS	-	expression tag	UNP Q6M145
B	333	HIS	-	expression tag	UNP Q6M145
B	334	HIS	-	expression tag	UNP Q6M145
C	48	ASN	LYS	engineered mutation	UNP Q6M145
C	50	VAL	ILE	engineered mutation	UNP Q6M145
C	93	ASN	ASP	engineered mutation	UNP Q6M145
C	101	SER	GLY	engineered mutation	UNP Q6M145
C	106	ASN	TYR	engineered mutation	UNP Q6M145
C	146	GLU	ASP	engineered mutation	UNP Q6M145
C	182	SER	LEU	engineered mutation	UNP Q6M145
C	187	ASP	ASN	engineered mutation	UNP Q6M145
C	254	ASN	THR	engineered mutation	UNP Q6M145
C	268	GLU	LYS	engineered mutation	UNP Q6M145
C	269	ASN	LYS	engineered mutation	UNP Q6M145
C	284	LEU	ILE	engineered mutation	UNP Q6M145
C	294	ALA	GLY	engineered mutation	UNP Q6M145
C	327	LEU	-	expression tag	UNP Q6M145
C	328	GLU	-	expression tag	UNP Q6M145
C	329	HIS	-	expression tag	UNP Q6M145
C	330	HIS	-	expression tag	UNP Q6M145
C	331	HIS	-	expression tag	UNP Q6M145
C	332	HIS	-	expression tag	UNP Q6M145
C	333	HIS	-	expression tag	UNP Q6M145
C	334	HIS	-	expression tag	UNP Q6M145

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	48	ASN	LYS	engineered mutation	UNP Q6M145
D	50	VAL	ILE	engineered mutation	UNP Q6M145
D	93	ASN	ASP	engineered mutation	UNP Q6M145
D	101	SER	GLY	engineered mutation	UNP Q6M145
D	106	ASN	TYR	engineered mutation	UNP Q6M145
D	146	GLU	ASP	engineered mutation	UNP Q6M145
D	182	SER	LEU	engineered mutation	UNP Q6M145
D	187	ASP	ASN	engineered mutation	UNP Q6M145
D	254	ASN	THR	engineered mutation	UNP Q6M145
D	268	GLU	LYS	engineered mutation	UNP Q6M145
D	269	ASN	LYS	engineered mutation	UNP Q6M145
D	284	LEU	ILE	engineered mutation	UNP Q6M145
D	294	ALA	GLY	engineered mutation	UNP Q6M145
D	327	LEU	-	expression tag	UNP Q6M145
D	328	GLU	-	expression tag	UNP Q6M145
D	329	HIS	-	expression tag	UNP Q6M145
D	330	HIS	-	expression tag	UNP Q6M145
D	331	HIS	-	expression tag	UNP Q6M145
D	332	HIS	-	expression tag	UNP Q6M145
D	333	HIS	-	expression tag	UNP Q6M145
D	334	HIS	-	expression tag	UNP Q6M145

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Ca 1 1	0	0
2	D	1	Total Ca 1 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	28	Total O 28 28	0	0
3	B	32	Total O 32 32	0	0
3	C	42	Total O 42 42	0	0
3	D	58	Total O 58 58	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Conserved archaeal protein Q6M145




- Molecule 1: Conserved archaeal protein Q6M145




- Molecule 1: Conserved archaeal protein Q6M145





● Molecule 1: Conserved archaeal protein Q6M145



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	157.40Å 87.27Å 123.99Å 90.00° 110.29° 90.00°	Depositor
Resolution (Å)	19.81 – 2.40 40.91 – 2.39	Depositor EDS
% Data completeness (in resolution range)	90.6 (19.81-2.40) 97.7 (40.91-2.39)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.14 (at 2.39Å)	Xtrriage
Refinement program	CNS 1.2	Depositor
R, $R_{free}$	0.214 , 0.259 0.227 , 0.273	Depositor DCC
$R_{free}$ test set	5979 reflections (4.92%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	31.2	Xtrriage
Anisotropy	0.294	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 53.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	9912	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	44.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.01% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.33	0/2469	0.57	0/3339
1	B	0.37	0/2486	0.60	0/3363
1	C	0.39	0/2474	0.64	0/3345
1	D	0.38	0/2442	0.64	0/3301
All	All	0.37	0/9871	0.61	0/13348

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2438	0	2418	108	0
1	B	2455	0	2430	119	0
1	C	2444	0	2427	117	0
1	D	2413	0	2402	61	0
2	B	1	0	0	0	0
2	D	1	0	0	0	0
3	A	28	0	0	3	0
3	B	32	0	0	2	0
3	C	42	0	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	58	0	0	3	0
All	All	9912	0	9677	396	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

The worst 5 of 396 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:89:VAL:HG11	1:B:103:LYS:HD3	1.25	1.08
1:C:108:LYS:HE3	1:C:108:LYS:HA	1.39	1.03
1:C:97:ILE:HD11	1:C:103:LYS:HG3	1.42	1.00
1:B:76:LEU:HD22	1:B:103:LYS:HZ2	1.28	0.98
1:A:72:VAL:HG21	1:A:109:VAL:HG21	1.46	0.94

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	308/334 (92%)	279 (91%)	29 (9%)	0	100	100
1	B	312/334 (93%)	286 (92%)	19 (6%)	7 (2%)	6	7
1	C	311/334 (93%)	292 (94%)	15 (5%)	4 (1%)	12	17
1	D	307/334 (92%)	291 (95%)	13 (4%)	3 (1%)	15	23
All	All	1238/1336 (93%)	1148 (93%)	76 (6%)	14 (1%)	14	20

5 of 14 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	101	SER

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	102	ALA
1	B	106	ASN
1	C	101	SER
1	C	102	ALA

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	275/287 (96%)	258 (94%)	17 (6%)	18	29
1	B	277/287 (96%)	261 (94%)	16 (6%)	20	32
1	C	276/287 (96%)	260 (94%)	16 (6%)	20	32
1	D	273/287 (95%)	259 (95%)	14 (5%)	24	39
All	All	1101/1148 (96%)	1038 (94%)	63 (6%)	20	33

5 of 63 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	179	ASN
1	D	105	ASN
1	C	32	MSE
1	D	72	VAL
1	D	213	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 38 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	113	ASN
1	D	269	ASN
1	D	124	ASN
1	D	176	HIS
1	D	324	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	312/334 (93%)	0.42	31 (9%) 7 6	21, 46, 105, 128	0
1	B	314/334 (94%)	0.07	18 (5%) 23 22	16, 38, 81, 121	0
1	C	312/334 (93%)	0.04	14 (4%) 33 31	13, 31, 82, 125	0
1	D	308/334 (92%)	0.06	19 (6%) 20 19	12, 31, 85, 116	0
All	All	1246/1336 (93%)	0.15	82 (6%) 18 17	12, 37, 96, 128	0

The worst 5 of 82 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	58	THR	9.2
1	A	9	GLY	7.3
1	A	43	LEU	5.7
1	B	101	SER	5.7
1	A	58	THR	5.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	CA	D	401	1/1	0.84	0.17	57,57,57,57	0
2	CA	B	402	1/1	0.87	0.21	66,66,66,66	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.