



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 6, 2023 – 08:44 am GMT

PDB ID : 2C18
Title : 5-(4-Carboxy-2-oxo-butane-1-sulfonyl)-4-oxo-pentanoic acid bound to Porphobilinogen synthase from *Pseudomonas aeruginosa*
Authors : Frere, F.; Nentwich, M.; Gacond, S.; Heinz, D.W.; Neier, R.; Frankenberg-Dinkel, N.
Deposited on : 2005-09-11
Resolution : 1.93 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.93 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4310 (1.96-1.92)
Clashscore	141614	1023 (1.94-1.94)
Ramachandran outliers	138981	1007 (1.94-1.94)
Sidechain outliers	138945	1007 (1.94-1.94)
RSRZ outliers	127900	4250 (1.96-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	337	 2% 91% 7% ...
1	B	337	 2% 90% 7% ..

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 6180 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DELTA-AMINOLEVULINIC ACID DEHYDRATASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	334	2753	1725	484	530	14	0	19	1
1	B	329	2757	1730	488	527	12	0	23	1

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	199	VAL	ILE	conflict	UNP Q59643
B	199	VAL	ILE	conflict	UNP Q59643

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	5	Total	Mg	0	0
			5	5		
2	B	2	Total	Mg	0	0
			2	2		

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Na	0	0
			1	1		
3	B	1	Total	Na	0	0
			1	1		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Cl	0	0
			2	2		

- Molecule 5 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: C₆H₁₄O₄).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf		
5	A	1	Total	C	O	0	0
			10	6	4		
5	B	1	Total	C	O	0	0
			10	6	4		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	
6	A	314	Total	O	0	0
			314	314		
6	B	325	Total	O	0	0
			325	325		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DELTA-AMINOLEVULINIC ACID DEHYDRATASE



- Molecule 1: DELTA-AMINOLEVULINIC ACID DEHYDRATASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 4 21 2	Depositor
Cell constants a, b, c, α , β , γ	125.60Å 125.60Å 86.57Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	89.80 – 1.93 39.51 – 1.89	Depositor EDS
% Data completeness (in resolution range)	94.3 (89.80-1.93) 93.0 (39.51-1.89)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.30 (at 1.89Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.149 , 0.186 0.149 , 0.185	Depositor DCC
R_{free} test set	2628 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	20.3	Xtrriage
Anisotropy	0.034	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 63.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6180	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.85% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, CL, PGE, MG, LSO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.73	0/2774	0.74	1/3763 (0.0%)
1	B	0.71	0/2777	0.70	1/3767 (0.0%)
All	All	0.72	0/5551	0.72	2/7530 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	B	0	2
All	All	0	5

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	51	ARG	NE-CZ-NH2	-5.65	117.47	120.30
1	A	333	LEU	N-CA-C	5.57	126.05	111.00

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	260	LSO	Mainchain,Peptide
1	A	333	LEU	Peptide
1	B	260	LSO	Mainchain,Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2753	0	2697	19	0
1	B	2757	0	2711	21	0
2	A	5	0	0	0	0
2	B	2	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	2	0	0	0	0
5	A	10	0	14	0	0
5	B	10	0	14	1	0
6	A	314	0	0	1	0
6	B	325	0	0	7	0
All	All	6180	0	5436	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:263[B]:MET:HG2	1:B:263[B]:MET:HG2	1.18	1.14
1:B:275[B]:GLU:OE2	6:B:2293:HOH:O	1.73	1.07
1:A:263[B]:MET:CG	1:B:263[B]:MET:HG2	1.97	0.94
1:A:330:ALA:O	1:A:334:ARG:HB2	1.69	0.93
1:A:263[B]:MET:HG2	1:B:263[B]:MET:CG	1.98	0.92
1:A:331:GLU:O	1:A:334:ARG:HB3	1.92	0.69
1:A:330:ALA:O	1:A:334:ARG:CB	2.44	0.66
1:A:263[B]:MET:SD	1:B:263[B]:MET:SD	2.94	0.65
5:B:1340:PGE:H2	6:B:2325:HOH:O	1.97	0.64
1:A:42[B]:VAL:HG11	1:A:68:LEU:HD13	1.81	0.62
1:A:42[B]:VAL:HG11	1:A:68:LEU:CD1	2.30	0.61
1:B:48[A]:VAL:O	1:B:50:GLN:HG2	2.03	0.59
1:B:314:LYS:HE3	6:B:2085:HOH:O	2.04	0.57
1:B:331[B]:GLU:O	1:B:335:ARG:HG2	2.05	0.56
1:B:58:PRO:HB2	1:B:297[B]:GLN:HE22	1.74	0.53
1:B:131:ASP:CG	6:B:2203:HOH:O	2.47	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:331[A]:GLU:O	1:B:335:ARG:HG2	2.11	0.51
1:A:105:GLU:O	1:A:110[A]:ARG:HD3	2.11	0.50
1:A:3:PHE:N	6:A:2001:HOH:O	2.45	0.50
1:B:93:LYS:HG2	6:B:2162:HOH:O	2.11	0.49
1:A:74[B]:GLU:HG2	1:A:327:LYS:HE3	1.97	0.46
1:B:216[A]:ASP:OD2	6:B:2257:HOH:O	2.19	0.44
1:A:301[A]:LEU:HG	1:A:305[A]:VAL:HG11	2.00	0.44
1:B:332:GLN:NE2	6:B:2323:HOH:O	2.50	0.44
1:B:334:ARG:HH11	1:B:335:ARG:HH21	1.66	0.44
1:B:51:ARG:NH2	1:B:71:GLU:OE2	2.42	0.43
1:B:130:LEU:HD12	1:B:140:GLY:HA2	2.00	0.43
1:A:296:ILE:HD11	1:A:305[B]:VAL:HG12	2.01	0.43
1:A:155:ASP:O	1:A:159[B]:ARG:HG3	2.19	0.42
1:A:72:ALA:HA	1:A:75:TRP:CE3	2.55	0.42
1:A:263[B]:MET:SD	1:B:263[B]:MET:HG2	2.59	0.42
1:A:4:THR:HA	1:A:5:PRO:HD3	1.91	0.41
1:B:48[B]:VAL:O	1:B:50:GLN:HG2	2.21	0.41
1:B:327:LYS:HE2	1:B:331[B]:GLU:OE2	2.20	0.41
1:B:72:ALA:HA	1:B:75:TRP:CE3	2.56	0.40
1:A:260:LSO:HEC1	1:A:260:LSO:H3C2	1.88	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	350/337 (104%)	343 (98%)	5 (1%)	2 (1%)	25 13
1	B	347/337 (103%)	336 (97%)	10 (3%)	1 (0%)	41 32
All	All	697/674 (103%)	679 (97%)	15 (2%)	3 (0%)	29 24

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	261	PRO
1	B	261	PRO
1	A	334	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	284/268 (106%)	276 (97%)	8 (3%)	43 29
1	B	285/268 (106%)	283 (99%)	2 (1%)	84 81
All	All	569/536 (106%)	559 (98%)	10 (2%)	60 47

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	THR
1	A	118	ARG
1	A	256	MET
1	A	277	ARG
1	A	283	TYR
1	A	301[A]	LEU
1	A	301[B]	LEU
1	A	334	ARG
1	B	256	MET
1	B	283	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	298	ASN
1	B	298	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	LSO	A	260	1	21,26,27	0.78	0	24,33,35	1.77	7 (29%)
1	LSO	B	260	1	21,26,27	0.77	0	24,33,35	1.57	5 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LSO	A	260	1	-	3/27/30/32	-
1	LSO	B	260	1	-	4/27/30/32	-

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	260	LSO	O5-S6-O6	-3.99	111.45	117.42
1	B	260	LSO	O5-S6-O6	-3.45	112.26	117.42
1	B	260	LSO	O6-S6-C5	3.28	111.85	107.84
1	A	260	LSO	O6-S6-C5	2.83	111.31	107.84
1	A	260	LSO	C5-S6-C7	2.80	110.84	105.83
1	A	260	LSO	C10-C9-C8	-2.69	110.20	113.61
1	B	260	LSO	C5-S6-C7	2.56	110.40	105.83
1	B	260	LSO	O4-C11-C10	2.31	121.44	114.03
1	A	260	LSO	O5-S6-C5	-2.19	105.16	107.84
1	A	260	LSO	CD-CE-NZ	-2.14	106.77	110.66
1	B	260	LSO	CE-NZ-C4	2.06	128.40	122.49
1	A	260	LSO	CG-CD-CE	-2.04	106.45	113.57

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	260	LSO	C4-C5-S6-O5
1	A	260	LSO	S6-C7-C8-O7
1	B	260	LSO	S6-C7-C8-O7
1	B	260	LSO	C9-C10-C11-O4
1	B	260	LSO	C9-C10-C11-O3
1	A	260	LSO	S6-C7-C8-C9
1	B	260	LSO	S6-C7-C8-C9

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	260	LSO	1	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 11 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	PGE	A	1341	-	9,9,9	0.52	0	8,8,8	0.38	0
5	PGE	B	1340	-	9,9,9	0.48	0	8,8,8	0.68	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '·' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	PGE	A	1341	-	-	5/7/7/7	-
5	PGE	B	1340	-	-	3/7/7/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	1341	PGE	C4-C3-O2-C2
5	A	1341	PGE	C6-C5-O3-C4
5	A	1341	PGE	O2-C3-C4-O3
5	B	1340	PGE	O3-C5-C6-O4
5	B	1340	PGE	O2-C3-C4-O3
5	B	1340	PGE	O1-C1-C2-O2
5	A	1341	PGE	O1-C1-C2-O2
5	A	1341	PGE	O3-C5-C6-O4

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	1340	PGE	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	333/337 (98%)	-0.29	7 (2%) 63 70	10, 16, 27, 47	4 (1%)
1	B	328/337 (97%)	-0.36	6 (1%) 68 74	11, 16, 29, 46	3 (0%)
All	All	661/674 (98%)	-0.33	13 (1%) 65 71	10, 16, 29, 47	7 (1%)

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	226	LYS	7.1
1	A	336	GLY	5.3
1	B	336	GLY	5.2
1	B	221	ALA	4.7
1	B	335	ARG	4.4
1	B	144	ASP	3.3
1	B	2	SER	3.3
1	A	334	ARG	3.0
1	A	48	VAL	2.9
1	A	225	GLY	2.7
1	A	49[A]	ASN	2.4
1	A	187[A]	GLU	2.3
1	B	334	ARG	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	LSO	B	260	27/28	0.95	0.10	11,18,34,37	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
1	LSO	A	260	27/28	0.96	0.10	12,18,26,27	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	MG	A	348	1/1	0.76	0.19	27,27,27,27	1
2	MG	A	347	1/1	0.81	0.13	53,53,53,53	0
5	PGE	B	1340	10/10	0.88	0.22	33,38,43,45	0
5	PGE	A	1341	10/10	0.91	0.18	33,38,44,45	0
2	MG	A	349	1/1	0.96	0.15	29,29,29,29	1
2	MG	B	1338	1/1	0.97	0.06	30,30,30,30	1
3	NA	B	1339	1/1	0.97	0.05	22,22,22,22	0
3	NA	A	1339	1/1	0.98	0.05	20,20,20,20	0
2	MG	A	1337	1/1	0.98	0.06	17,17,17,17	0
4	CL	A	1342	1/1	0.99	0.04	24,24,24,24	0
2	MG	B	1337	1/1	1.00	0.10	15,15,15,15	0
2	MG	A	1338	1/1	1.00	0.02	23,23,23,23	0
4	CL	A	1340	1/1	1.00	0.04	18,18,18,18	0

6.5 Other polymers [i](#)

There are no such residues in this entry.