



Full wwPDB X-ray Structure Validation Report

Dec 10, 2023 – 03:36 am GMT

PDB ID : 2C40
Title : CRYSTAL STRUCTURE OF INOSINE-URIDINE PREFERRING NUCLEOSIDE HYDROLASE FROM BACILLUS ANTHRACIS AT 2.2A RESOLUTION
Authors : Moroz, O.V.; Blagova, E.V.; Fogg, M.J.; Levdikov, V.M.; Brannigan, J.A.; Wilkinson, A.J.; Wilson, K.S.
Deposited on : 2005-10-13
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	312	<div style="display: flex; align-items: center;"> <div style="width: 3%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 75%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 17%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 5%; height: 10px; background-color: orange; margin-right: 5px;"></div> <div style="width: 0%; height: 10px; background-color: grey;"></div> </div> <p style="margin-left: 10px;">3% 75% 17% 5% .</p>
1	B	312	<div style="display: flex; align-items: center;"> <div style="width: 9%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 77%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 16%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 0%; height: 10px; background-color: orange;"></div> <div style="width: 0%; height: 10px; background-color: grey;"></div> </div> <p style="margin-left: 10px;">9% 77% 16% . .</p>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4867 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called INOSINE-URIDINE PREFERRING NUCLEOSIDE HYDROLASE FAMILY PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	302	Total 2380	C 1532	N 407	O 434	S 7	0	3	0
1	B	300	Total 2342	C 1507	N 396	O 432	S 7	0	1	0

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total 1	Ca 1	0	0
2	B	1	Total 1	Ca 1	0	0

- Molecule 3 is alpha-D-ribofuranose (three-letter code: RIB) (formula: C₅H₁₀O₅).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			9	5	4		
3	B	1	Total	C	O	0	0
			9	5	4		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	77	Total	O	0	0
			77	77		
4	B	48	Total	O	0	0
			48	48		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: INOSINE-URIDINE PREFERRED NUCLEOSIDE HYDROLASE FAMILY PROTEIN



- Molecule 1: INOSINE-URIDINE PREFERRED NUCLEOSIDE HYDROLASE FAMILY PROTEIN



4 Data and refinement statistics i

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	78.15Å 78.15Å 203.63Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	67.88 – 2.20 24.81 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.6 (67.88-2.20) 99.7 (24.81-2.20)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.37 (at 2.19Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.205 , 0.242 0.214 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	38.7	Xtrriage
Anisotropy	0.244	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 54.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	0.045 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4867	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: RIB, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.82	1/2440 (0.0%)	0.89	7/3331 (0.2%)
1	B	0.82	2/2402 (0.1%)	0.87	5/3280 (0.2%)
All	All	0.82	3/4842 (0.1%)	0.88	12/6611 (0.2%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	288	VAL	CB-CG1	-6.07	1.40	1.52
1	B	283	ASP	CG-OD1	5.64	1.38	1.25
1	B	283	ASP	CG-OD2	5.50	1.38	1.25

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	26	ASN	N-CA-CB	-7.10	97.83	110.60
1	A	286	ARG	NE-CZ-NH2	-7.07	116.76	120.30
1	A	276	GLN	CB-CA-C	-6.67	97.07	110.40
1	A	286	ARG	NE-CZ-NH1	6.22	123.41	120.30
1	B	293	ASP	CB-CG-OD1	-6.04	112.87	118.30
1	B	54	ARG	NE-CZ-NH2	-5.98	117.31	120.30
1	A	147	ARG	NE-CZ-NH1	5.66	123.13	120.30
1	A	15	LEU	CB-CG-CD1	5.42	120.21	111.00
1	A	45	MET	CG-SD-CE	-5.27	91.76	100.20
1	B	286	ARG	NE-CZ-NH2	-5.20	117.70	120.30
1	A	298	ARG	NE-CZ-NH1	5.08	122.84	120.30
1	B	13	ASP	CB-CG-OD2	-5.03	113.78	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2380	0	2312	53	0
1	B	2342	0	2264	37	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	9	0	0	0	0
3	B	9	0	0	0	0
4	A	77	0	0	5	0
4	B	48	0	0	4	0
All	All	4867	0	4576	89	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (89) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:113:LEU:HD23	1:B:120:THR:HG23	1.44	0.99
1:A:66:ASN:ND2	1:A:133:ARG:HH22	1.71	0.89
1:A:66:ASN:HD22	1:A:133:ARG:HH22	1.27	0.83
1:A:89:ILE:HD13	1:A:219:ILE:HD11	1.60	0.82
1:A:34:VAL:HB	1:A:45:MET:HE2	1.64	0.79
1:B:24:MET:CE	1:B:27:VAL:HG21	2.13	0.79
1:A:300:PHE:O	1:A:304:THR:HG23	1.83	0.78
1:B:113:LEU:HD23	1:B:120:THR:CG2	2.13	0.77
1:B:24:MET:HE2	1:B:27:VAL:HG21	1.65	0.77
1:B:66:ASN:ND2	1:B:133:ARG:HH22	1.83	0.77
1:B:300:PHE:O	1:B:304:THR:HG23	1.85	0.76
1:A:24:MET:HE3	1:A:27:VAL:HG21	1.68	0.76
1:A:45:MET:CE	1:A:64:ALA:HA	2.17	0.75
1:B:53:ASP:OD1	1:B:98:THR:HG23	1.88	0.73
1:A:181:VAL:HG12	1:A:269:VAL:HG21	1.72	0.72
1:A:24:MET:CE	1:A:27:VAL:HG21	2.20	0.72
1:A:53:ASP:OD1	1:A:98:THR:HG23	1.89	0.71
1:A:24:MET:HE3	1:A:27:VAL:CG2	2.20	0.71
1:A:185:TRP:O	1:A:286:ARG:HD2	1.92	0.70

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:113:LEU:CD2	1:B:120:THR:HG23	2.22	0.69
1:A:34:VAL:HB	1:A:45:MET:CE	2.23	0.67
1:A:45:MET:HE3	1:A:64:ALA:HA	1.77	0.66
1:B:185:TRP:O	1:B:286:ARG:HD2	1.96	0.65
1:A:262:VAL:HG11	1:A:289:HIS:HB3	1.76	0.65
1:A:53:ASP:OD1	1:A:98:THR:CG2	2.45	0.64
1:A:262:VAL:CG1	1:A:289:HIS:HB3	2.29	0.63
1:B:112:THR:O	1:B:116:THR:HG23	1.99	0.62
1:B:123:LEU:HG	1:B:254:VAL:HG21	1.82	0.61
1:A:45:MET:HE1	1:A:64:ALA:HA	1.82	0.61
1:A:300:PHE:O	1:A:304:THR:CG2	2.49	0.61
1:B:231:VAL:HG22	1:B:231:VAL:O	2.01	0.60
1:B:166:HIS:HE1	1:B:170:ALA:H	1.48	0.60
1:B:205:THR:HG22	1:B:208:ILE:HD12	1.83	0.60
1:A:113:LEU:HD22	1:A:120:THR:HG23	1.83	0.59
1:B:106:HIS:HD2	4:B:2023:HOH:O	1.86	0.59
1:A:115:GLN:CB	4:A:2029:HOH:O	2.50	0.59
1:A:45:MET:HE3	1:A:64:ALA:CB	2.34	0.58
1:A:231:VAL:O	1:A:231:VAL:HG13	2.04	0.56
1:A:298:ARG:HD2	4:A:2066:HOH:O	2.04	0.56
1:A:45:MET:HE3	1:A:64:ALA:CA	2.35	0.56
1:B:53:ASP:OD1	1:B:98:THR:CG2	2.53	0.55
1:B:83:TYR:HB3	1:B:230:ILE:HD13	1.88	0.55
1:A:120:THR:HG22	4:A:2030:HOH:O	2.06	0.53
1:B:66:ASN:HD21	1:B:133:ARG:HH22	1.54	0.53
1:A:166:HIS:HE1	1:A:170:ALA:H	1.57	0.52
1:B:66:ASN:HD22	1:B:66:ASN:C	2.13	0.52
1:B:66:ASN:ND2	1:B:133:ARG:NH2	2.56	0.51
1:A:112:THR:O	1:A:116:THR:HG23	2.10	0.51
1:A:120:THR:CG2	4:A:2030:HOH:O	2.58	0.51
1:B:185:TRP:HZ3	1:B:267:SER:HG	1.59	0.50
1:A:123:LEU:HG	1:A:254:VAL:HG21	1.95	0.49
1:B:83:TYR:CB	1:B:230:ILE:HD13	2.42	0.49
1:A:113:LEU:CD2	1:A:120:THR:HG23	2.43	0.49
1:B:6:PHE:CD2	1:B:123:LEU:HD13	2.48	0.48
1:B:13:ASP:HB3	1:B:247:ASP:HB2	1.96	0.48
1:B:199:THR:HG21	1:B:246:TRP:NE1	2.29	0.47
1:A:21:LEU:HD23	1:A:24:MET:CE	2.45	0.46
1:B:296:HIS:HD2	1:B:297:ASP:OD1	1.99	0.46
1:A:76:ASP:CB	4:A:2021:HOH:O	2.63	0.46
1:B:116:THR:HG21	4:B:2006:HOH:O	2.15	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:155:PHE:N	1:B:197:GLU:OE2	2.48	0.46
1:B:202:VAL:HG23	1:B:202:VAL:O	2.16	0.45
1:A:24:MET:HE3	1:A:27:VAL:CB	2.47	0.45
1:A:181:VAL:CG1	1:A:269:VAL:HG21	2.45	0.44
1:B:120:THR:HG21	4:B:2006:HOH:O	2.17	0.44
1:A:45:MET:HE3	1:A:64:ALA:HB2	1.99	0.44
1:A:45:MET:CE	1:A:64:ALA:CA	2.93	0.44
1:A:199:THR:HG21	1:A:244:TYR:HB3	1.99	0.44
1:B:166:HIS:CE1	1:B:170:ALA:H	2.32	0.44
1:B:166:HIS:CE1	1:B:170:ALA:HB3	2.53	0.43
1:B:24:MET:CE	1:B:27:VAL:CG2	2.91	0.43
1:B:209:ARG:NH1	4:B:2037:HOH:O	2.50	0.43
1:A:6:PHE:CD2	1:A:123:LEU:HD13	2.53	0.43
1:A:193:LEU:HD13	1:A:195:THR:HG22	2.01	0.43
1:A:21:LEU:HA	1:A:24:MET:HE2	2.00	0.43
1:A:54[A]:ARG:NH1	1:A:89:ILE:HD12	2.34	0.43
1:A:129:THR:HB	1:A:177:ASP:OD2	2.19	0.43
1:A:24:MET:HE1	1:A:27:VAL:HG21	1.97	0.43
1:A:83:TYR:HB2	1:A:230:ILE:HD13	2.00	0.42
1:A:199:THR:HG23	1:A:244:TYR:CD2	2.55	0.42
1:A:296:HIS:HD2	1:A:297:ASP:OD1	2.03	0.42
1:A:74:PRO:O	1:A:78:ARG:HG3	2.19	0.41
1:A:225:GLY:HA3	1:B:88:PRO:HG3	2.02	0.41
1:B:282:THR:HG22	1:B:284:ASP:H	1.85	0.41
1:A:54[A]:ARG:CZ	1:A:89:ILE:HD12	2.50	0.41
1:A:194:ILE:HD12	1:A:194:ILE:N	2.36	0.40
1:B:206:ILE:HD11	1:B:233:PRO:CB	2.52	0.40
1:A:231:VAL:O	1:A:231:VAL:CG1	2.68	0.40
1:A:21:LEU:HD23	1:A:24:MET:HE1	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	301/312 (96%)	293 (97%)	8 (3%)	0	100	100
1	B	297/312 (95%)	289 (97%)	8 (3%)	0	100	100
All	All	598/624 (96%)	582 (97%)	16 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	247/269 (92%)	220 (89%)	27 (11%)	6	5
1	B	243/269 (90%)	219 (90%)	24 (10%)	8	7
All	All	490/538 (91%)	439 (90%)	51 (10%)	7	6

All (51) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	2	LYS
1	A	15	LEU
1	A	29	LEU
1	A	59	THR
1	A	66	ASN
1	A	89	ILE
1	A	98	THR
1	A	120	THR
1	A	123	LEU
1	A	129	THR
1	A	141	ILE
1	A	147	ARG
1	A	148	LEU
1	A	151	MET
1	A	183	ARG
1	A	185	TRP
1	A	204	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	205	THR
1	A	231	VAL
1	A	249	LEU
1	A	258	ASP
1	A	261	LYS
1	A	269	VAL
1	A	271	THR
1	A	283	ASP
1	A	304	THR
1	B	1	MET
1	B	13	ASP
1	B	15	LEU
1	B	29	LEU
1	B	59	THR
1	B	66	ASN
1	B	98	THR
1	B	120	THR
1	B	123	LEU
1	B	129	THR
1	B	133	ARG
1	B	147	ARG
1	B	148	LEU
1	B	151	MET
1	B	157	THR
1	B	186	GLU
1	B	193	LEU
1	B	197	GLU
1	B	204	LEU
1	B	205	THR
1	B	249	LEU
1	B	271	THR
1	B	298	ARG
1	B	304	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	7	ASN
1	A	66	ASN
1	A	106	HIS
1	A	166	HIS
1	A	296	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	7	ASN
1	B	66	ASN
1	B	106	HIS
1	B	166	HIS
1	B	296	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	RIB	A	1312	2	9,9,10	2.00	3 (33%)	10,12,14	2.72	3 (30%)
3	RIB	B	1312	2	9,9,10	1.70	2 (22%)	10,12,14	2.50	2 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '·' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	RIB	A	1312	2	-	0/2/15/18	0/1/1/1
3	RIB	B	1312	2	-	0/2/15/18	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1312	RIB	C2-C3	3.77	1.59	1.53
3	B	1312	RIB	C2-C3	3.48	1.59	1.53
3	A	1312	RIB	O2-C2	3.47	1.50	1.43
3	B	1312	RIB	O2-C2	3.13	1.50	1.43
3	A	1312	RIB	C1-C2	2.17	1.55	1.51

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1312	RIB	O2-C2-C3	7.15	124.81	111.27
3	A	1312	RIB	O2-C2-C3	6.72	123.99	111.27
3	A	1312	RIB	O4-C4-C3	3.84	108.10	104.70
3	A	1312	RIB	O3-C3-C2	2.10	117.10	112.04
3	B	1312	RIB	O3-C3-C2	2.00	116.87	112.04

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	302/312 (96%)	0.13	10 (3%) 46 44	40, 43, 49, 54	0
1	B	300/312 (96%)	0.50	29 (9%) 7 6	39, 43, 49, 52	0
All	All	602/624 (96%)	0.31	39 (6%) 18 17	39, 43, 49, 54	0

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	128	LEU	5.1
1	B	272	TYR	4.7
1	B	124	PHE	4.6
1	B	283	ASP	4.0
1	B	123	LEU	3.9
1	B	205	THR	3.9
1	A	18	LEU	3.6
1	A	283	ASP	3.4
1	B	150	TRP	3.3
1	B	282	THR	3.3
1	A	193	LEU	3.3
1	B	136	TYR	3.3
1	B	232	PRO	3.2
1	A	194	ILE	3.0
1	B	149	VAL	2.9
1	B	201	GLN	2.9
1	B	202	VAL	2.8
1	A	243	TYR	2.8
1	A	217	LYS	2.8
1	B	285	GLY	2.7
1	B	204	LEU	2.7
1	B	233	PRO	2.6
1	B	18	LEU	2.6
1	B	281	GLU	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	203	PRO	2.6
1	B	94	GLY	2.5
1	A	150	TRP	2.4
1	B	127	PRO	2.4
1	A	1	MET	2.4
1	B	225	GLY	2.4
1	B	157	THR	2.4
1	A	15	LEU	2.3
1	B	185	TRP	2.3
1	B	284	ASP	2.3
1	B	1	MET	2.3
1	B	194	ILE	2.1
1	B	244	TYR	2.0
1	A	124	PHE	2.0
1	B	125	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	RIB	A	1312	9/10	0.71	0.27	42,48,51,52	0
3	RIB	B	1312	9/10	0.78	0.22	38,47,50,51	0
2	CA	B	1311	1/1	0.99	0.10	33,33,33,33	0
2	CA	A	1311	1/1	1.00	0.04	31,31,31,31	0

6.5 Other polymers [i](#)

There are no such residues in this entry.