



## Full wwPDB X-ray Structure Validation Report ⓘ

Dec 10, 2023 – 10:26 am GMT

PDB ID : 2C4L  
Title : Structure of Neuraminidase Subtype N9 Complexed with 30 MM Sialic Acid (NANA, NEU5AC), Crystal Soaked for 24 Hours at 291 K and Finally Back-soaked for 30 Min in a Cryoprotectant Solution which did not contain NEU5AC  
Authors : Rudino-Pinera, E.; Tunnah, P.; Crennell, S.J.; Webster, R.G.; Laver, W.G.; Garman, E.F.  
Deposited on : 2005-10-20  
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1479 (2.16-2.16)
Clashscore	141614	1585 (2.16-2.16)
Ramachandran outliers	138981	1560 (2.16-2.16)
Sidechain outliers	138945	1559 (2.16-2.16)
RSRZ outliers	127900	1456 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	388	 89% 10% .
2	B	2	 50% 50%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	NAG	A	1471	-	-	X	-
5	BMA	A	1480	-	-	X	-

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 3490 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NEURAMINIDASE SUBTYPE N9.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	388	3067	1914	538	592	23	0	0	0

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
2	B	2	22	12	10	0	0	0

- Molecule 3 is N-acetyl-alpha-neuraminic acid (three-letter code: SIA) (formula: C<sub>11</sub>H<sub>19</sub>NO<sub>9</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	21	11	1	9	0	0
3	A	1	21	11	1	9	0	0

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
4	A	1	14	8	1	5	0	0
4	A	1	14	8	1	5	0	0
4	A	1	14	8	1	5	0	0
4	A	1	14	8	1	5	0	0
4	A	1	14	8	1	5	0	0

- Molecule 5 is beta-D-mannopyranose (three-letter code: BMA) (formula:  $C_6H_{12}O_6$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 11 6 5	0	0
5	A	1	Total C O 11 6 5	0	0
5	A	1	Total C O 11 6 5	0	0
5	A	1	Total C O 11 6 5	0	0
5	A	1	Total C O 11 6 5	0	0

- Molecule 6 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total Ca 1 1	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	233	Total O 233 233	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: NEURAMINIDASE SUBTYPE N9



- Molecule 2: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose





## 4 Data and refinement statistics

Property	Value	Source
Space group	I 4 3 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	181.18Å 181.18Å 181.18Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.70 – 2.15 42.70 – 2.15	Depositor EDS
% Data completeness (in resolution range)	100.0 (42.70-2.15) 100.0 (42.70-2.15)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	6.59 (at 2.16Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.155 , 0.196 0.157 , 0.195	Depositor DCC
$R_{free}$ test set	1398 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	16.5	Xtrriage
Anisotropy	0.000	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 30.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3490	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	15.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.80% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, SIA, CA, BMA, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	1.05	6/3150 (0.2%)	0.87	8/4290 (0.2%)

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	416	GLU	CD-OE1	-27.43	0.95	1.25
1	A	333	VAL	C-N	19.11	1.67	1.33
1	A	392	GLN	C-N	15.62	1.61	1.33
1	A	463	LYS	CE-NZ	-7.21	1.31	1.49
1	A	416	GLU	CD-OE2	6.35	1.32	1.25
1	A	174	GLU	CB-CG	5.33	1.62	1.52

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	416	GLU	OE1-CD-OE2	21.47	149.07	123.30
1	A	416	GLU	CG-CD-OE2	-12.55	93.21	118.30
1	A	127	ASP	CB-CG-OD2	-9.20	110.02	118.30
1	A	364	ARG	NE-CZ-NH1	9.05	124.83	120.30
1	A	172	ARG	NE-CZ-NH2	-8.23	116.18	120.30
1	A	364	ARG	NE-CZ-NH2	-8.20	116.20	120.30
1	A	172	ARG	NE-CZ-NH1	6.08	123.34	120.30
1	A	125	ASP	CB-CG-OD1	5.94	123.64	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3067	0	2895	38	0
2	B	22	0	19	3	0
3	A	42	0	36	0	0
4	A	70	0	63	16	0
5	A	55	0	49	12	0
6	A	1	0	0	0	0
7	A	233	0	0	1	0
All	All	3490	0	3062	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:146:ASN:HD21	4:A:1475:NAG:C1	0.99	1.57
1:A:200:ASN:HD21	4:A:1471:NAG:C1	0.90	1.50
1:A:200:ASN:ND2	4:A:1471:NAG:C1	1.75	1.44
4:A:1472:NAG:O4	5:A:1476:BMA:C1	1.67	1.43
4:A:1471:NAG:O4	4:A:1472:NAG:C1	1.67	1.42
5:A:1479:BMA:C1	2:B:2:MAN:O2	1.73	1.35
1:A:146:ASN:ND2	4:A:1475:NAG:C1	1.85	1.34
4:A:1473:NAG:O4	4:A:1474:NAG:C1	1.84	1.23
5:A:1476:BMA:O6	5:A:1480:BMA:C1	2.01	1.09
1:A:172:ARG:HD3	1:A:209:ARG:NH2	1.69	1.08
1:A:87:LEU:H	1:A:233:HIS:HD2	1.13	0.93
4:A:1473:NAG:HO4	4:A:1474:NAG:C1	1.76	0.91
1:A:168:THR:H	1:A:170:ASN:HD21	1.17	0.90
5:A:1480:BMA:O3	5:A:1482:BMA:C1	2.23	0.86
1:A:172:ARG:HD3	1:A:209:ARG:HH22	1.43	0.81
1:A:87:LEU:H	1:A:233:HIS:CD2	1.99	0.79
4:A:1472:NAG:C4	5:A:1476:BMA:C1	2.62	0.76
4:A:1471:NAG:C4	4:A:1472:NAG:C1	2.64	0.76
1:A:172:ARG:CD	1:A:209:ARG:NH2	2.49	0.75
5:A:1480:BMA:O6	5:A:1481:BMA:H3	1.90	0.70

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:292:ARG:HE	1:A:294:ASN:HD22	1.42	0.68
5:A:1476:BMA:C6	5:A:1480:BMA:C1	2.76	0.63
1:A:200:ASN:CG	4:A:1471:NAG:C1	2.65	0.63
1:A:98:HIS:HE1	1:A:419:ARG:HH11	1.47	0.61
5:A:1479:BMA:C2	2:B:2:MAN:O2	2.49	0.60
5:A:1479:BMA:C1	2:B:2:MAN:C2	2.77	0.60
1:A:98:HIS:CE1	1:A:419:ARG:HH11	2.19	0.59
1:A:97:TRP:H	1:A:395:GLN:HE22	1.49	0.58
1:A:168:THR:H	1:A:170:ASN:ND2	1.95	0.57
1:A:172:ARG:CD	1:A:209:ARG:HH22	2.15	0.57
1:A:200:ASN:ND2	4:A:1471:NAG:O5	2.32	0.57
1:A:200:ASN:ND2	4:A:1471:NAG:C2	2.65	0.54
1:A:465:GLU:H	1:A:465:GLU:CD	2.10	0.54
1:A:172:ARG:HD3	1:A:209:ARG:CZ	2.36	0.53
4:A:1472:NAG:O4	5:A:1476:BMA:C2	2.52	0.52
1:A:168:THR:OG1	1:A:170:ASN:ND2	2.43	0.51
1:A:292:ARG:HH21	1:A:294:ASN:ND2	2.10	0.50
5:A:1480:BMA:O6	5:A:1481:BMA:H5	2.12	0.50
1:A:170:ASN:HD22	1:A:170:ASN:H	1.61	0.48
1:A:82:ARG:O	1:A:187:LYS:HE2	2.13	0.48
5:A:1480:BMA:O3	5:A:1482:BMA:C2	2.61	0.47
1:A:284:ARG:HG2	1:A:284:ARG:HH11	1.80	0.46
1:A:140:ILE:HD11	7:A:2202:HOH:O	2.15	0.46
1:A:121:TYR:CG	1:A:228:SER:HA	2.52	0.45
1:A:364:ARG:HD2	1:A:375:GLU:OE2	2.16	0.45
1:A:367:SER:HB2	1:A:400:ASN:HD21	1.82	0.45
1:A:228:SER:HB3	1:A:350:LYS:HE2	1.99	0.43
1:A:292:ARG:HH21	1:A:294:ASN:HD21	1.66	0.43
1:A:204:VAL:HG22	1:A:206:TRP:NE1	2.34	0.43
1:A:330:ASP:CG	1:A:364:ARG:HH22	2.22	0.43
1:A:437:TRP:CD1	4:A:1475:NAG:H82	2.55	0.41
1:A:320:PRO:O	1:A:331:PRO:HD2	2.21	0.41
1:A:98:HIS:HD2	1:A:99:ILE:O	2.03	0.41
4:A:1473:NAG:C4	4:A:1474:NAG:C1	2.93	0.41
1:A:278:CYS:HB3	1:A:289:CYS:HB3	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	386/388 (100%)	366 (95%)	20 (5%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	341/341 (100%)	333 (98%)	8 (2%)	50	53

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	82	ARG
1	A	165	SER
1	A	170	ASN
1	A	204	VAL
1	A	221	ASN
1	A	364	ARG
1	A	416	GLU
1	A	465	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	95	ASN
1	A	98	HIS
1	A	146	ASN
1	A	170	ASN
1	A	200	ASN
1	A	216	ASN
1	A	221	ASN
1	A	233	HIS
1	A	294	ASN
1	A	346	ASN
1	A	392	GLN
1	A	395	GLN
1	A	400	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	MAN	B	1	2	11,11,12	0.48	0	15,15,17	2.29	4 (26%)
2	MAN	B	2	2	11,11,12	0.53	0	15,15,17	1.56	3 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MAN	B	1	2	-	0/2/19/22	0/1/1/1
2	MAN	B	2	2	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
2	B	1	MAN	C1-O5-C5	6.11	120.47	112.19
2	B	1	MAN	O2-C2-C1	-4.77	99.39	109.15
2	B	2	MAN	C1-O5-C5	3.24	116.59	112.19
2	B	2	MAN	C1-C2-C3	2.95	113.29	109.67
2	B	2	MAN	O5-C1-C2	2.77	115.05	110.77
2	B	1	MAN	C1-C2-C3	2.75	113.05	109.67
2	B	1	MAN	O6-C6-C5	-2.18	103.82	111.29

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	2	MAN	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 1 is monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	SIA	A	1470	-	21,21,21	1.12	2 (9%)	25,31,31	1.58	6 (24%)
3	SIA	A	1469	-	21,21,21	1.47	2 (9%)	25,31,31	1.92	6 (24%)
5	BMA	A	1481	-	11,11,12	0.77	0	15,15,17	1.75	4 (26%)
5	BMA	A	1476	-	11,11,12	0.30	0	15,15,17	1.48	3 (20%)
4	NAG	A	1473	-	14,14,15	0.39	0	17,19,21	1.38	2 (11%)
5	BMA	A	1480	-	11,11,12	0.40	0	15,15,17	1.78	2 (13%)
4	NAG	A	1475	-	14,14,15	0.44	0	17,19,21	1.41	2 (11%)



Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	BMA	A	1479	-	11,11,12	0.72	0	15,15,17	1.71	2 (13%)
4	NAG	A	1474	-	14,14,15	0.72	0	17,19,21	2.44	3 (17%)
5	BMA	A	1482	-	11,11,12	0.57	0	15,15,17	1.85	3 (20%)
4	NAG	A	1472	-	14,14,15	0.46	0	17,19,21	1.69	5 (29%)
4	NAG	A	1471	-	14,14,15	0.50	0	17,19,21	1.85	6 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SIA	A	1470	-	-	7/20/38/38	0/1/1/1
3	SIA	A	1469	-	-	3/20/38/38	0/1/1/1
5	BMA	A	1481	-	-	0/2/19/22	0/1/1/1
5	BMA	A	1476	-	-	0/2/19/22	0/1/1/1
4	NAG	A	1473	-	-	0/6/23/26	0/1/1/1
5	BMA	A	1480	-	-	0/2/19/22	0/1/1/1
4	NAG	A	1475	-	-	2/6/23/26	0/1/1/1
5	BMA	A	1479	-	-	0/2/19/22	0/1/1/1
4	NAG	A	1474	-	-	2/6/23/26	0/1/1/1
5	BMA	A	1482	-	-	2/2/19/22	0/1/1/1
4	NAG	A	1472	-	-	0/6/23/26	0/1/1/1
4	NAG	A	1471	-	-	0/6/23/26	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1469	SIA	O2-C2	4.30	1.45	1.39
3	A	1470	SIA	O2-C2	3.40	1.44	1.39
3	A	1469	SIA	C3-C2	3.29	1.55	1.51
3	A	1470	SIA	C3-C2	2.47	1.54	1.51

All (44) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1474	NAG	C2-N2-C7	6.99	132.86	122.90
3	A	1469	SIA	O1A-C1-C2	-6.13	114.30	123.59
4	A	1474	NAG	C4-C3-C2	-5.45	103.03	111.02
5	A	1480	BMA	C1-O5-C5	5.08	119.08	112.19

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	1479	BMA	C1-O5-C5	4.82	118.72	112.19
5	A	1482	BMA	C1-O5-C5	4.81	118.71	112.19
3	A	1470	SIA	O1A-C1-C2	-4.27	117.12	123.59
4	A	1475	NAG	C1-C2-N2	-4.10	103.49	110.49
5	A	1479	BMA	O5-C1-C2	3.64	116.38	110.77
5	A	1481	BMA	C1-C2-C3	3.58	114.07	109.67
3	A	1469	SIA	O6-C6-C5	3.55	113.24	109.78
5	A	1482	BMA	C1-C2-C3	3.49	113.95	109.67
4	A	1472	NAG	O5-C1-C2	3.37	116.61	111.29
4	A	1474	NAG	O5-C1-C2	-3.17	106.29	111.29
4	A	1471	NAG	C4-C3-C2	-3.10	106.47	111.02
5	A	1481	BMA	O5-C1-C2	3.05	115.48	110.77
5	A	1481	BMA	C1-O5-C5	3.05	116.32	112.19
3	A	1470	SIA	C6-C5-N5	-3.03	105.88	110.91
4	A	1473	NAG	O5-C5-C6	3.01	111.92	107.20
5	A	1480	BMA	C3-C4-C5	-3.01	104.88	110.24
4	A	1471	NAG	O4-C4-C3	-2.96	103.50	110.35
5	A	1476	BMA	C1-C2-C3	-2.96	106.03	109.67
5	A	1482	BMA	O5-C1-C2	2.94	115.31	110.77
4	A	1471	NAG	O5-C5-C6	2.93	111.79	107.20
3	A	1470	SIA	C3-C4-C5	2.81	114.30	109.98
3	A	1469	SIA	O2-C2-C3	2.79	113.43	109.40
5	A	1476	BMA	O3-C3-C2	-2.79	104.66	109.99
4	A	1475	NAG	O5-C1-C2	2.77	115.66	111.29
4	A	1473	NAG	C1-C2-N2	-2.69	105.89	110.49
5	A	1481	BMA	O5-C5-C6	2.68	111.40	107.20
4	A	1471	NAG	C1-C2-N2	-2.64	105.98	110.49
3	A	1469	SIA	O2-C2-C1	-2.58	105.51	110.76
4	A	1472	NAG	C2-N2-C7	-2.58	119.23	122.90
4	A	1471	NAG	C1-O5-C5	-2.56	108.73	112.19
3	A	1469	SIA	C6-C5-N5	-2.47	106.82	110.91
4	A	1472	NAG	C4-C3-C2	-2.30	107.64	111.02
3	A	1470	SIA	O10-C10-N5	2.22	126.04	121.95
4	A	1472	NAG	C8-C7-N2	2.19	119.81	116.10
5	A	1476	BMA	O6-C6-C5	-2.15	103.92	111.29
3	A	1470	SIA	O6-C6-C5	2.15	111.87	109.78
4	A	1472	NAG	C3-C4-C5	2.15	114.07	110.24
3	A	1469	SIA	O6-C6-C7	2.14	110.59	107.29
3	A	1470	SIA	O2-C2-C1	-2.09	106.52	110.76
4	A	1471	NAG	C8-C7-N2	-2.00	112.70	116.10

There are no chirality outliers.

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1470	SIA	O8-C8-C9-O9
3	A	1470	SIA	C6-C7-C8-O8
3	A	1470	SIA	C7-C8-C9-O9
3	A	1470	SIA	O7-C7-C8-C9
3	A	1470	SIA	C6-C7-C8-C9
4	A	1475	NAG	C4-C5-C6-O6
3	A	1470	SIA	O7-C7-C8-O8
5	A	1482	BMA	C4-C5-C6-O6
4	A	1475	NAG	O5-C5-C6-O6
5	A	1482	BMA	O5-C5-C6-O6
4	A	1474	NAG	C3-C2-N2-C7
3	A	1469	SIA	O1A-C1-C2-C3
4	A	1474	NAG	C1-C2-N2-C7
3	A	1469	SIA	O1A-C1-C2-O2
3	A	1470	SIA	O1A-C1-C2-O2
3	A	1469	SIA	O1B-C1-C2-O6


There are no ring outliers.


10 monomers are involved in 25 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	1481	BMA	2	0
5	A	1476	BMA	5	0
4	A	1473	NAG	3	0
5	A	1480	BMA	6	0
4	A	1475	NAG	3	0
5	A	1479	BMA	3	0
4	A	1474	NAG	3	0
5	A	1482	BMA	2	0
4	A	1472	NAG	5	0
4	A	1471	NAG	7	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	2

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	333:VAL	C	335:GLY	N	1.67
1	A	392:GLN	C	394:GLY	N	1.61

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	388/388 (100%)	-0.63	1 (0%) 94   95	8, 13, 20, 41	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	82	ARG	3.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	MAN	B	1	11/12	0.94	0.09	16,19,23,24	0
2	MAN	B	2	11/12	0.97	0.08	14,17,19,19	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	SIA	A	1470	21/21	0.53	0.34	55,62,63,64	0
5	BMA	A	1481	11/12	0.54	0.23	74,76,77,77	0
4	NAG	A	1474	14/15	0.75	0.25	49,51,53,54	0
5	BMA	A	1482	11/12	0.83	0.21	41,46,47,47	0
3	SIA	A	1469	21/21	0.87	0.16	18,26,28,33	0
4	NAG	A	1475	14/15	0.88	0.21	40,42,44,44	0
5	BMA	A	1480	11/12	0.92	0.16	30,32,34,34	0
4	NAG	A	1473	14/15	0.95	0.15	20,24,27,28	0
4	NAG	A	1471	14/15	0.95	0.11	14,18,23,27	0
4	NAG	A	1472	14/15	0.96	0.09	14,17,19,24	0
5	BMA	A	1476	11/12	0.96	0.09	14,18,19,22	0
5	BMA	A	1479	11/12	0.96	0.09	13,15,16,16	0
6	CA	A	1483	1/1	1.00	0.03	15,15,15,15	0

The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.