



Full wwPDB EM Validation Report ⓘ

Nov 19, 2023 – 04:08 PM JST

PDB ID : 7C4V
EMDB ID : EMD-30289
Title : MicroED structure of anorthic Vancomycin at 1.05 Å resolution
Authors : Fan, Q.; Zhou, H.; Li, X.; Wang, J.
Deposited on : 2020-05-18
Resolution : 1.05 Å (reported)
Based on initial model : 1FVM

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev70
Mogul : 1.8.5 (274361), CSD as541be (2020)
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : **NOT EXECUTED**
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON CRYSTALLOGRAPHY

The reported resolution of this entry is 1.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	7	
1	B	7	
2	C	2	
2	D	2	

2 Entry composition i


There are 4 unique types of molecules in this entry. The entry contains 246 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Vancomycin.

Mol	Chain	Residues	Atoms				AltConf	Trace	
			Total	C	Cl	N			O
1	A	7	80	53	2	8	17	0	0
1	B	7	80	53	2	8	17	0	0

- Molecule 2 is an oligosaccharide called vancosamine-(1-2)-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
2	C	2	21	13	1	7	0	0
2	D	2	21	13	1	7	0	0

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
3	A	1	Total	Cl	0
			1	1	

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		AltConf
4	A	20	Total	O	0
			20	20	
4	B	23	Total	O	0
			23	23	

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Vancomycin

Chain A: 



- Molecule 1: Vancomycin

Chain B: 



- Molecule 2: vancosamine-(1-2)-beta-D-glucopyranose

Chain C: 



- Molecule 2: vancosamine-(1-2)-beta-D-glucopyranose

Chain D: 



4 Experimental information

Property	Value	Source
EM reconstruction method	CRYSTALLOGRAPHY	Depositor
Imposed symmetry	3D CRYSTAL, $a=13.97 \text{ \AA}$, $b=18.55 \text{ \AA}$, $c=23.86 \text{ \AA}$, $\alpha=109.52^\circ$, $\beta=97.33^\circ$, $\gamma=106.58^\circ$, space group=P 1	Depositor
Number of images used	Not provided	
Resolution determination method	DIFFRACTION PATTERN/LAYERLINES	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TECNAI F20	Depositor
Voltage (kV)	200	Depositor
Electron dose ($e^-/\text{\AA}^2$)	0.0286	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN ULTRASCAN 4000 (4k x 4k)	Depositor
Maximum map value	12.975	Depositor
Minimum map value	-2.182	Depositor
Average map value	0.011	Depositor
Map value standard deviation	1.003	Depositor
Recommended contour level	1.6	Depositor
Map size (\AA)	54.327003, 32.0788, 43.979004	wwPDB
Map dimensions	170, 124, 210	wwPDB
Map angles ($^\circ$)	109.517, 97.327, 106.58	wwPDB
Pixel spacing (\AA)	0.2587, 0.2587, 0.2587	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 3FG, OMY, GHP, BGC, OMZ, CL, RER, MLU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.06	0/7	1.33	0/8
1	B	1.31	0/7	1.64	0/8
All	All	1.19	0/14	1.49	0/16

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	80	0	43	2	0
1	B	80	0	42	0	0
2	C	21	0	22	0	0
2	D	21	0	22	2	0
3	A	1	0	0	1	0
4	A	20	0	0	0	1
4	B	23	0	0	1	2
All	All	246	0	129	4	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (4) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:1:BGC:O6	2:D:1:BGC:O4	2.23	0.55
1:A:4:GHP:HA	3:A:101:CL:CL	2.47	0.52
1:A:3:ASN:C	1:A:4:GHP:H6	2.28	0.52
4:B:123:HOH:O	2:D:2:RER:N3	2.35	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:201:HOH:O	4:B:113:HOH:O[1_655]	2.15	0.05
4:B:107:HOH:O	4:B:112:HOH:O[1_455]	2.16	0.04

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1/7 (14%)	0	0	1 (100%)	0	0
1	B	1/7 (14%)	0	0	1 (100%)	0	0
All	All	2/14 (14%)	0	0	2 (100%)	0	0

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	3	ASN
1	B	3	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1/1 (100%)	1 (100%)	0	100	100
1	B	1/1 (100%)	1 (100%)	0	100	100
All	All	2/2 (100%)	2 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

12 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	GHP	A	5	1	10,11,12	2.37	2 (20%)	11,14,16	1.24	1 (9%)
1	3FG	B	7	1	12,13,13	2.12	2 (16%)	14,18,18	1.29	2 (14%)
1	MLU	B	1	1	7,8,9	1.36	1 (14%)	6,9,11	1.48	2 (33%)
1	3FG	A	7	1	12,13,13	2.40	3 (25%)	14,18,18	1.26	2 (14%)
1	OMY	A	6	1	12,14,15	1.53	2 (16%)	17,19,21	2.60	4 (23%)
1	OMZ	B	2	1	12,14,15	1.89	2 (16%)	17,19,21	1.27	2 (11%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	OMZ	A	2	1	12,14,15	2.90	5 (41%)	17,19,21	1.33	2 (11%)
1	OMY	B	6	1	12,14,15	1.47	1 (8%)	17,19,21	3.50	10 (58%)
1	GHP	A	4	2,1	10,11,12	2.03	3 (30%)	11,14,16	1.20	1 (9%)
1	GHP	B	4	2,1	10,11,12	2.38	4 (40%)	11,14,16	0.90	0
1	MLU	A	1	1	7,8,9	1.07	1 (14%)	6,9,11	1.35	1 (16%)
1	GHP	B	5	1	10,11,12	2.93	1 (10%)	11,14,16	0.95	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	GHP	A	5	1	-	0/4/6/8	0/1/1/1
1	3FG	B	7	1	-	2/8/8/8	0/1/1/1
1	MLU	B	1	1	-	5/5/8/10	-
1	3FG	A	7	1	-	2/8/8/8	0/1/1/1
1	OMY	A	6	1	-	1/9/10/12	0/1/1/1
1	OMZ	B	2	1	-	1/9/10/12	0/1/1/1
1	OMZ	A	2	1	-	1/9/10/12	0/1/1/1
1	OMY	B	6	1	-	1/9/10/12	0/1/1/1
1	GHP	A	4	2,1	-	2/4/6/8	0/1/1/1
1	GHP	B	4	2,1	-	0/4/6/8	0/1/1/1
1	MLU	A	1	1	-	0/5/8/10	-
1	GHP	B	5	1	-	0/4/6/8	0/1/1/1

All (27) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	5	GHP	C1-CA	-8.62	1.43	1.52
1	A	7	3FG	CB-CA	-6.81	1.44	1.52
1	A	5	GHP	C1-CA	-6.49	1.45	1.52
1	B	4	GHP	C1-CA	-5.47	1.46	1.52
1	A	2	OMZ	CG-CB	-5.42	1.43	1.51
1	B	7	3FG	CB-CA	-5.24	1.46	1.52
1	A	2	OMZ	OC-CB	-5.02	1.32	1.42
1	B	2	OMZ	OC-CB	-4.72	1.33	1.42
1	A	2	OMZ	CA-N	-4.32	1.34	1.47
1	A	4	GHP	C1-CA	-4.29	1.48	1.52
1	A	6	OMY	OCZ-CZ	4.23	1.45	1.36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	6	OMY	OCZ-CZ	3.65	1.43	1.36
1	B	4	GHP	CA-C	-3.28	1.44	1.51
1	B	2	OMZ	CA-N	-3.18	1.37	1.47
1	B	4	GHP	C6-C1	-2.99	1.34	1.39
1	A	4	GHP	CA-C	-2.74	1.45	1.51
1	A	2	OMZ	OH-CZ	-2.67	1.31	1.36
1	A	2	OMZ	CE1-CL	-2.66	1.67	1.73
1	B	7	3FG	CG1-CD1	-2.59	1.35	1.39
1	B	1	MLU	CA-N	-2.40	1.43	1.47
1	A	4	GHP	C6-C5	-2.40	1.34	1.38
1	A	6	OMY	CG-CB	-2.16	1.48	1.51
1	A	5	GHP	C3-C2	-2.15	1.34	1.38
1	A	7	3FG	OD1-CD1	-2.15	1.32	1.37
1	A	7	3FG	CG1-CB	-2.14	1.35	1.39
1	A	1	MLU	O-C	2.09	1.28	1.19
1	B	4	GHP	C6-C5	-2.02	1.35	1.38

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	6	OMY	CG-CB-CA	-11.96	95.48	111.49
1	A	6	OMY	CG-CB-CA	-8.97	99.48	111.49
1	B	6	OMY	CD2-CG-CD1	4.08	123.47	118.76
1	B	2	OMZ	O-C-CA	-3.53	115.54	124.78
1	A	2	OMZ	O-C-CA	-3.32	116.08	124.78
1	A	6	OMY	CD2-CG-CD1	3.31	122.58	118.76
1	B	6	OMY	CD1-CG-CB	-2.82	114.74	119.55
1	B	6	OMY	CE2-CZ-CE1	2.75	121.22	118.55
1	B	7	3FG	CD2-CG2-CB	-2.65	117.83	120.11
1	B	6	OMY	CZ-CE1-CL	2.55	124.32	119.53
1	A	4	GHP	C1-CA-N	-2.45	106.54	112.40
1	B	2	OMZ	OC-CB-CA	2.40	112.35	107.28
1	A	2	OMZ	OC-CB-CA	2.38	112.31	107.28
1	B	7	3FG	OXT-C-O	2.35	129.41	124.09
1	B	1	MLU	CN-N-CA	-2.24	106.66	113.64
1	B	6	OMY	CD2-CE2-CZ	-2.24	118.20	120.50
1	B	6	OMY	CD1-CE1-CL	-2.22	114.90	118.49
1	A	6	OMY	CD2-CE2-CZ	-2.18	118.27	120.50
1	B	6	OMY	CE2-CD2-CG	-2.16	119.03	121.20
1	A	1	MLU	O-C-CA	-2.12	119.23	124.78
1	B	6	OMY	ODE-CB-CA	2.12	111.75	107.28
1	A	5	GHP	C2-C1-CA	-2.09	117.28	120.65

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	6	OMY	CE1-CD1-CG	-2.07	117.28	120.29
1	A	6	OMY	CE1-CD1-CG	-2.07	117.28	120.29
1	A	7	3FG	CD2-CZ-CD1	-2.06	116.74	119.20
1	A	7	3FG	CG2-CB-CA	2.02	122.92	119.71
1	B	1	MLU	CD2-CG-CB	-2.02	103.69	111.11

There are no chirality outliers.

All (15) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	B	1	MLU	O-C-CA-CB
1	A	2	OMZ	O-C-CA-CB
1	B	2	OMZ	O-C-CA-CB
1	A	6	OMY	O-C-CA-CB
1	B	6	OMY	O-C-CA-CB
1	B	1	MLU	N-CA-CB-CG
1	B	1	MLU	CA-CB-CG-CD2
1	B	1	MLU	CA-CB-CG-CD1
1	A	7	3FG	C-CA-CB-CG2
1	B	7	3FG	C-CA-CB-CG1
1	B	7	3FG	C-CA-CB-CG2
1	A	4	GHP	C2-C1-CA-C
1	A	4	GHP	C6-C1-CA-C
1	B	1	MLU	C-CA-CB-CG
1	A	7	3FG	C-CA-CB-CG1

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	4	GHP	2	0

5.5 Carbohydrates [i](#)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	BGC	C	1	2,1	11,11,12	2.14	4 (36%)	15,15,17	2.17	5 (33%)
2	RER	C	2	2	6,10,11	1.77	2 (33%)	6,15,17	1.63	1 (16%)
2	BGC	D	1	2,1	11,11,12	1.54	4 (36%)	15,15,17	2.18	5 (33%)
2	RER	D	2	2	6,10,11	1.00	0	6,15,17	2.07	2 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	C	1	2,1	-	2/2/19/22	0/1/1/1
2	RER	C	2	2	-	-	0/1/1/1
2	BGC	D	1	2,1	-	1/2/19/22	0/1/1/1
2	RER	D	2	2	-	-	1/1/1/1

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	1	BGC	O5-C1	-4.24	1.36	1.43
2	C	1	BGC	C2-C3	-3.02	1.48	1.52
2	C	2	RER	O5-C1	-2.89	1.37	1.44
2	D	1	BGC	O2-C2	-2.53	1.38	1.43
2	C	2	RER	O5-C5	-2.41	1.38	1.43
2	C	1	BGC	O3-C3	-2.36	1.37	1.43
2	C	1	BGC	O5-C5	-2.33	1.38	1.43
2	D	1	BGC	O5-C5	-2.18	1.39	1.43
2	D	1	BGC	O5-C1	-2.15	1.40	1.43
2	D	1	BGC	O3-C3	-2.12	1.38	1.43

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	1	BGC	C1-C2-C3	-5.54	102.86	109.67
2	C	1	BGC	C1-C2-C3	-5.40	103.03	109.67
2	C	1	BGC	C6-C5-C4	-3.52	104.76	113.00
2	C	2	RER	C5A-C5-C4	-3.33	106.88	112.57
2	D	2	RER	O5-C5-C4	3.18	115.71	110.03

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	1	BGC	C6-C5-C4	-3.10	105.74	113.00
2	D	1	BGC	C2-C3-C4	-2.85	105.97	110.89
2	D	1	BGC	O3-C3-C4	-2.81	103.84	110.35
2	C	1	BGC	C2-C3-C4	-2.46	106.64	110.89
2	C	1	BGC	O2-C2-C3	-2.45	105.23	110.14
2	C	1	BGC	C3-C4-C5	-2.44	105.89	110.24
2	D	2	RER	C1-O5-C5	2.28	118.23	110.95
2	D	1	BGC	C3-C4-C5	-2.09	106.52	110.24

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	1	BGC	O5-C5-C6-O6
2	C	1	BGC	C4-C5-C6-O6
2	D	1	BGC	O5-C5-C6-O6



All (1) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	2	RER	C1-C2-C3-C4-C5-O5

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	1	BGC	1	0
2	D	2	RER	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-30289. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 105



Y Index: 62




Z Index: 85


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 16



Y Index: 2




Z Index: 11


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 1.6. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.

7.2 Volume estimate [i](#)



The volume at the recommended contour level is 5 nm³; this corresponds to an approximate mass of 5 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)

This section was not generated. The rotationally averaged power spectrum is only generated for cubic maps.


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-30289 and PDB model 7C4V. Per-residue inclusion information can be found in section 3 on page 4.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 1.6 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model


Q-score is only calculated for structures below 1.25 Å in resolution. The reported resolution for this entry is above 1.25 Å and so Q-score has not been calculated.

9.3 Atom inclusion mapped to coordinate model [\(i\)](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (1.6).






9.4 Atom inclusion [i](#)



At the recommended contour level, 71% of all backbone atoms, 85% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary [i](#)

The table lists the average atom inclusion at the recommended contour level (1.6) and Q-score for the entire model and for each chain.

Chain	Atom inclusion
All	 0.8460
A	 1.0000
B	 0.7850
C	 1.0000
D	 0.3810

