

wwPDB X-ray Structure Validation Summary Report (i)

Nov 19, 2023 – 11:35 PM JST

PDB ID : 7C7K

Title: Crystal Structure of Thioacyl-Glyceraldehyde-3-phosphate dehydrogenase

1(GAPDH 1) from Escherichia coli at 1.77 Angstrom resolution

Authors : Zhang, L.; Liu, M.R.; Bao, L.Y.; Yao, Y.C.; Bostrom, I.K.; Wang, Y.D.; Chen,

A.Q.; Li, J.X.; Gu, S.H.; Ji, C.N.

Deposited on : 2020-05-26

Resolution : 1.77 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 1.77 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
R_{free}	130704	9185 (1.80-1.76)
Clashscore	141614	10184 (1.80-1.76)
Ramachandran outliers	138981	10051 (1.80-1.76)
Sidechain outliers	138945	10050 (1.80-1.76)
RSRZ outliers	127900	9032 (1.80-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain				
1	О	352	91%	5% •			
1	Р	352	88%	6% 5%			
1	Q	352	86%	9% • 5%			
1	R	352	88%	7% 5%			



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	EDO	Q	404	-	-	X	-
5	PEG	O	407	-	-	X	-
6	ACN	P	404	-	-	X	-
6	ACN	Q	406	-	-	X	-



2 Entry composition (i)

There are 8 unique types of molecules in this entry. The entry contains 11660 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Glyceraldehyde-3-phosphate dehydrogenase.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	0	337	Total	С	N	Ο	S	0	5	0
1		337	2586	1631	437	510	8	0		U
1	Р	333	Total	С	N	О	S	0	5	0
1	1	333	2554	1613	430	504	7	0	9	U
1	0	335	Total	С	N	О	S	0	9	0
1	Q	333	2583	1631	436	509	7	U	9	U
1	R	334	Total	С	N	О	S	0	3	0
1	11	334	2547	1608	428	504	7		3	U

There are 76 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
О	-18	HIS	-	expression tag	UNP A0A140NCK4
О	-17	HIS	-	expression tag	UNP A0A140NCK4
О	-16	HIS	-	expression tag	UNP A0A140NCK4
О	-15	HIS	-	expression tag	UNP A0A140NCK4
О	-14	HIS	-	expression tag	UNP A0A140NCK4
О	-13	HIS	-	expression tag	UNP A0A140NCK4
О	-12	SER	-	expression tag	UNP A0A140NCK4
О	-11	SER	-	expression tag	UNP A0A140NCK4
О	-10	GLY	-	expression tag	UNP A0A140NCK4
О	-9	LEU	-	expression tag	UNP A0A140NCK4
О	-8	VAL	-	expression tag	UNP A0A140NCK4
О	-7	PRO	-	expression tag	UNP A0A140NCK4
О	-6	ARG	-	expression tag	UNP A0A140NCK4
О	-5	GLY	-	expression tag	UNP A0A140NCK4
О	-4	SER	-	expression tag	UNP A0A140NCK4
О	-3	HIS	-	expression tag	UNP A0A140NCK4
О	-2	MET	-	expression tag	UNP A0A140NCK4
О	-1	ALA	-	expression tag	UNP A0A140NCK4
О	0	SER		expression tag	UNP A0A140NCK4
Р	-18	HIS	-	expression tag	UNP A0A140NCK4
Р	-17	HIS	-	expression tag	UNP A0A140NCK4



 $Continued\ from\ previous\ page...$


P -16 HIS P -15 HIS P -14 HIS P -13 HIS P -12 SER P -11 SER P -10 GLY	- - - - -	expression tag expression tag expression tag expression tag expression tag expression tag	UNP A0A140NCK4 UNP A0A140NCK4 UNP A0A140NCK4 UNP A0A140NCK4 UNP A0A140NCK4
P -14 HIS P -13 HIS P -12 SER P -11 SER P -10 GLY	-	expression tag expression tag expression tag	UNP A0A140NCK4 UNP A0A140NCK4
P -13 HIS P -12 SER P -11 SER P -10 GLY	-	expression tag expression tag	UNP A0A140NCK4
P -12 SER P -11 SER P -10 GLY	-	expression tag	
P -11 SER P -10 GLY		1 0	UNP A0A140NCK4
P -10 GLY	-	evaression tag	
	-	capicosion dag	UNP A0A140NCK4
		expression tag	UNP A0A140NCK4
P -9 LEU	-	expression tag	UNP A0A140NCK4
P -8 VAL	-	expression tag	UNP A0A140NCK4
P -7 PRO	-	expression tag	UNP A0A140NCK4
P -6 ARG	-	expression tag	UNP A0A140NCK4
P -5 GLY	-	expression tag	UNP A0A140NCK4
P -4 SER	-	expression tag	UNP A0A140NCK4
P -3 HIS	-	expression tag	UNP A0A140NCK4
P -2 MET	-	expression tag	UNP A0A140NCK4
P -1 ALA	-	expression tag	UNP A0A140NCK4
P 0 SER	-	expression tag	UNP A0A140NCK4
Q -18 HIS	-	expression tag	UNP A0A140NCK4
Q -17 HIS	-	expression tag	UNP A0A140NCK4
Q -16 HIS	-	expression tag	UNP A0A140NCK4
Q -15 HIS	-	expression tag	UNP A0A140NCK4
Q -14 HIS	-	expression tag	UNP A0A140NCK4
Q -13 HIS	-	expression tag	UNP A0A140NCK4
Q -12 SER	-	expression tag	UNP A0A140NCK4
Q -11 SER	-	expression tag	UNP A0A140NCK4
Q -10 GLY	-	expression tag	UNP A0A140NCK4
Q -9 LEU	-	expression tag	UNP A0A140NCK4
Q -8 VAL	-	expression tag	UNP A0A140NCK4
Q -7 PRO	-	expression tag	UNP A0A140NCK4
Q -6 ARG	-	expression tag	UNP A0A140NCK4
Q -5 GLY	-	expression tag	UNP A0A140NCK4
Q -4 SER	-	expression tag	UNP A0A140NCK4
Q -3 HIS	-	expression tag	UNP A0A140NCK4
Q -2 MET	-	expression tag	UNP A0A140NCK4
Q -1 ALA	-	expression tag	UNP A0A140NCK4
Q 0 SER	-	expression tag	UNP A0A140NCK4
R -18 HIS	-	expression tag	UNP A0A140NCK4
R -17 HIS	-	expression tag	UNP A0A140NCK4
R -16 HIS	-	expression tag	UNP A0A140NCK4
R -15 HIS	-	expression tag	UNP A0A140NCK4
R -14 HIS	-	expression tag	UNP A0A140NCK4
R -13 HIS	-	expression tag	UNP A0A140NCK4



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
R	-12	SER	-	expression tag	UNP A0A140NCK4
R	-11	SER	-	expression tag	UNP A0A140NCK4
R	-10	GLY	-	expression tag	UNP A0A140NCK4
R	-9	LEU	-	expression tag	UNP A0A140NCK4
R	-8	VAL	-	expression tag	UNP A0A140NCK4
R	-7	PRO	-	expression tag	UNP A0A140NCK4
R	-6	ARG	-	expression tag	UNP A0A140NCK4
R	-5	GLY	-	expression tag	UNP A0A140NCK4
R	-4	SER	-	expression tag	UNP A0A140NCK4
R	-3	HIS	-	expression tag	UNP A0A140NCK4
R	-2	MET	-	expression tag	UNP A0A140NCK4
R	-1	ALA	-	expression tag	UNP A0A140NCK4
R	0	SER	-	expression tag	UNP A0A140NCK4


• Molecule 2 is GLYCERALDEHYDE-3-PHOSPHATE (three-letter code: G3H) (formula: $C_3H_7O_6P$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	\circ	1	Total C O P	0	0
	O	1	10 3 6 1	0	0
2	Р	1	Total C O P	0	0
	1	1	10 3 6 1	0	U
2	0	1	Total C O P	0	0
	Q	1	10 3 6 1	0	
2	R	1	Total C O P	0	0
	11	1	10 3 6 1	U	U




• Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O_4P) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	О	1	Total O P 5 4 1	0	0
3	О	1	Total O P 5 4 1	0	0


• Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	О	1	Total C O 4 2 2	0	0
4	О	1	Total C O 4 2 2	0	0
4	О	1	Total C O 4 2 2	0	0
4	Р	1	Total C O 4 2 2	0	0
4	Р	1	Total C O 4 2 2	0	0
4	Р	1	Total C O 4 2 2	0	0
4	Q	1	Total C O 4 2 2	0	0
4	Q	1	Total C O 4 2 2	0	0
4	Q	1	Total C O 4 2 2	0	0
4	Q	1	Total C O 4 2 2	0	0
4	R	1	Total C O 4 2 2	0	0
4	R	1	Total C O 4 2 2	0	0


 \bullet Molecule 5 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3)$ (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	О	1	Total C O 7 4 3	0	0
5	R	1	Total C O 7 4 3	0	0


• Molecule 6 is ACETONE (three-letter code: ACN) (formula: C_3H_6O) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	Р	1	Total C O 4 3 1	0	0
6	Q	1	Total C O 4 3 1	0	0
6	R	1	Total C O 4 3 1	0	0

• Molecule 7 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: $C_{21}H_{27}N_7O_{14}P_2$) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	Q	1	Total C N O P 44 21 7 14 2	0	0
7	R	1	Total C O P 10 1 7 2	0	0


• Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	О	365	Total O 365 365	0	0
8	Р	322	Total O 323 323	0	1
8	Q	277	Total O 278 278	0	1
8	R	245	Total O 246 246	0	1



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.









4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants	90.16Å 90.16Å 345.06Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.49 - 1.77	Depositor
rtesolution (A)	48.49 - 1.77	EDS
% Data completeness	99.6 (48.49-1.77)	Depositor
(in resolution range)	99.6 (48.49-1.77)	EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	4.84 (at 1.77Å)	Xtriage
Refinement program	REFMAC 5.8.0131	Depositor
D D.	0.143 , 0.181	Depositor
R, R_{free}	0.158 , 0.190	DCC
R_{free} test set	6788 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å ²)	21.4	Xtriage
Anisotropy	0.019	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.34, 41.3	EDS
L-test for twinning ²	$ < L > = 0.50, < L^2> = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	11660	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.46% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: G3H, PO4, PEG, ACN, NAD, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	Bond lengths		ond angles
WIOI		RMSZ	# Z > 5	RMSZ	# Z >5
1	О	1.00	2/2629~(0.1%)	1.04	7/3566~(0.2%)
1	Р	1.00	3/2595~(0.1%)	1.08	$10/3521 \ (0.3\%)$
1	Q	1.06	4/2624~(0.2%)	1.13	$12/3560 \ (0.3\%)$
1	R	0.98	0/2588	0.99	5/3511 (0.1%)
All	All	1.01	9/10436 (0.1%)	1.06	34/14158 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	Q	0	1

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}(\text{\AA})$
1	Q	56	TRP	CE3-CZ3	8.63	1.53	1.38
1	Q	202	GLU	CD-OE2	7.69	1.34	1.25
1	Р	202	GLU	CD-OE2	6.79	1.33	1.25
1	О	216	GLY	CA-C	-6.54	1.41	1.51
1	O	96	CYS	CB-SG	-5.94	1.72	1.81

The worst 5 of 34 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$Ideal(^{o})$
1	P	18	ARG	NE-CZ-NH1	15.80	128.20	120.30
1	Q	18	ARG	NE-CZ-NH1	12.49	126.55	120.30
1	Q	18	ARG	NE-CZ-NH2	-10.65	114.97	120.30
1	Р	18	ARG	NE-CZ-NH2	-10.19	115.21	120.30



Continued from previous page...

I	Mol	Chain	Res	Type	Atoms	${f Z}$	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
	1	R	193	ASP	CB-CG-OD1	9.38	126.75	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

	Mol	Chain	Res	Type	Group
ſ	1	Q	34	ASP	Mainchain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	О	2586	0	2595	12	0
1	Р	2554	0	2573	13	0
1	Q	2583	0	2611	30	0
1	R	2547	0	2564	17	0
2	О	10	0	4	0	0
2	Р	10	0	4	0	0
2	Q	10	0	4	3	0
2	R	10	0	4	0	0
3	О	10	0	0	0	0
4	О	12	0	18	4	0
4	Р	12	0	18	1	0
4	Q	16	0	24	6	0
4	R	8	0	12	2	0
5	О	7	0	10	6	0
5	R	7	0	10	0	0
6	Р	4	0	6	5	0
6	Q	4	0	6	7	0
6	R	4	0	6	3	0
7	Q	44	0	26	12	0
7	R	10	0	0	1	0
8	О	365	0	0	4	0
8	Р	323	0	0	6	0
8	Q	278	0	0	7	0
8	R	246	0	0	9	0
All	All	11660	0	10495	89	0



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 89 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$
1:R:232:ARG:HH12	6:R:406:ACN:H23	1.20	1.06
1:Q:232:ARG:HH12	6:Q:406:ACN:H13	1.28	0.96
7:Q:401:NAD:C5N	2:Q:402:G3H:O1	2.20	0.89
4:Q:404:EDO:H22	8:Q:682:HOH:O	1.79	0.81
7:Q:401:NAD:C4N	2:Q:402:G3H:C1	2.59	0.80

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percer	ntiles
1	О	340/352~(97%)	331 (97%)	9 (3%)	0	100	100
1	Р	$336/352 \ (96\%)$	325 (97%)	11 (3%)	0	100	100
1	Q	342/352 (97%)	332 (97%)	10 (3%)	0	100	100
1	R	335/352~(95%)	322 (96%)	13 (4%)	0	100	100
All	All	1353/1408 (96%)	1310 (97%)	43 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Perce	entiles
1	О	281/289 (97%)	281 (100%)	0	100	100
1	Р	278/289 (96%)	277 (100%)	1 (0%)	91	88
1	Q	279/289 (96%)	277 (99%)	2 (1%)	84	79
1	R	277/289 (96%)	275 (99%)	2 (1%)	84	79
All	All	1115/1156 (96%)	1110 (100%)	5 (0%)	88	88

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	Р	229	HIS
1	Q	229	HIS
1	Q	276	GLU
1	R	229	HIS
1	R	313	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

25 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the



expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trmo	Chain	Res	Link	Во	nd leng	ths	В	ond ang	gles
IVIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2
2	G3H	Р	401	1	8,9,9	1.32	1 (12%)	10,12,12	1.03	0
2	G3H	R	401	1	8,9,9	1.86	3 (37%)	10,12,12	0.96	0
4	EDO	О	404	-	3,3,3	0.76	0	2,2,2	1.29	0
4	EDO	Р	405	-	3,3,3	0.61	0	2,2,2	0.72	0
5	PEG	О	407	-	6,6,6	0.70	0	5,5,5	1.03	0
6	ACN	R	406	-	3,3,3	0.53	0	3,3,3	0.87	0
4	EDO	О	405	ı	3,3,3	1.12	0	2,2,2	1.38	0
4	EDO	Р	402	-	3,3,3	0.45	0	2,2,2	0.48	0
3	PO4	О	402	-	4,4,4	1.48	1 (25%)	6,6,6	2.00	2 (33%)
4	EDO	Q	407	-	3,3,3	1.22	0	2,2,2	1.92	1 (50%)
5	PEG	R	404	-	6,6,6	0.74	0	5,5,5	0.70	0
3	PO4	О	406	1	4,4,4	0.93	0	6,6,6	1.01	0
4	EDO	Q	403	-	3,3,3	0.52	0	2,2,2	1.52	1 (50%)
6	ACN	P	404	-	3,3,3	0.93	0	3,3,3	1.19	0
2	G3H	Q	402	1	8,9,9	1.42	1 (12%)	10,12,12	1.18	1 (10%)
4	EDO	R	403	-	3,3,3	0.49	0	2,2,2	0.24	0
4	EDO	Р	403	-	3,3,3	0.34	0	2,2,2	0.55	0
7	NAD	R	402	-	7,9,48	1.70	2 (28%)	12,14,73	1.98	4 (33%)
4	EDO	О	403	-	3,3,3	0.70	0	2,2,2	1.93	1 (50%)
4	EDO	Q	404	-	3,3,3	0.85	0	2,2,2	1.15	0
7	NAD	Q	401	-	42,48,48	2.11	7 (16%)	50,73,73	2.45	16 (32%)
6	ACN	Q	406	-	3,3,3	0.34	0	3,3,3	0.93	0
2	G3H	О	401	1	8,9,9	1.94	3 (37%)	10,12,12	1.62	3 (30%)
4	EDO	R	405	-	3,3,3	0.33	0	2,2,2	1.50	0
4	EDO	Q	405	-	3,3,3	0.17	0	2,2,2	1.99	1 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	G3H	Р	401	1	-	1/7/8/8	-
2	G3H	R	401	1	-	1/7/8/8	-
4	EDO	О	404	-	-	1/1/1/1	-
4	EDO	Р	405	-	-	1/1/1/1	-
5	PEG	О	407	-	-	0/4/4/4	-



Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	О	405	-	-	0/1/1/1	-
4	EDO	Р	402	-	-	1/1/1/1	-
4	EDO	Q	407	-	-	1/1/1/1	-
5	PEG	R	404	-	-	3/4/4/4	-
4	EDO	Q	403	-	-	1/1/1/1	-
2	G3H	Q	402	1	-	0/7/8/8	-
4	EDO	R	403	-	-	1/1/1/1	-
4	EDO	Р	403	-	-	1/1/1/1	-
7	NAD	R	402	-	-	0/9/9/62	-
4	EDO	О	403	-	-	1/1/1/1	-
4	EDO	Q	404	-	-	1/1/1/1	-
7	NAD	Q	401	ı	-	5/26/62/62	0/5/5/5
2	G3H	О	401	1	-	0/7/8/8	-
4	EDO	R	405	_	-	0/1/1/1	-
4	EDO	Q	405	_	_	0/1/1/1	_

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	Observed(Å)	$Ideal(\AA)$
7	Q	401	NAD	C2N-N1N	-9.17	1.23	1.35
7	Q	401	NAD	C2A-N3A	5.55	1.41	1.32
2	О	401	G3H	O1-C1	4.09	1.36	1.19
2	R	401	G3H	C3-C2	3.33	1.56	1.51
7	Q	401	NAD	C2A-N1A	3.31	1.40	1.33

The worst 5 of 30 bond angle outliers are listed below:

\mathbf{M}	ol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^{o})$	$ \operatorname{Ideal}(^o) $
7	7	Q	401	NAD	N3A-C2A-N1A	-9.64	113.61	128.68
7	7	Q	401	NAD	N6A-C6A-N1A	4.93	128.81	118.57
7	7	Q	401	NAD	C2A-N1A-C6A	4.68	126.77	118.75
7	7	R	402	NAD	O2A-PA-O1A	4.49	128.27	110.68
7	7	Q	401	NAD	C5A-C6A-N6A	-4.44	113.61	120.35

There are no chirality outliers.

5 of 19 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	Q	401	NAD	O4D-C1D-N1N-C2N
7	Q	401	NAD	O4D-C1D-N1N-C6N
7	Q	401	NAD	C2D-C1D-N1N-C2N



Continued from previous page...

Mol	Chain	Res	Type	Atoms
7	Q	401	NAD	C2D-C1D-N1N-C6N
5	R	404	PEG	O1-C1-C2-O2


There are no ring outliers.


13 monomers are involved in 44 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	О	404	EDO	2	0
4	Р	405	EDO	1	0
5	О	407	PEG	6	0
6	R	406	ACN	3	0
4	О	405	EDO	2	0
4	Q	407	EDO	1	0
6	Р	404	ACN	5	0
2	Q	402	G3H	3	0
4	R	403	EDO	2	0
7	R	402	NAD	1	0
4	Q	404	EDO	5	0
7	Q	401	NAD	12	0
6	Q	406	ACN	7	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











































































































5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\#\mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q < 0.9
1	О	337/352 (95%)	-0.42	2 (0%) 89 89	15, 19, 33, 53	0
1	Р	333/352 (94%)	-0.48	1 (0%) 94 93	14, 20, 35, 49	0
1	Q	335/352~(95%)	-0.31	1 (0%) 94 93	15, 20, 37, 57	0
1	R	334/352 (94%)	-0.16	7 (2%) 63 63	15, 24, 43, 64	0
All	All	1339/1408 (95%)	-0.34	11 (0%) 86 86	14, 21, 39, 64	0

The worst 5 of 11 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	R	0	SER	3.7
1	Q	-1	ALA	3.0
1	О	-2	MET	2.7
1	R	81	LYS	2.3
1	R	1	MET	2.3

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
5	PEG	R	404	7/7	0.69	0.16	49,55,57,59	0
4	EDO	Р	405	4/4	0.72	0.17	42,43,44,49	0
4	EDO	О	403	4/4	0.75	0.15	47,48,49,52	0
6	ACN	Р	404	4/4	0.77	0.18	21,22,30,37	0
3	PO4	O	406	5/5	0.78	0.21	59,64,64,86	0
4	EDO	R	403	4/4	0.79	0.15	40,44,53,54	0
4	EDO	Q	405	4/4	0.83	0.16	46,52,52,52	0
4	EDO	Р	402	4/4	0.85	0.13	39,43,44,48	0
4	EDO	Q	403	4/4	0.86	0.12	34,39,44,47	0
6	ACN	R	406	4/4	0.86	0.12	26,28,29,31	0
4	EDO	R	405	4/4	0.88	0.13	49,51,57,58	0
4	EDO	О	404	4/4	0.88	0.14	33,38,42,44	0
4	EDO	О	405	4/4	0.89	0.14	24,33,39,41	0
7	NAD	R	402	10/44	0.89	0.18	21,24,30,32	10
5	PEG	О	407	7/7	0.90	0.14	30,39,45,47	0
7	NAD	Q	401	44/44	0.91	0.14	13,23,27,27	44
4	EDO	Q	404	4/4	0.91	0.37	40,40,42,45	0
4	EDO	Q	407	4/4	0.94	0.19	20,28,31,36	0
3	PO4	О	402	5/5	0.95	0.18	25,30,34,39	0
4	EDO	Р	403	4/4	0.95	0.18	22,30,32,48	0
6	ACN	Q	406	4/4	0.95	0.17	21,21,25,29	0
2	G3H	Q	402	10/10	0.96	0.11	20,24,28,29	10
2	G3H	O	401	10/10	0.97	0.08	19,21,27,33	0
2	G3H	Р	401	10/10	0.98	0.06	20,21,24,24	0
2	G3H	R	401	10/10	0.99	0.06	22,24,27,30	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.














Electron density around EDO O 403:


 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)















Electron density around PO4 O 406: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${ m mF}_o{ m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)














Electron density around EDO P 402:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around EDO Q 403:


 $2 \text{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\text{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)
























Electron density around EDO O 405: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_{o}\text{-}\mathrm{DF}_{c}$ (at 3 rmsd) in purple (negative) and green (positive)



Electron density around NAD R 402: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${ m mF}_o{ m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









Electron density around NAD Q 401: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${ m mF}_o{ m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive) Electron density around EDO Q 404:

Electron density around EDO Q 404: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)
























Electron density around G3H Q 402:


 $2mF_o$ -DF_c (at 0.7 rmsd) in gray mF_o -DF_c (at 3 rmsd) in purple (negative) and green (positive)















 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)











6.5 Other polymers (i)

There are no such residues in this entry.

