



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 13, 2024 – 03:33 PM JST

PDB ID : 5CAD
Title : Crystal structure of the vicilin from Solanum melongena revealed existence of different anionic ligands in structurally similar pockets
Authors : Jain, A.; Kumar, A.; Salunke, D.M.
Deposited on : 2015-06-29
Resolution : 1.49 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2936 (1.50-1.50)
Clashscore	141614	3144 (1.50-1.50)
Ramachandran outliers	138981	3066 (1.50-1.50)
Sidechain outliers	138945	3064 (1.50-1.50)
RSRZ outliers	127900	2884 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	392	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	ACT	A	402	-	X	-	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3182 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called SM80.1 Vicilin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	379	2912	1858	521	525	8	0	0	0

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
2	A	1	1	1	0	0

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
3	A	1	4	2	2	0	0

- Molecule 4 is PYROGLUTAMIC ACID (three-letter code: PCA) (formula: C₅H₇NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
4	A	1	9	5	1	3	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	256	256	256	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: SM80.1 Vicilin



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	119.33Å 119.33Å 158.19Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	22.71 – 1.49 22.71 – 1.49	Depositor EDS
% Data completeness (in resolution range)	99.9 (22.71-1.49) 99.9 (22.71-1.49)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.87 (at 1.49Å)	Xtrriage
Refinement program	CNS 1.3	Depositor
R, R_{free}	0.199 , 0.210 0.193 , 0.204	Depositor DCC
R_{free} test set	3557 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	16.0	Xtrriage
Anisotropy	0.420	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 60.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.36$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3182	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.86% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, MG, PCA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.44	0/2974	0.87	10/4012 (0.2%)

There are no bond length outliers.

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	204	ALA	N-CA-C	-9.47	85.44	111.00
1	A	205	ALA	N-CA-C	9.18	135.79	111.00
1	A	203	ALA	N-CA-C	-8.06	89.23	111.00
1	A	194	GLY	N-CA-C	7.93	132.93	113.10
1	A	6	GLN	N-CA-C	-7.55	90.61	111.00
1	A	198	ALA	N-CA-C	7.38	130.91	111.00
1	A	195	HIS	N-CA-C	-5.91	95.06	111.00
1	A	331	ASP	N-CA-C	-5.33	96.61	111.00
1	A	199	ALA	N-CA-C	-5.33	96.62	111.00
1	A	7	GLU	N-CA-C	5.08	124.72	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2912	0	2889	56	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	1	0	0	0	0
3	A	4	0	3	1	0
4	A	9	0	5	2	0
5	A	256	0	0	16	0
All	All	3182	0	2897	57	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:83:VAL:HG21	1:A:110:PHE:CZ	1.86	1.08
1:A:89:LYS:HE2	1:A:91:ASN:HD21	1.45	0.79
1:A:83:VAL:HG21	1:A:110:PHE:CE2	2.23	0.74
1:A:155:VAL:HG11	1:A:170:GLU:OE2	1.86	0.74
1:A:83:VAL:CG2	1:A:110:PHE:CZ	2.70	0.73
1:A:327:VAL:HG21	5:A:504:HOH:O	1.95	0.66
1:A:83:VAL:HG13	5:A:618:HOH:O	1.95	0.66
1:A:287:ALA:HB2	1:A:300:HIS:H	1.61	0.66
1:A:162:PHE:O	1:A:164:VAL:HG23	1.96	0.65
1:A:202:ALA:CB	1:A:210:GLY:H	2.10	0.65
1:A:283:PRO:O	1:A:287:ALA:N	2.30	0.65
1:A:81:VAL:HG22	1:A:94:ASN:OD1	1.96	0.64
1:A:17:VAL:HG13	5:A:660:HOH:O	1.98	0.63
1:A:165:LYS:HE3	1:A:167:ASP:H	1.62	0.63
1:A:77:LYS:HG2	5:A:553:HOH:O	1.98	0.62
1:A:83:VAL:CG2	1:A:110:PHE:CE2	2.84	0.61
1:A:322:ILE:HG13	1:A:323:THR:N	2.16	0.60
1:A:234:GLN:NE2	1:A:343:HIS:H	2.00	0.58
1:A:202:ALA:O	1:A:203:ALA:HB2	2.05	0.56
1:A:89:LYS:CE	1:A:91:ASN:HD21	2.18	0.56
1:A:112:LEU:HD21	1:A:189:ALA:HA	1.89	0.55
1:A:154:VAL:HG22	1:A:157:ILE:HD12	1.87	0.55
1:A:328:ALA:HB3	5:A:627:HOH:O	2.07	0.54
1:A:320:HIS:NE2	5:A:501:HOH:O	2.00	0.54
1:A:270:ALA:HB1	5:A:748:HOH:O	2.08	0.53
1:A:267:ARG:HD2	3:A:402:ACT:O	2.08	0.53
1:A:16:VAL:HG12	5:A:556:HOH:O	2.09	0.52
1:A:202:ALA:HB2	1:A:210:GLY:H	1.75	0.51
1:A:188:GLN:HB3	1:A:192:LYS:NZ	2.25	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:403:PCA:HG3	5:A:730:HOH:O	2.11	0.51
1:A:83:VAL:HG21	1:A:110:PHE:CE1	2.44	0.50
1:A:83:VAL:HG11	5:A:620:HOH:O	2.11	0.49
1:A:154:VAL:HG23	1:A:157:ILE:H	1.78	0.49
1:A:81:VAL:HG23	5:A:736:HOH:O	2.12	0.48
1:A:77:LYS:NZ	5:A:506:HOH:O	2.46	0.48
1:A:234:GLN:HE22	1:A:343:HIS:H	1.62	0.47
1:A:53:VAL:HG11	5:A:733:HOH:O	2.15	0.47
1:A:192:LYS:HB2	1:A:192:LYS:HE2	1.74	0.46
1:A:146:PRO:HD3	1:A:179:GLY:C	2.37	0.45
1:A:165:LYS:CE	1:A:167:ASP:H	2.29	0.45
1:A:83:VAL:CG2	1:A:110:PHE:CE1	2.99	0.45
1:A:215:MET:N	1:A:215:MET:HE3	2.32	0.45
1:A:94:ASN:HB2	1:A:198:ALA:HA	2.00	0.44
1:A:202:ALA:O	1:A:203:ALA:CB	2.65	0.44
1:A:188:GLN:HB3	1:A:192:LYS:HZ2	1.81	0.44
1:A:67:CYS:SG	4:A:403:PCA:CD	3.06	0.43
1:A:121:LYS:HD3	5:A:733:HOH:O	2.18	0.43
1:A:165:LYS:HE3	1:A:166:ARG:HG2	2.00	0.43
1:A:215:MET:HE3	5:A:558:HOH:O	2.19	0.43
1:A:123:TYR:CZ	1:A:270:ALA:HB2	2.55	0.42
1:A:14:PHE:HB3	1:A:33:LEU:HD11	2.02	0.41
1:A:384:LEU:HD13	5:A:633:HOH:O	2.20	0.41
1:A:81:VAL:HG11	1:A:193:ALA:HA	2.02	0.41
1:A:7:GLU:HB2	1:A:13:LEU:HD23	2.03	0.41
1:A:61:PHE:CD2	1:A:192:LYS:HE3	2.55	0.41
1:A:154:VAL:CG2	1:A:157:ILE:HD12	2.50	0.41
1:A:237:VAL:HG23	1:A:240:ARG:HD2	2.03	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:191:ALA:O	1:A:194:GLY:O[17_555]	2.16	0.04

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	375/392 (96%)	358 (96%)	14 (4%)	3 (1%)	19 5

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	194	GLY
1	A	203	ALA
1	A	195	HIS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	296/302 (98%)	290 (98%)	6 (2%)	55 25

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	165	LYS
1	A	192	LYS
1	A	197	SER
1	A	267	ARG
1	A	322	ILE
1	A	342	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	91	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	ACT	A	402	-	3,3,3	3.10	2 (66%)	3,3,3	3.10	2 (66%)
4	PCA	A	403	-	9,9,9	2.45	3 (33%)	12,12,12	2.48	4 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PCA	A	403	-	-	0/4/13/13	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	403	PCA	CA-C	-5.04	1.39	1.52
3	A	402	ACT	O-C	4.61	1.43	1.22
4	A	403	PCA	CG-CD	-4.48	1.38	1.50
3	A	402	ACT	OXT-C	2.52	1.42	1.30
4	A	403	PCA	CA-N	2.13	1.48	1.45

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	403	PCA	CA-N-CD	-6.47	108.91	114.48
3	A	402	ACT	O-C-CH3	-3.80	107.55	122.33
3	A	402	ACT	OXT-C-O	-3.50	109.17	122.05
4	A	403	PCA	CB-CA-N	2.89	105.02	102.70
4	A	403	PCA	CG-CD-N	2.37	114.54	108.39
4	A	403	PCA	OE-CD-CG	-2.26	122.81	126.76

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	402	ACT	1	0
4	A	403	PCA	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	379/392 (96%)	0.79	41 (10%) 5 5	10, 18, 42, 67	14 (3%)

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	204	ALA	18.7
1	A	198	ALA	14.7
1	A	205	ALA	14.4
1	A	199	ALA	12.9
1	A	203	ALA	12.3
1	A	201	GLY	10.4
1	A	206	ALA	10.3
1	A	202	ALA	9.7
1	A	200	GLY	8.0
1	A	297	ALA	7.8
1	A	287	ALA	7.2
1	A	392	ALA	6.8
1	A	286	GLY	5.8
1	A	5	GLN	5.6
1	A	194	GLY	5.4
1	A	285	LEU	5.2
1	A	189	ALA	5.2
1	A	190	ALA	4.9
1	A	193	ALA	4.8
1	A	191	ALA	4.8
1	A	196	ALA	4.6
1	A	195	HIS	4.3
1	A	8	SER	3.9
1	A	6	GLN	3.5
1	A	123	TYR	3.5
1	A	186	GLY	3.4
1	A	192	LYS	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	328	ALA	3.3
1	A	197	SER	3.3
1	A	218	VAL	3.2
1	A	216	LYS	3.0
1	A	185	SER	2.8
1	A	208	VAL	2.6
1	A	117	ASN	2.5
1	A	284	HIS	2.4
1	A	187	ALA	2.4
1	A	214	LEU	2.4
1	A	184	ALA	2.3
1	A	298	VAL	2.3
1	A	207	ARG	2.1
1	A	329	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	PCA	A	403	9/9	0.60	0.27	39,45,49,49	9
3	ACT	A	402	4/4	0.90	0.16	12,13,14,15	0
2	MG	A	401	1/1	1.00	0.06	14,14,14,14	0

6.5 Other polymers [i](#)

There are no such residues in this entry.