

# Full wwPDB X-ray Structure Validation Report (i)

#### Jul 31, 2023 – 04:42 AM EDT

:	1CBW
:	BOVINE CHYMOTRYPSIN COMPLEXED TO BPTI
:	Hynes, T.R.; Scheidig, A.J.; Kossiakoff, A.A.
:	1996-12-22
:	2.60  Å(reported)
	: : : :

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
$\mathrm{EDS}$	:	2.34
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.34

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY \, DIFFRACTION$ 

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Motria	Whole archive	Similar resolution
Metric	$(\# { m Entries})$	$(\# { m Entries},  { m resolution}  { m range}({ m \AA}))$
R <sub>free</sub>	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain						
1	А	13	54%	31%	15%	-			
1	F	13	46%	38%	15%				
2	В	131	2% 66%		33% •				
2	G	131	<sup>2%</sup> 70%		28% •				
3	С	97	% 69%		29% •				



Mol	Chain	Length	Quality of chain		
3	Н	97	% 66%	29%	5%
4	D	58	69%	29%	•
4	Ι	58	74%	19%	5% •



#### $1\mathrm{CBW}$

## 2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 4588 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Δ	11	Total	С	Ν	Ο	$\mathbf{S}$	0	0	0
1	A	11	74	48	12	13	1	0		
1	Б	11	Total	С	Ν	0	S	0	0	0
1	Г	11	74	48	12	13	1	0		0

• Molecule 1 is a protein called BOVINE CHYMOTRYPSIN.

• Molecule 2 is a protein called BOVINE CHYMOTRYPSIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
9	В	121	Total	С	Ν	Ο	S	10	0	0
	D	101	980	618	162	196	4	19	0	0
9	С	121	Total	С	Ν	Ο	S	10	0	0
	2 G	131	980	618	162	196	4	19	0	0

• Molecule 3 is a protein called BOVINE CHYMOTRYPSIN.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
3	С	97	Total 702	C 436	N 123	O 136	S 7	5	0	0
3	Н	97	Total 702	C 436	N 123	O 136	S 7	23	0	0

• Molecule 4 is a protein called BPTI.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
4	л	59	Total	С	Ν	Ο	S	10	0	0
4	D	- 00	454	284	84	79	$\overline{7}$	10		
4	т	I 58	Total	С	Ν	Ο	S	10	0	0
4			454	284	84	79	7	10	0	0

• Molecule 5 is SULFATE ION (three-letter code: SO4) (formula:  $O_4S$ ).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	Ι	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
5	Ι	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0

• Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	А	5	$\begin{array}{cc} \text{Total} & \text{O} \\ 5 & 5 \end{array}$	0	0
6	В	31	Total         O           31         31	0	0
6	С	27	Total O 27 27	0	0
6	D	21	TotalO2121	0	0
6	F	3	Total O 3 3	0	0
6	G	30	Total O 30 30	0	0
6	Н	18	Total         O           18         18	0	0
6	Ι	23	Total O 23 23	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Chain A: 54% 31% 15% 07 V9 L1 • Molecule 1: BOVINE CHYMOTRYPSIN Chain F: 46% 38% 15% • Molecule 2: BOVINE CHYMOTRYPSIN Chain B: 66% 33% • Molecule 2: BOVINE CHYMOTRYPSIN Chain G: 70% 28% • Molecule 3: BOVINE CHYMOTRYPSIN Chain C: 69% 29%
- Molecule 1: BOVINE CHYMOTRYPSIN







# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 61	Depositor
Cell constants	101.60Å 101.60Å 205.90Å	Deperitor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $120.00^{\circ}$	Depositor
$\mathbf{P}_{\text{acclution}}(\hat{\mathbf{A}})$	10.00 - 2.60	Depositor
Resolution (A)	9.99 - 2.60	EDS
% Data completeness	89.0 (10.00-2.60)	Depositor
(in resolution range)	89.0 (9.99-2.60)	EDS
R <sub>merge</sub>	0.09	Depositor
R <sub>sym</sub>	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.01 (at 2.60Å)	Xtriage
Refinement program	X-PLOR 3.1	Depositor
D D.	0.192 , $0.251$	Depositor
$\Pi, \Pi_{free}$	0.185 , $0.241$	DCC
$R_{free}$ test set	3221 reflections $(9.99%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	42.0	Xtriage
Anisotropy	0.004	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.26 , $80.9$	EDS
L-test for twinning <sup>2</sup>	$< L >=0.50, < L^2>=0.33$	Xtriage
Estimated twinning fraction	0.043 for h,-h-k,-l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4588	wwPDB-VP
Average B, all atoms $(Å^2)$	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.25% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bo	ond angles
	Unain	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	А	0.69	0/75	0.95	0/103
1	F	0.65	0/75	1.07	0/103
2	В	0.70	0/1000	0.94	0/1361
2	G	0.69	0/1000	0.86	1/1361~(0.1%)
3	С	0.70	0/715	0.93	1/973~(0.1%)
3	Н	0.70	0/715	0.90	0/973
4	D	0.81	0/465	0.91	1/622~(0.2%)
4	Ι	0.80	0/465	0.99	1/622~(0.2%)
All	All	0.72	0/4510	0.92	4/6118 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms		$Observed(^{o})$	$Ideal(^{o})$
3	С	199	LEU	N-CA-C	-6.30	94.00	111.00
4	Ι	29	LEU	CA-CB-CG	5.32	127.54	115.30
4	D	17	ARG	NE-CZ-NH2	-5.16	117.72	120.30
2	G	143	LEU	CB-CG-CD1	-5.13	102.27	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	74	0	81	9	0
1	F	74	0	81	6	0
2	В	980	0	951	37	0
2	G	980	0	951	23	0
3	С	702	0	698	19	0
3	Н	702	0	698	17	0
4	D	454	0	438	18	0
4	Ι	454	0	438	17	0
5	Ι	10	0	0	0	0
6	А	5	0	0	0	0
6	В	31	0	0	2	0
6	С	27	0	0	1	0
6	D	21	0	0	2	0
6	F	3	0	0	0	0
6	G	30	0	0	0	0
6	Н	18	0	0	0	0
6	Ι	23	0	0	1	0
All	All	4588	0	4336	120	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (120) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
4:I:1:ARG:HA	4:I:58:ALA:HB1	1.53	0.89
2:B:28:PRO:HB2	2:B:119:SER:H	1.48	0.78
2:G:16:ILE:O	2:G:144:THR:HA	1.83	0.78
1:F:9:VAL:HG23	2:G:23:VAL:HG21	1.70	0.73
3:H:164:SER:OG	3:H:167:ASN:HB2	1.91	0.70
6:B:161:HOH:O	4:I:52:MET:HG3	1.94	0.67
1:A:8:PRO:HG3	2:B:26:SER:HB2	1.77	0.65
2:G:56:ALA:HA	2:G:104:THR:OG1	1.96	0.65
1:A:8:PRO:HG3	2:B:26:SER:O	1.96	0.64
3:C:213:VAL:HG22	3:C:228:TYR:HE2	1.62	0.63
4:D:21:TYR:CE2	4:D:48:ALA:HB2	2.34	0.63
2:G:52:VAL:HB	2:G:106:LEU:HD12	1.82	0.61
2:B:28:PRO:CG	2:B:119:SER:HB3	2.31	0.61
3:H:166:THR:HG22	3:H:170:LYS:NZ	2.16	0.60
1:A:5:ALA:CB	2:B:116:GLN:HB2	2.31	0.60
3:H:165:ASN:O	3:H:169:LYS:HG3	2.03	0.59
4:I:1:ARG:HA	4:I:58:ALA:CB	2.28	0.59



		Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
3:C:213:VAL:HG22	3:C:228:TYR:CE2	2.38	0.59
2:G:46:LEU:O	2:G:121:VAL:HG12	2.04	0.58
1:F:9:VAL:CG2	2:G:23:VAL:HG21	2.33	0.58
3:C:230:ARG:O	3:C:234:LEU:HD22	2.04	0.58
4:D:12:GLY:HA3	6:D:59:HOH:O	2.03	0.57
4:D:29:LEU:CD2	4:I:19:ILE:HD12	2.34	0.57
1:F:4:PRO:O	1:F:7:GLN:NE2	2.38	0.57
2:G:28:PRO:HB2	2:G:119:SER:H	1.69	0.57
4:D:25:ALA:HA	6:D:73:HOH:O	2.03	0.56
3:H:202:LYS:HA	3:H:206:ALA:O	2.05	0.56
4:I:18:ILE:HD12	4:I:18:ILE:N	2.20	0.56
3:H:169:LYS:O	3:H:173:GLY:HA2	2.05	0.56
4:D:53:ARG:HG2	4:D:53:ARG:HH11	1.71	0.55
4:D:53:ARG:HG2	4:D:53:ARG:NH1	2.22	0.55
3:H:235:VAL:HG13	3:H:236:ASN:N	2.22	0.54
2:B:88:VAL:HG22	2:B:106:LEU:CD2	2.37	0.53
2:B:57:HIS:HD2	4:D:37:GLY:HA3	1.74	0.53
1:A:8:PRO:CA	2:B:26:SER:HB2	2.38	0.53
2:B:81:GLN:HE21	2:B:118:VAL:HG21	1.73	0.53
3:C:192:MET:HE3	3:C:192:MET:H	1.74	0.53
2:B:124:PRO:HD3	3:C:209:LEU:O	2.09	0.52
2:B:44:GLY:HA2	3:C:196:GLY:O	2.10	0.52
3:C:213:VAL:HA	3:C:228:TYR:CD2	2.44	0.52
4:D:29:LEU:HD23	4:I:19:ILE:CD1	2.41	0.51
2:B:114:PHE:HA	2:B:118:VAL:O	2.10	0.51
4:I:10:TYR:HB2	4:I:41:LYS:HD2	1.92	0.51
4:D:29:LEU:HD22	4:I:19:ILE:HD12	1.92	0.51
4:D:49:GLU:O	4:D:53:ARG:HG3	2.11	0.51
3:H:238:VAL:O	3:H:242:LEU:HG	2.11	0.51
2:G:27:TRP:HE3	2:G:29:TRP:CZ2	2.29	0.50
2:G:94:TYR:CD1	2:G:102:ASP:HA	2.47	0.50
1:A:8:PRO:HA	2:B:26:SER:HB2	1.94	0.50
2:B:60:VAL:HG12	2:B:61:THR:N	2.26	0.49
4:I:23:TYR:CE2	4:I:25:ALA:HA	2.48	0.49
3:H:195:SER:OG	4:I:15:LYS:C	2.51	0.49
1:A:8:PRO:HA	2:B:26:SER:CB	2.42	0.49
2:B:28:PRO:HG2	2:B:119:SER:HB3	1.94	0.49
3:C:195:SER:OG	4:D:15:LYS:C	2.50	0.49
2:B:17:VAL:O	3:C:188:VAL:HA	2.13	0.48
2:B:94:TYR:HA	2:B:100:ASN:O	2.13	0.48
2:B:81:GLN:NE2	2:B:118:VAL:HG21	2.28	0.48



		Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
2:G:114:PHE:HA	2:G:118:VAL:O	2.13	0.48
1:F:5:ALA:CB	2:G:116:GLN:HA	2.43	0.48
3:C:201:CYS:SG	3:C:210:VAL:HG21	2.53	0.48
3:H:216:GLY:O	3:H:217:SER:C	2.51	0.48
4:D:23:TYR:CE2	4:D:25:ALA:HA	2.49	0.47
1:F:9:VAL:O	1:F:10:LEU:HB3	2.15	0.47
3:H:192:MET:HE3	3:H:220:CYS:SG	2.54	0.47
2:B:64:ASP:O	2:B:85:ILE:HD12	2.14	0.47
2:G:53:VAL:HG23	2:G:105:LEU:HD23	1.97	0.47
3:H:184:GLY:C	3:H:186:SER:N	2.68	0.47
2:B:141:TRP:O	3:C:152:PRO:HD2	2.15	0.47
3:H:234:LEU:O	3:H:238:VAL:HG23	2.15	0.47
2:B:121:VAL:HG22	2:B:122:CYS:N	2.30	0.46
4:I:44:ASN:OD1	4:I:44:ASN:C	2.53	0.46
2:B:17:VAL:O	2:B:18:ASN:HB2	2.16	0.46
4:I:10:TYR:CB	4:I:41:LYS:HD2	2.45	0.46
3:C:211:GLY:HA2	3:C:229:ALA:O	2.15	0.46
4:D:31:GLN:HG2	6:I:402:HOH:O	2.15	0.46
2:B:86:ALA:HB2	2:B:109:SER:HA	1.97	0.46
4:D:29:LEU:HD23	4:I:19:ILE:HD12	1.98	0.45
1:A:8:PRO:CG	2:B:26:SER:HB2	2.43	0.45
3:C:211:GLY:HA2	3:C:231:VAL:HG23	1.99	0.45
2:B:112:ALA:HB3	2:B:114:PHE:HE1	1.82	0.45
2:G:75:SER:OG	2:G:78:GLU:HG3	2.17	0.45
2:G:126:ALA:HB2	3:H:236:ASN:OD1	2.16	0.45
3:C:232:THR:HG23	6:C:38:HOH:O	2.16	0.45
4:D:21:TYR:HA	4:D:31:GLN:O	2.16	0.44
4:D:19:ILE:HG12	4:I:29:LEU:HD23	1.99	0.44
2:B:123:LEU:HA	2:B:123:LEU:HD23	1.79	0.43
2:B:23:VAL:HB	6:B:152:HOH:O	2.19	0.43
2:B:48:ASN:OD1	2:B:48:ASN:C	2.57	0.43
2:B:56:ALA:HA	2:B:104:THR:OG1	2.19	0.43
4:I:53:ARG:HH11	4:I:53:ARG:HB3	1.82	0.43
2:B:48:ASN:OD1	2:B:50:ASN:N	2.44	0.43
2:G:60:VAL:HG12	2:G:61:THR:N	2.32	0.43
3:C:162:LEU:C	3:C:163:LEU:HD23	2.39	0.43
2:B:98:THR:C	2:B:99:ILE:HG13	2.38	0.42
4:D:7:GLU:HA	4:D:8:PRO:HD2	1.85	0.42
2:G:121:VAL:HG22	2:G:122:CYS:N	2.34	0.42
2:G:136:CYS:HB3	3:H:200:VAL:O	2.19	0.42
3:C:203:LYS:O	3:C:204:ASN:HB2	2.19	0.42



A 4 am 1	A.t.a.m. D	Interatomic	Clash
Atom-1	Atom-2	distance $(\text{\AA})$	overlap (Å)
2:B:108:LEU:HA	2:B:108:LEU:HD23	1.72	0.42
1:A:9:VAL:HG22	2:B:23:VAL:HG23	2.01	0.42
3:C:209:LEU:HG	3:C:231:VAL:HG21	2.01	0.42
4:D:4:PHE:CD1	4:D:5:CYS:N	2.88	0.42
4:I:1:ARG:N	4:I:2:PRO:CD	2.82	0.42
2:G:45:SER:OG	3:H:198:PRO:HB3	2.20	0.41
2:G:100:ASN:HD22	2:G:101:ASN:ND2	2.18	0.41
2:B:65:VAL:HG13	2:B:83:LEU:O	2.20	0.41
2:G:125:SER:O	2:G:128:ASP:HB2	2.20	0.41
2:G:27:TRP:CE3	2:G:29:TRP:CZ2	3.08	0.41
3:C:232:THR:O	3:C:235:VAL:HG12	2.20	0.41
4:I:53:ARG:HH11	4:I:53:ARG:CB	2.33	0.41
2:B:121:VAL:HG22	2:B:122:CYS:H	1.86	0.41
2:G:24:PRO:HG3	2:G:71:PHE:CZ	2.56	0.41
2:G:89:PHE:HE1	2:G:107:LYS:HB2	1.86	0.41
3:H:160:LEU:HD23	3:H:160:LEU:H	1.85	0.40
2:B:60:VAL:CG1	2:B:61:THR:N	2.84	0.40
3:C:184:GLY:O	3:C:185:ALA:HB3	2.21	0.40
1:F:10:LEU:C	1:F:10:LEU:HD12	2.42	0.40
3:H:171:TYR:HE1	3:H:172:TRP:CZ2	2.40	0.40
1:A:7:GLN:HA	1:A:8:PRO:HD3	1.91	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	9/13~(69%)	9 (100%)	0	0	100 100
1	F	9/13~(69%)	$8 \ (89\%)$	1 (11%)	0	100 100
2	В	129/131~(98%)	116 (90%)	11 (8%)	2(2%)	9 19
2	G	129/131~(98%)	120 (93%)	7 (5%)	2(2%)	9 19



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
3	С	95/97~(98%)	90~(95%)	4 (4%)	1 (1%)	14	30
3	Н	95/97~(98%)	83~(87%)	10 (10%)	2(2%)	7	13
4	D	56/58~(97%)	52 (93%)	4 (7%)	0	100	100
4	Ι	56/58~(97%)	51 (91%)	5 (9%)	0	100	100
All	All	578/598~(97%)	529~(92%)	42 (7%)	7(1%)	13	27

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	Н	244	ALA
3	С	198	PRO
2	G	71	PHE
3	Н	198	PRO
2	В	143	LEU
2	В	99	ILE
2	G	99	ILE

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	9/10~(90%)	9 (100%)	0	100 100
1	F	9/10~(90%)	9~(100%)	0	100 100
2	В	109/109~(100%)	103~(94%)	6~(6%)	21 43
2	G	109/109~(100%)	107~(98%)	2(2%)	59 80
3	С	77/77~(100%)	70~(91%)	7 (9%)	9 18
3	Н	77/77~(100%)	67~(87%)	10 (13%)	4 7
4	D	46/46~(100%)	44 (96%)	2(4%)	29 54
4	Ι	46/46~(100%)	41 (89%)	5 (11%)	6 11
All	All	482/484~(100%)	450 (93%)	32(7%)	16 33

All (32) residues with a non-rotameric sidechain are listed below:



Mol	Chain	$\mathbf{Res}$	Type
2	В	20	GLU
2	В	32	SER
2	В	45	SER
2	В	77	SER
2	В	127	SER
2	В	144	THR
3	С	153	ASP
3	С	160	LEU
3	С	192	MET
3	С	198	PRO
3	С	217	SER
3	С	236	ASN
3	С	239	GLN
4	D	1	ARG
4	D	29	LEU
2	G	32	SER
2	G	75	SER
3	Н	153	ASP
3	Н	159	SER
3	Н	160	LEU
3	Н	162	LEU
3	Н	167	ASN
3	Н	177	LYS
3	Н	192	MET
3	Н	217	SER
3	Н	239	GLN
3	Н	240	GLN
4	Ι	1	ARG
4	Ι	3	ASP
4	Ι	29	LEU
4	Ι	44	ASN
4	Ι	53	ARG

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such side chains are listed below:

Mol	Chain	Res	Type
2	В	34	GLN
2	В	73	GLN
2	В	101	ASN
3	С	167	ASN
1	F	7	GLN
2	G	101	ASN
3	Н	240	GLN



#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type	Turne	Chain	Res	Tink	Bond lengths			Bond angles		
	туре	Chan			Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	SO4	Ι	301	-	4,4,4	0.62	0	6,6,6	0.64	0
5	SO4	Ι	401	-	4,4,4	0.60	0	6,6,6	0.50	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers (i)

There are no such residues in this entry.



## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	< <b>RSRZ</b> >	#RSRZ>2	$OWAB(Å^2)$	Q < 0.9
1	А	11/13 (84%)	-0.65	0 100 100	26, 42, 66, 68	0
1	F	11/13 (84%)	-0.61	0 100 100	37, 51, 63, 69	0
2	В	131/131 (100%)	-1.10	3 (2%) 60 54	16, 35, 58, 70	5~(3%)
2	G	131/131 (100%)	-1.03	2 (1%) 73 70	16,  36,  62,  70	5 (3%)
3	С	97/97~(100%)	-1.07	1 (1%) 82 80	12, 31, 63, 68	1 (1%)
3	Η	95/97~(97%)	-1.00	1 (1%) 80 78	14,  35,  62,  70	2(2%)
4	D	58/58~(100%)	-1.25	0 100 100	11, 24, 48, 62	2 (3%)
4	Ι	58/58~(100%)	-1.27	0 100 100	10, 23, 44, 61	2 (3%)
All	All	592/598~(98%)	-1.08	7 (1%) 79 76	10, 32, 62, 70	17 (2%)

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	Н	243	ALA	3.1
2	G	76	SER	3.0
2	В	77	SER	2.6
2	В	75	SER	2.5
2	В	76	SER	2.3
2	G	77	SER	2.2
3	С	243	ALA	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.



## 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathrm{\AA}^2)$	Q<0.9
5	SO4	Ι	301	5/5	0.95	0.12	45,46,59,62	0
5	SO4	Ι	401	5/5	0.99	0.16	43,49,59,62	0

### 6.5 Other polymers (i)

There are no such residues in this entry.

