



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 22, 2023 – 06:04 AM EDT

PDB ID : 3CDG
Title : Human CD94/NKG2A in complex with HLA-E
Authors : Petrie, E.J.; Clements, C.S.; Lin, J.; Sullivan, L.C.; Johnson, D.; Huyton, T.; Heroux, A.; Hoare, H.L.; Beddoe, T.; Reid, H.H.; Wilce, M.C.J.; Brooks, A.G.; Rossjohn, J.
Deposited on : 2008-02-26
Resolution : 3.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1026 (3.48-3.32)
Clashscore	141614	1055 (3.48-3.32)
Ramachandran outliers	138981	1038 (3.48-3.32)
Sidechain outliers	138945	1038 (3.48-3.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	273	56% (green), 38% (yellow), 6% (orange)
1	C	273	58% (green), 36% (yellow), 5% (orange)
2	B	100	59% (green), 36% (yellow), 5% (orange)
2	D	100	62% (green), 33% (yellow), 5% (orange)
3	E	123	52% (green), 39% (yellow), 8% (orange), . (red)
3	J	123	52% (green), 38% (yellow), 9% (orange), . (red)
4	F	120	38% (green), 52% (yellow), 8% (orange), . (red)

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	K	120	 40% 52% . .
5	P	9	 22% 56% 22%
5	Q	9	 22% 67% 11%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 10200 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA class I histocompatibility antigen, alpha chain E.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	273	Total	C	N	O	S	0	0	0
			2234	1396	401	430	7			
1	C	273	Total	C	N	O	S	0	0	0
			2234	1396	401	430	7			

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	100	Total	C	N	O	S	0	0	0
			837	533	141	159	4			
2	D	100	Total	C	N	O	S	0	0	0
			837	533	141	159	4			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	expression tag	UNP P61769
D	0	MET	-	expression tag	UNP P61769

- Molecule 3 is a protein called Natural killer cells antigen CD94.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	J	123	Total	C	N	O	S	0	0	0
			1010	632	167	201	10			
3	E	123	Total	C	N	O	S	0	0	0
			1010	632	167	201	10			

- Molecule 4 is a protein called NKG2-A/NKG2-B type II integral membrane protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	K	116	Total	C	N	O	S	0	0	0
			931	585	166	171	9			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	F	120	961	600	171	181	9	0	0	0


- Molecule 5 is a protein called leader peptide of HLA class I histocompatibility antigen, alpha chain G.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
5	P	9	73	49	12	11	1	0	0	0
5	Q	9	73	49	12	11	1	0	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HLA class I histocompatibility antigen, alpha chain E




- Molecule 1: HLA class I histocompatibility antigen, alpha chain E




- Molecule 2: Beta-2-microglobulin




- Molecule 2: Beta-2-microglobulin




- Molecule 3: Natural killer cells antigen CD94




- Molecule 3: Natural killer cells antigen CD94




- Molecule 4: NKG2-A/NKG2-B type II integral membrane protein




- Molecule 4: NKG2-A/NKG2-B type II integral membrane protein



- Molecule 5: leader peptide of HLA class I histocompatibility antigen, alpha chain G

Chain P:  22% 56% 22%



- Molecule 5: leader peptide of HLA class I histocompatibility antigen, alpha chain G

Chain Q:  22% 67% 11%



4 Data and refinement statistics

Property	Value	Source
Space group	I 41 3 2	Depositor
Cell constants a, b, c, α , β , γ	345.20Å 345.20Å 345.20Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 3.40 140.93 – 3.40	Depositor EDS
% Data completeness (in resolution range)	99.9 (50.00-3.40) 99.9 (140.93-3.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.58 (at 3.41Å)	Xtrriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.249 , 0.279 0.302 , 0.330	Depositor DCC
R_{free} test set	2434 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	100.7	Xtrriage
Anisotropy	0.000	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 107.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.26$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.84	EDS
Total number of atoms	10200	wwPDB-VP
Average B, all atoms (Å ²)	118.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.34% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.36	0/2300	0.52	1/3127 (0.0%)
1	C	0.36	0/2300	0.53	1/3127 (0.0%)
2	B	0.36	0/860	0.47	0/1162
2	D	0.35	0/860	0.46	0/1162
3	E	0.36	0/1036	0.51	1/1402 (0.1%)
3	J	0.37	0/1036	0.51	1/1402 (0.1%)
4	F	0.30	0/985	0.48	0/1330
4	K	0.30	0/954	0.47	0/1286
5	P	0.30	0/74	0.62	0/98
5	Q	0.31	0/74	0.59	0/98
All	All	0.35	0/10479	0.51	4/14194 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
4	F	3	1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	227	ASP	CB-CG-OD2	5.24	123.02	118.30
1	C	227	ASP	CB-CG-OD2	5.22	123.00	118.30
3	J	57	ASP	CB-CG-OD2	5.20	122.98	118.30
3	E	57	ASP	CB-CG-OD2	5.20	122.97	118.30

All (3) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	F	199	LYS	CA
4	F	200	ASP	CA
4	F	203	ASN	CA

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
4	F	205	GLU	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2234	0	2072	148	0
1	C	2234	0	2072	135	0
2	B	837	0	803	36	0
2	D	837	0	803	36	0
3	E	1010	0	913	54	0
3	J	1010	0	913	63	0
4	F	961	0	915	81	0
4	K	931	0	895	58	0
5	P	73	0	83	11	0
5	Q	73	0	83	8	0
All	All	10200	0	9552	554	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 28.

The worst 5 of 554 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:221:GLY:O	1:A:222:GLU:HG3	1.07	1.23
1:C:221:GLY:O	1:C:222:GLU:HG3	1.08	1.22
1:A:219:GLN:HE21	1:A:225:THR:CG2	1.58	1.16
1:A:219:GLN:HE21	1:A:225:THR:HG22	1.14	1.13
1:C:219:GLN:HE21	1:C:225:THR:CG2	1.61	1.13

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	271/273 (99%)	225 (83%)	35 (13%)	11 (4%)	3	18
1	C	271/273 (99%)	226 (83%)	35 (13%)	10 (4%)	3	20
2	B	98/100 (98%)	90 (92%)	7 (7%)	1 (1%)	15	46
2	D	98/100 (98%)	89 (91%)	9 (9%)	0	100	100
3	E	121/123 (98%)	101 (84%)	16 (13%)	4 (3%)	4	22
3	J	121/123 (98%)	101 (84%)	16 (13%)	4 (3%)	4	22
4	F	118/120 (98%)	91 (77%)	21 (18%)	6 (5%)	2	14
4	K	112/120 (93%)	91 (81%)	17 (15%)	4 (4%)	3	21
5	P	7/9 (78%)	4 (57%)	3 (43%)	0	100	100
5	Q	7/9 (78%)	5 (71%)	2 (29%)	0	100	100
All	All	1224/1250 (98%)	1023 (84%)	161 (13%)	40 (3%)	4	22

5 of 40 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	213	ILE
1	A	215	LEU
1	A	222	GLU
1	A	224	HIS
3	J	62	GLN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	236/236 (100%)	214 (91%)	22 (9%)	9	31
1	C	236/236 (100%)	216 (92%)	20 (8%)	10	35
2	B	95/95 (100%)	87 (92%)	8 (8%)	11	36
2	D	95/95 (100%)	87 (92%)	8 (8%)	11	36
3	E	114/114 (100%)	102 (90%)	12 (10%)	7	25
3	J	114/114 (100%)	101 (89%)	13 (11%)	5	21
4	F	109/109 (100%)	99 (91%)	10 (9%)	9	31
4	K	105/109 (96%)	98 (93%)	7 (7%)	16	46
5	P	8/8 (100%)	6 (75%)	2 (25%)	0	2
5	Q	8/8 (100%)	6 (75%)	2 (25%)	0	2
All	All	1120/1124 (100%)	1016 (91%)	104 (9%)	9	31

5 of 104 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	88	SER
2	D	58	LYS
4	F	214	ASN
1	C	134	THR
1	C	196	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 57 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	32	GLN
4	F	228	HIS
1	C	197	HIS
4	F	214	ASN
3	E	177	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.