



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 13, 2023 – 03:18 pm GMT

PDB ID : 8CE9
Title : Crystal structure of human Cd11b I domain in C121 space group
Authors : Koekemoer, L.; Williams, E.; Ni, X.; Gao, Q.; Coker, J.; MacLean, E.M.;
Wright, N.D.; Marsden, B.; Von Delft, F.; Schwenzer, A.; Midwood, K.
Deposited on : 2023-02-01
Resolution : 2.11 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.32.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.32.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.11 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6241 (2.14-2.10)
Clashscore	141614	6778 (2.14-2.10)
Ramachandran outliers	138981	6705 (2.14-2.10)
Sidechain outliers	138945	6706 (2.14-2.10)
RSRZ outliers	127900	6112 (2.14-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	215	 75% 13% 12%
1	B	215	 79% 9% 12%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	EDO	B	209	-	-	X	-

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 6337 atoms, of which 3130 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Integrin alpha-M.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	B	190	2990	958	1493	256	280	3	47	3	0
1	A	190	3039	969	1517	265	285	3	44	4	0

There are 52 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-21	MET	-	initiating methionine	UNP P11215
B	-20	HIS	-	expression tag	UNP P11215
B	-19	HIS	-	expression tag	UNP P11215
B	-18	HIS	-	expression tag	UNP P11215
B	-17	HIS	-	expression tag	UNP P11215
B	-16	HIS	-	expression tag	UNP P11215
B	-15	HIS	-	expression tag	UNP P11215
B	-14	SER	-	expression tag	UNP P11215
B	-13	SER	-	expression tag	UNP P11215
B	-12	GLY	-	expression tag	UNP P11215
B	-11	VAL	-	expression tag	UNP P11215
B	-10	ASP	-	expression tag	UNP P11215
B	-9	LEU	-	expression tag	UNP P11215
B	-8	GLY	-	expression tag	UNP P11215
B	-7	THR	-	expression tag	UNP P11215
B	-6	GLU	-	expression tag	UNP P11215
B	-5	ASN	-	expression tag	UNP P11215
B	-4	LEU	-	expression tag	UNP P11215
B	-3	TYR	-	expression tag	UNP P11215
B	-2	PHE	-	expression tag	UNP P11215
B	-1	GLN	-	expression tag	UNP P11215
B	0	SER	-	expression tag	UNP P11215
B	190	GLU	-	expression tag	UNP P11215
B	191	PRO	-	expression tag	UNP P11215
B	192	GLU	-	expression tag	UNP P11215

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	193	ALA	-	expression tag	UNP P11215
A	-21	MET	-	initiating methionine	UNP P11215
A	-20	HIS	-	expression tag	UNP P11215
A	-19	HIS	-	expression tag	UNP P11215
A	-18	HIS	-	expression tag	UNP P11215
A	-17	HIS	-	expression tag	UNP P11215
A	-16	HIS	-	expression tag	UNP P11215
A	-15	HIS	-	expression tag	UNP P11215
A	-14	SER	-	expression tag	UNP P11215
A	-13	SER	-	expression tag	UNP P11215
A	-12	GLY	-	expression tag	UNP P11215
A	-11	VAL	-	expression tag	UNP P11215
A	-10	ASP	-	expression tag	UNP P11215
A	-9	LEU	-	expression tag	UNP P11215
A	-8	GLY	-	expression tag	UNP P11215
A	-7	THR	-	expression tag	UNP P11215
A	-6	GLU	-	expression tag	UNP P11215
A	-5	ASN	-	expression tag	UNP P11215
A	-4	LEU	-	expression tag	UNP P11215
A	-3	TYR	-	expression tag	UNP P11215
A	-2	PHE	-	expression tag	UNP P11215
A	-1	GLN	-	expression tag	UNP P11215
A	0	SER	-	expression tag	UNP P11215
A	190	GLU	-	expression tag	UNP P11215
A	191	PRO	-	expression tag	UNP P11215
A	192	GLU	-	expression tag	UNP P11215
A	193	ALA	-	expression tag	UNP P11215

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	H	O	1	0
			10	2	6	2		
3	B	1	Total	C	H	O	1	0
			10	2	6	2		
3	B	1	Total	C	H	O	1	0
			10	2	6	2		
3	B	1	Total	C	H	O	1	0
			10	2	6	2		
3	B	1	Total	C	H	O	1	0
			10	2	6	2		
3	B	1	Total	C	H	O	1	0
			10	2	6	2		
3	B	1	Total	C	H	O	1	0
			10	2	6	2		
3	B	1	Total	C	H	O	1	0
			10	2	6	2		
3	B	1	Total	C	H	O	1	0
			10	2	6	2		
3	A	1	Total	C	H	O	1	0
			10	2	6	2		
3	A	1	Total	C	H	O	1	0
			10	2	6	2		
3	A	1	Total	C	H	O	1	0
			10	2	6	2		
3	A	1	Total	C	H	O	1	0
			10	2	6	2		
3	A	1	Total	C	H	O	1	0
			10	2	6	2		
3	A	1	Total	C	H	O	1	0
			10	2	6	2		
3	A	1	Total	C	H	O	1	0
			10	2	6	2		


- Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	41	Total O 41 41	0	0
4	A	42	Total O 42 42	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Integrin alpha-M

Chain B: 



- Molecule 1: Integrin alpha-M

Chain A: 



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	128.50Å 51.49Å 90.17Å 90.00° 132.96° 90.00°	Depositor
Resolution (Å)	65.99 – 2.11 65.99 – 2.11	Depositor EDS
% Data completeness (in resolution range)	92.5 (65.99-2.11) 92.5 (65.99-2.11)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.69 (at 2.10Å)	Xtrriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.188 , 0.239 0.195 , 0.241	Depositor DCC
R_{free} test set	1085 reflections (4.68%)	wwPDB-VP
Wilson B-factor (Å ²)	31.4	Xtrriage
Anisotropy	0.079	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.43 , 55.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.029 for h+2*k,-h-l 0.016 for h,-k,-h-l 0.023 for -h-2*k,-k,l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6337	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.76	0/1554	0.86	0/2100
1	B	0.74	0/1532	0.85	0/2072
All	All	0.75	0/3086	0.85	0/4172

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1522	1517	1494	23	0
1	B	1497	1493	1467	12	0
2	A	15	0	0	0	0
2	B	10	0	0	0	0
3	A	40	60	60	6	0
3	B	40	60	60	6	0
4	A	42	0	0	3	0
4	B	41	0	0	0	0
All	All	3207	3130	3081	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:167:VAL:HG13	1:A:172:ALA:HB3	1.77	0.66
1:B:175:THR:O	3:B:209:EDO:H22	1.95	0.66
1:A:92:ASN:OD1	1:A:94[A]:THR:HG22	2.00	0.61
1:A:12:SER:O	3:A:204:EDO:H22	1.99	0.61
1:A:133:ILE:HD13	1:A:187:ILE:HD13	1.83	0.59
1:A:178:ASN:OD1	1:A:181:ARG:NH2	2.35	0.59
1:A:71:THR:HG23	4:A:301:HOH:O	2.02	0.57
3:B:209:EDO:C1	1:A:187:ILE:HA	2.34	0.57
1:A:101:ALA:O	3:A:212:EDO:H21	2.05	0.56
1:A:130:GLU:O	3:A:212:EDO:H11	2.06	0.56
1:A:173:LEU:HB2	3:A:213:EDO:H11	1.89	0.54
3:B:209:EDO:H12	1:A:187:ILE:HA	1.90	0.54
1:B:91:PHE:CD1	1:B:103:LYS:HD3	2.46	0.51
1:B:17:ASP:HB3	1:B:170:PHE:CD2	2.45	0.51
1:A:171:GLU:HG2	4:A:331:HOH:O	2.12	0.48
1:B:178:ASN:HB3	3:B:209:EDO:O2	2.14	0.48
1:A:31:GLN:HB3	1:A:181:ARG:HD3	1.96	0.47
1:A:184:ILE:O	1:A:188:GLU:HG3	2.14	0.47
1:A:93:ILE:HD11	1:A:99:LYS:HG3	1.98	0.46
1:B:52:PHE:HA	1:B:56:GLU:OE1	2.16	0.45
1:B:68:LYS:N	1:B:69:PRO:CD	2.80	0.45
1:A:136:VAL:HG22	1:A:155:ILE:HG22	1.99	0.45
1:B:140:GLY:HA2	1:B:168:ASN:O	2.17	0.44
3:B:209:EDO:H11	1:A:187:ILE:HA	1.99	0.44
1:A:112:GLU:HA	1:A:151:GLU:OE1	2.17	0.44
1:B:31:GLN:HB3	1:B:181:ARG:HD3	1.99	0.44
1:B:176:ILE:C	3:B:209:EDO:H21	2.38	0.44
1:B:68:LYS:N	1:B:69:PRO:HD2	2.33	0.44
1:A:68:LYS:N	1:A:69:PRO:CD	2.81	0.43
1:B:167:VAL:HG13	1:B:172:ALA:HB3	1.99	0.43
1:A:174:LYS:N	3:A:213:EDO:H12	2.34	0.42
1:B:28:VAL:HG12	1:B:32:LEU:HD12	2.02	0.41
1:A:171:GLU:CG	4:A:331:HOH:O	2.67	0.41
1:A:103:LYS:HG2	3:A:212:EDO:H21	2.02	0.40
1:A:68:LYS:N	1:A:69:PRO:HD2	2.37	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	192/215 (89%)	188 (98%)	4 (2%)	0	100	100
1	B	191/215 (89%)	184 (96%)	6 (3%)	1 (0%)	29	25
All	All	383/430 (89%)	372 (97%)	10 (3%)	1 (0%)	41	40

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	99	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	163/191 (85%)	160 (98%)	3 (2%)	59	63
1	B	159/191 (83%)	159 (100%)	0	100	100
All	All	322/382 (84%)	319 (99%)	3 (1%)	86	83

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	150[A]	GLN
1	A	150[B]	GLN
1	A	171	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such

sidechains are listed below:

Mol	Chain	Res	Type
1	B	179	GLN
1	A	177	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

25 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	B	201	-	4,4,4	0.50	0	6,6,6	0.25	0
2	SO4	A	205	-	4,4,4	0.33	0	6,6,6	0.19	0
3	EDO	B	203	-	3,3,3	0.20	0	2,2,2	0.08	0
3	EDO	B	209	-	3,3,3	0.22	0	2,2,2	0.80	0
3	EDO	A	212	-	3,3,3	0.32	0	2,2,2	0.60	0
3	EDO	A	207	-	3,3,3	0.18	0	2,2,2	0.17	0
3	EDO	B	208	-	3,3,3	0.18	0	2,2,2	0.29	0
2	SO4	A	206	-	4,4,4	0.29	0	6,6,6	0.11	0
3	EDO	A	203	-	3,3,3	0.33	0	2,2,2	0.39	0
3	EDO	B	202	-	3,3,3	0.23	0	2,2,2	0.29	0
3	EDO	B	204	-	3,3,3	0.32	0	2,2,2	0.37	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	EDO	A	204	-	3,3,3	0.25	0	2,2,2	0.41	0
3	EDO	A	209	-	3,3,3	0.59	0	2,2,2	0.63	0
3	EDO	B	206	-	3,3,3	0.25	0	2,2,2	0.16	0
3	EDO	B	205	-	3,3,3	0.17	0	2,2,2	0.28	0
2	SO4	B	207	-	4,4,4	0.24	0	6,6,6	0.22	0
3	EDO	B	211	-	3,3,3	0.57	0	2,2,2	0.75	0
3	EDO	A	211	-	3,3,3	0.25	0	2,2,2	0.40	0
3	EDO	A	213	-	3,3,3	0.23	0	2,2,2	0.35	0
3	EDO	A	210	-	3,3,3	0.22	0	2,2,2	0.52	0
3	EDO	A	208	-	3,3,3	0.40	0	2,2,2	0.33	0
3	EDO	B	212	-	3,3,3	0.07	0	2,2,2	0.10	0
2	SO4	A	202	-	4,4,4	0.55	0	6,6,6	0.16	0
3	EDO	A	201	-	3,3,3	0.21	0	2,2,2	0.10	0
3	EDO	B	210	-	3,3,3	0.21	0	2,2,2	0.24	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	B	203	-	-	1/1/1/1	-
3	EDO	B	209	-	-	1/1/1/1	-
3	EDO	A	212	-	-	1/1/1/1	-
3	EDO	A	207	-	-	0/1/1/1	-
3	EDO	B	208	-	-	0/1/1/1	-
3	EDO	A	203	-	-	1/1/1/1	-
3	EDO	B	202	-	-	0/1/1/1	-
3	EDO	B	204	-	-	0/1/1/1	-
3	EDO	A	204	-	-	1/1/1/1	-
3	EDO	A	209	-	-	0/1/1/1	-
3	EDO	B	206	-	-	1/1/1/1	-
3	EDO	B	205	-	-	1/1/1/1	-
3	EDO	B	211	-	-	1/1/1/1	-
3	EDO	A	211	-	-	1/1/1/1	-
3	EDO	A	213	-	-	1/1/1/1	-
3	EDO	A	208	-	-	0/1/1/1	-
3	EDO	B	212	-	-	1/1/1/1	-
3	EDO	B	210	-	-	0/1/1/1	-
3	EDO	A	201	-	-	1/1/1/1	-
3	EDO	A	210	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	203	EDO	O1-C1-C2-O2
3	B	205	EDO	O1-C1-C2-O2
3	B	209	EDO	O1-C1-C2-O2
3	A	201	EDO	O1-C1-C2-O2
3	A	204	EDO	O1-C1-C2-O2
3	A	210	EDO	O1-C1-C2-O2
3	A	211	EDO	O1-C1-C2-O2
3	A	212	EDO	O1-C1-C2-O2
3	B	211	EDO	O1-C1-C2-O2
3	A	203	EDO	O1-C1-C2-O2
3	B	206	EDO	O1-C1-C2-O2
3	A	213	EDO	O1-C1-C2-O2
3	B	212	EDO	O1-C1-C2-O2

There are no ring outliers.

4 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	209	EDO	6	0
3	A	212	EDO	3	0
3	A	204	EDO	1	0
3	A	213	EDO	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	190/215 (88%)	-0.25	0 100 100	23, 31, 46, 56	0
1	B	190/215 (88%)	-0.27	0 100 100	24, 33, 49, 57	0
All	All	380/430 (88%)	-0.26	0 100 100	23, 32, 47, 57	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	EDO	A	211	4/4	0.69	0.25	47,57,58,60	1
3	EDO	B	210	4/4	0.76	0.21	55,56,57,57	1
3	EDO	B	212	4/4	0.81	0.19	51,54,56,56	1
3	EDO	A	203	4/4	0.82	0.19	38,43,45,45	1
3	EDO	B	211	4/4	0.84	0.16	39,48,50,50	1
3	EDO	B	209	4/4	0.84	0.41	40,52,53,54	1
3	EDO	B	206	4/4	0.86	0.42	47,53,57,59	1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	EDO	B	205	4/4	0.87	0.10	42,45,46,47	1
3	EDO	A	209	4/4	0.88	0.17	35,42,43,45	1
3	EDO	A	212	4/4	0.89	0.41	51,53,57,58	1
3	EDO	A	213	4/4	0.89	0.27	46,48,52,53	1
3	EDO	A	201	4/4	0.91	0.17	54,58,66,66	1
3	EDO	A	208	4/4	0.91	0.16	48,52,60,64	1
2	SO4	B	207	5/5	0.92	0.13	54,63,68,72	0
3	EDO	A	204	4/4	0.93	0.19	37,41,44,46	1
3	EDO	B	202	4/4	0.93	0.17	51,52,57,57	1
3	EDO	B	204	4/4	0.93	0.26	42,46,49,49	1
3	EDO	A	210	4/4	0.94	0.19	46,50,52,54	1
3	EDO	B	203	4/4	0.94	0.12	45,46,48,48	1
2	SO4	A	206	5/5	0.95	0.22	60,61,66,68	0
3	EDO	B	208	4/4	0.96	0.30	39,47,52,55	1
3	EDO	A	207	4/4	0.97	0.12	35,36,38,38	1
2	SO4	B	201	5/5	0.98	0.12	37,40,47,48	0
2	SO4	A	205	5/5	0.99	0.10	51,51,61,66	0
2	SO4	A	202	5/5	0.99	0.10	37,38,43,46	0

6.5 Other polymers [i](#)

There are no such residues in this entry.