

wwPDB EM Validation Summary Report (i)

Nov 11, 2023 – 03:39 pm GMT

PDB ID	:	8CGK
EMDB ID	:	EMD-16646
Title	:	Lincomycin and Avilamycin bound to the 50S subunit
Authors	:	Paternoga, H.; Crowe-McAuliffe, C.; Beckert, B.; Wilson, D.N.
Deposited on	:	2023-02-05
Resolution	:	1.64 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/EMValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis	:	0.0.1.dev70
Mogul	:	1.8.4, CSD as541be (2020)
MolProbity	:	4.02b-467
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ	:	1.9.9
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 1.64 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f EM\ structures}\ (\#{f Entries})$		
Clashscore	158937	4297		
Ramachandran outliers	154571	4023		
Sidechain outliers	154315	3826		
RNA backbone	4643	859		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for $\geq=3, 2, 1$ and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq=5\%$ The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	0	55	91%	9%
2	1	46	98%	•
3	2	65	95%	• •
4	3	38	92%	5% •
5	a	2904	80% 12%	o 7%
6	b	120	5% 31% • 68%	
7	с	273	99%	



Mol	Chain	Length	Quality of chain	
8	d	209	5% 99%	•
9	е	201	18%	
10	g	177	60%	5%
11	h	149	15% 28% 72%	
12	i	142	99%	
13	j	123	99%	·
14	k	144	6%	
15	1	136	<mark>6%</mark> 99%	·
16	m	127	93%	7%
17	О	115	97%	••
18	р	118	98%	
19	q	103	9%	
20	r	110	5% 99%	
21	s	100	92%	• 7%
22	t	104	8%	• 11%
23	u	94	20%	•
24	V	85	88%	12%
25	W	78	97%	••
26	x	63	33%	5%
27	У	59	8%	
28	Z	57	95%	5%

Continued from previous page...

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
32	6UQ	a	3003	Х	-	-	-



2 Entry composition (i)

There are 35 unique types of molecules in this entry. The entry contains 88150 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Large ribosomal subunit protein bL33.

Mol	Chain	Residues		Aton	ns	AltConf	Trace	
1	0	50	Total 413	C 267	N 75	0 71	0	0

• Molecule 2 is a protein called Large ribosomal subunit protein bL34.

Mol	Chain	Residues		Ato	\mathbf{ms}	AltConf	Trace		
2	1	45	Total	С	N	0	S	0	0
			367	222	88	55	2		

• Molecule 3 is a protein called Large ribosomal subunit protein bL35.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	2	64	Total 504	C 323	N 105	0 74	${ m S} { m 2}$	0	0

• Molecule 4 is a protein called Large ribosomal subunit protein bL36A.

Mol	Chain	Residues		Ato	\mathbf{ms}	AltConf	Trace		
4	3	38	Total 302	C 185	N 65	O 48	$\begin{array}{c} \mathrm{S} \\ \mathrm{4} \end{array}$	0	0

• Molecule 5 is a RNA chain called 23S rRNA.

Mol	Chain	Residues			AltConf	Trace			
5	a	2699	Total 57983	C 25871	N 10696	O 18717	Р 2699	0	0

• Molecule 6 is a RNA chain called 5S rRNA.

Mol	Chain	Residues		\mathbf{A}^{\dagger}	AltConf	Trace			
6	b	39	Total 840	C 373	N 153	0 275	Р 39	0	0



• Molecule 7 is a protein called Large ribosomal subunit protein uL2.

Mol	Chain	Residues		Ate	oms			AltConf	Trace
7	С	271	Total 2082	C 1288	N 423	0 364	${f S}7$	0	0

• Molecule 8 is a protein called Large ribosomal subunit protein uL3.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	d	207	Total 1552	C 972	N 286	O 291	${ m S} { m 3}$	0	0

• Molecule 9 is a protein called Large ribosomal subunit protein uL4.

Mol	Chain	Residues		At	oms	AltConf	Trace		
9	е	201	Total 1552	С 974	N 283	O 290	${ m S}{ m 5}$	0	0

• Molecule 10 is a protein called Large ribosomal subunit protein uL6.

Mol	Chain	Residues		At	oms			AltConf	Trace
10	g	168	Total 1255	C 791	N 228	0 234	${ m S} { m 2}$	0	0

• Molecule 11 is a protein called Large ribosomal subunit protein bL9.

Mol	Chain	Residues		Atc	\mathbf{ms}	AltConf	Trace		
11	h	41	Total 303	C 194	N 54	0 54	S 1	0	0

• Molecule 12 is a protein called Large ribosomal subunit protein uL13.

Mol	Chain	Residues	Atoms					AltConf	Trace
12	i	141	Total 1121	C 709	N 211	0 198	${ m S} { m 3}$	0	0

• Molecule 13 is a protein called Large ribosomal subunit protein uL14.

Mol	Chain	Residues		At	oms	AltConf	Trace		
13	j	123	Total 946	C 593	N 181	0 166	S 6	0	0

• Molecule 14 is a protein called 50S ribosomal protein L15.



Mol	Chain	Residues		At	oms			AltConf	Trace
14	k	144	Total 1053	$\begin{array}{c} \mathrm{C} \\ 654 \end{array}$	N 207	O 190	${ m S} { m 2}$	0	0

• Molecule 15 is a protein called Large ribosomal subunit protein uL16.

Mol	Chain	Residues		At	oms	AltConf	Trace		
15	1	136	Total 1075	C 686	N 205	0 177	${ m S} 7$	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
1	82	MS6	MET	modified residue	UNP P0ADY7

• Molecule 16 is a protein called Large ribosomal subunit protein bL17.

Mol	Chain	Residues		At	\mathbf{oms}			AltConf	Trace
16	m	118	Total 945	C 585	N 194	0 161	${ m S}{ m 5}$	0	0

• Molecule 17 is a protein called Large ribosomal subunit protein bL19.

Mol	Chain	Residues		At	oms	AltConf	Trace		
17	О	114	Total 917	C 574	N 179	0 163	S 1	0	0

• Molecule 18 is a protein called Large ribosomal subunit protein bL20.

Mol	Chain	Residues	Atoms				AltConf	Trace
18	р	117	Total 947	C 604	N 192	O 151	0	0

• Molecule 19 is a protein called Large ribosomal subunit protein bL21.

Mol	Chain	Residues	Atoms					AltConf	Trace
19	q	103	Total 816	C 516	N 153	0 145	${ m S} { m 2}$	0	0

• Molecule 20 is a protein called Large ribosomal subunit protein uL22.



Mol	Chain	Residues	Atoms					AltConf	Trace
20	r	109	Total 845	C 526	N 162	O 154	${ m S} { m 3}$	0	0

• Molecule 21 is a protein called Large ribosomal subunit protein uL23.

Mol	Chain	Residues	Atoms					AltConf	Trace
21	s	93	Total 738	C 466	N 139	0 131	${ m S} { m 2}$	0	0

• Molecule 22 is a protein called Large ribosomal subunit protein uL24.

Mol	Chain	Residues		Ato	ms	AltConf	Trace	
22	t	93	Total 717	C 452	N 135	O 130	0	0

• Molecule 23 is a protein called 50S ribosomal protein L25.

Mol	Chain	Residues	Atoms					AltConf	Trace
23	u	93	Total 745	C 474	N 136	0 133	${ m S} { m 2}$	0	0

• Molecule 24 is a protein called Large ribosomal subunit protein bL27.

Mol	Chain	Residues		Atoms					Trace
24	V	75	Total 569	C 353	N 113	0 102	S 1	0	0

• Molecule 25 is a protein called Large ribosomal subunit protein bL28.

Mol	Chain	Residues	Atoms					AltConf	Trace
25	W	77	Total 625	C 388	N 129	O 106	${ m S} { m 2}$	0	0

• Molecule 26 is a protein called Large ribosomal subunit protein uL29.

Mol	Chain	Residues	Atoms					AltConf	Trace
26	x	60	Total	С	Ν	0	\mathbf{S}	0	0
20		00	491	303	96	91	1	Ŭ	0

• Molecule 27 is a protein called Large ribosomal subunit protein uL30.



Mol	Chain	Residues	Atoms					AltConf	Trace
97		59	Total	С	Ν	Ο	S	0	0
21	У		449	281	87	79	2	0	0

• Molecule 28 is a protein called Large ribosomal subunit protein bL32.

Mol	Chain	Residues	Atoms					AltConf	Trace
28	Z	54	Total 429	C 260	N 91	0 77	S 1	0	0

• Molecule 29 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	AltConf
29	3	1	Total Zn 1 1	0

• Molecule 30 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms	AltConf
30	a	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0

• Molecule 31 is LINCOMYCIN (three-letter code: 3QB) (formula: $C_{18}H_{34}N_2O_6S$) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms				AltConf	
21	9	1	Total	С	Ν	0	S	0
51	a	1	27	18	2	6	1	0

• Molecule 32 is (2R,3S,4R,6S)-4-hydroxy-6-{[(2R,3aR,4R,4'R,5'S,6S,6'R,7aR)-4'-hydroxy-6-{[(2S,3R,4R,5S,6R)-3-hydroxy-2-{[(2R,3S,4S,5S,6S)-4-hydroxy-6-({(2R,3aS,3a'R,6S,6'R,7R,7'R,7aR,7a'R)-7'-hydroxy-7'-[(1S)-1-hydroxyethyl]-6'-methyl-7-[(2-methylpropanoyl)oxy]o ctahydro-4H-2,4'-spirobi[[1,3]dioxolo[4,5-c]pyran]-6-yl}oxy)-5-methoxy-2-(methoxymethyl) tetrahydro-2H-pyran-3-yl]oxy}-5-methoxy-6-methyltetrahydro-2H-pyran-4-yl]oxy}-4,6',7 a-trimethyloctahydro-4H-spiro[1,3-dioxolo[4,5-c]pyran-2,2'-pyran]-5'-yl]oxy}-2-methyltetrahydro-2H-pyran-3-yl 3,5-dichloro-4-hydroxy-2-methoxy-6-methylbenzoate (non-preferred name) (three-letter code: 6UQ) (formula: C₆₁H₉₀Cl₂O₃₂) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms				AltConf
32	a	1	Total 95	C 61	Cl 2	O 32	0

• Molecule 33 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	AltConf
33	a	251	Total Mg 251 251	0
33	b	3	Total Mg 3 3	0
33	с	3	Total Mg 3 3	0
33	d	1	Total Mg 1 1	0
33	р	1	Total Mg 1 1	0
33	Z	1	Total Mg 1 1	0

• Molecule 34 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms	AltConf
34	a	77	Total K 77 77	0
34	с	3	Total K 3 3	0
34	е	1	Total K 1 1	0

• Molecule 35 is water.

Mol	Chain	Residues	Atoms	AltConf
35	0	6	Total O 6 6	0
35	1	36	$\begin{array}{cc} \text{Total} & \text{O} \\ 36 & 36 \end{array}$	0
35	2	31	$\begin{array}{cc} \text{Total} & \text{O} \\ 31 & 31 \end{array}$	0
35	3	16	Total O 16 16	0
35	a	6026	Total O 6026 6026	0



Continued from previous page...

Mol	Chain	Residues	Atoms	AltConf
25	h	00	Total O	0
50	D	00	83 83	0
25	0	122	Total O	0
50	С	155	133 133	0
25	d	05	Total O	0
- 55	u	90	$95 ext{ }95$	0
35	0	63	Total O	0
- 55	е	05	63 63	0
35	ď	9	Total O	0
- 55	g	2	2 2	0
35	h	1	Total O	0
- 55	11	T	1 1	0
35	i	56	Total O	0
- 00	1	50	56 - 56	0
35	i	32	Total O	0
- 00	J	52	32 32	0
35	k	62	Total O	0
- 00	К	02	62 62	0
35	1	56	Total O	0
- 55	I	50	56 - 56	0
35	m	52	Total O	0
- 55	111	52	52 52	0
35	0	30	Total O	0
00	0	50	30 30	0
35	n	56	Total O	0
- 00	Р	50	56 - 56	0
35	n	32	Total O	0
00	Ч		32 32	0
35	r	54	Total O	0
	1		54 54	Ŭ
35	S	27	Total O	0
			27 27	Ŭ
35	t	14	Total O	0
	Ŭ	**	14 14	
35	11	19	Total O	0
	~		19 19	
35	v	20	Total O	0
			20 20	
35	w	27	Total O	0
			27 27	
35	x	10	Total O	0
00	л	10	10 10	



Continued from previous page...

Mol	Chain	Residues	Atoms	AltConf
35	У	20	Total O 20 20	0
35	Z	42	$\begin{array}{cc} \text{Total} & \text{O} \\ 42 & 42 \end{array}$	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Large ribosomal subunit protein bL33















• Molecule 15: Large ribosomal subunit protein uL16



6%		
Jhain I:	99%	•
MI K58 K58 R59 Q60 Q60 T134 V135 M136		
Molecule 16: Large ribo	osomal subunit protein bL17	
Chain m:	93%	7%
R118 SER CLU CLU CLU ALA ALA ALA ALA CLU CLU		
Molecule 17: Large ribc	osomal subunit protein bL19	
Chain o:	97%	••
22 015 116 116 116 116 116 116 116 116 116 1	N115	
Molecule 18: Large ribo	somal subunit protein bL20	
Chain p:	98%	
A2 861 A118		
Molecule 19: Large ribe	somal subunit protein bL21	
9% Chain q:	100%	
E34 F35 A35 A35 A35 C44 E45 E45 E45 A102 A103		
Molecule 20: Large ribc	osomal subunit protein uL22	
Chain r:	99%	.
De5 De7 De7 R95 ARC		
Molecule 21: Large ribc	somal subunit protein uL23	
thain s:	92%	• 7%
• • • • • • • • • • • • • • • • • • • •	•	
н1 E52 H70 H70 G71 G71 G71 G71 G72 C73 G72 G90 G91 G91	L93 PARP PARP CLY CLY CLY CLY CLY	

R L D W I D E PDB TEIN DATA BANK

• Molecule 22: Large ribosomal subunit protein uL24	
Chain t: 88%	11%
MET A2 A2 A2 A2 A1 A1 A1 A1 A1 A1 A1 A1 A1 A1 A1 A1 A1	
• Molecule 23: 50S ribosomal protein L25	
Chain u: 99%	·
MET T3 T3 T4 K10 K10 K10 G33 K53 K53 K53 K53 K53 K53 K53 K	
\bullet Molecule 24: Large ribosomal subunit protein bL27	
25% Chain v: 88%	12%
MET ALA ALA ALA ALA ALA ALA ALA ALA ALA AL	
\bullet Molecule 25: Large ribosomal subunit protein bL28	
Chain w: 97%	
MET 82 84 84 84 84 84 84 84 84 84 84 84 84 84	
\bullet Molecule 26: Large ribosomal subunit protein uL29	
Chain x: 95%	5%
MET K2 K2 K3 E5 E5 C1 C1 E1 E1 C3 C3 C3 C3 C3 C3 C3 C3 C3 C3 C3 C3 C3	
\bullet Molecule 27: Large ribosomal subunit protein uL30	
Chain y: 98%	·
MET A 2 B 40 B 55 B 55 B 55 B 55 B 55 B 55 B 55 B 5	
\bullet Molecule 28: Large ribosomal subunit protein bL32	
Chain z: 95%	5%







4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	419159	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE	Depositor
	CORRECTION	
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{\AA}^2)$	60	Depositor
Minimum defocus (nm)	400	Depositor
Maximum defocus (nm)	1000	Depositor
Magnification	Not provided	
Image detector	FEI FALCON IV $(4k \ge 4k)$	Depositor
Maximum map value	0.208	Depositor
Minimum map value	-0.078	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.002	Depositor
Recommended contour level	0.0164	Depositor
Map size (Å)	544.8, 544.8, 544.8	wwPDB
Map dimensions	800, 800, 800	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.681, 0.681, 0.681	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: H2U, MEQ, 1MG, 2MG, OMU, MG, 4D4, 3QB, OMC, 5MC, K, PSU, ZN, 6MZ, 5MU, 6UQ, ACT, G7M, 2MA, MS6, OMG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond lengths		Bond angles		
WIOI	Ullaili	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	0	0.30	0/420	0.60	0/560	
2	1	0.31	0/370	0.72	0/487	
3	2	0.31	0/513	0.66	0/676	
4	3	0.29	0/303	0.70	0/397	
5	a	0.49	0/64435	1.13	66/100513~(0.1%)	
6	b	0.54	0/938	1.15	0/1460	
7	с	0.31	0/2121	0.69	0/2852	
8	d	0.29	0/1562	0.64	0/2102	
9	е	0.28	0/1571	0.62	0/2113	
10	g	0.31	0/1273	0.63	0/1725	
11	h	0.32	0/306	0.69	0/413	
12	i	0.29	0/1144	0.64	0/1541	
13	j	0.30	0/955	0.68	0/1279	
14	k	0.32	0/1062	0.64	0/1413	
15	1	0.28	0/1073	0.66	0/1433	
16	m	0.28	0/958	0.63	0/1281	
17	0	0.30	0/929	0.64	0/1242	
18	р	0.30	0/960	0.66	0/1278	
19	q	0.31	0/829	0.66	0/1107	
20	r	0.29	0/852	0.63	0/1142	
21	s	0.27	0/744	0.64	0/994	
22	t	0.28	0/721	0.63	0/956	
23	u	0.29	0/758	0.65	0/1015	
24	V	0.33	0/576	0.66	0/762	
25	W	0.32	0/635	0.69	0/848	
26	X	0.27	0/492	0.54	0/655	
27	У	0.28	0/453	0.65	0/605	
28	Z	0.32	0/435	0.66	0/581	
All	All	0.45	0/87388	1.04	$66/\overline{131430}~(0.1\%)$	



Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	2	0	1
4	3	0	1
5	a	0	1
17	0	0	1
18	р	0	1
21	s	0	1
25	W	0	1
All	All	0	7

There are no bond length outliers.

The worst 5 of 66 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
5	a	512	G	O4'-C1'-N9	11.51	117.41	108.20
5	a	2429	G	O3'-P-O5'	-11.19	82.75	104.00
5	a	196	А	O5'-P-OP1	-8.28	98.24	105.70
5	a	1971	U	O3'-P-O5'	-8.05	88.70	104.00
5	a	2848	G	O4'-C1'-N9	7.60	114.28	108.20

There are no chirality outliers.

5 of 7 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	2	13	ARG	Sidechain
4	3	36	ARG	Sidechain
5	а	512	G	Sidechain
17	0	88	ARG	Sidechain
18	р	51	ARG	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	0	413	0	448	0	0
2	1	367	0	405	0	0
3	2	504	0	572	0	0
4	3	302	0	340	1	0
5	a	57983	0	29161	0	0
6	b	840	0	423	0	0
7	с	2082	0	2153	0	0
8	d	1552	0	1601	0	0
9	е	1552	0	1619	0	0
10	g	1255	0	1296	0	0
11	h	303	0	327	0	0
12	i	1121	0	1150	0	0
13	j	946	0	1023	0	0
14	k	1053	0	1129	0	0
15	1	1075	0	1145	0	0
16	m	945	0	989	0	0
17	0	917	0	962	0	0
18	р	947	0	1019	0	0
19	q	816	0	839	0	0
20	r	845	0	909	0	0
21	s	738	0	807	0	0
22	t	717	0	766	0	0
23	u	745	0	768	0	0
24	V	569	0	581	0	0
25	W	625	0	652	0	0
26	Х	491	0	523	0	0
27	у	449	0	488	0	0
28	Z	429	0	440	0	0
29	3	1	0	0	0	0
30	a	4	0	3	0	0
31	a	27	0	34	0	0
32	a	95	0	0	0	0
33	a	251	0	0	0	0
33	b	3	0	0	0	0
33	с	3	0	0	0	0
33	d	1	0	0	0	0
33	р	1	0	0	0	0
33	Z	1	0	0	0	0
34	a	77	0	0	0	0
34	с	3	0	0	0	0
34	е	1	0	0	0	0
35	0	6	0	0	0	0
35	1	36	0	0	0	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
35	2	31	0	0	0	0
35	3	16	0	0	0	0
35	a	6026	0	0	0	0
35	b	83	0	0	0	0
35	с	133	0	0	0	0
35	d	95	0	0	0	0
35	е	63	0	0	0	0
35	g	2	0	0	0	0
35	h	1	0	0	0	0
35	i	56	0	0	0	0
35	j	32	0	0	0	0
35	k	62	0	0	0	0
35	l	56	0	0	0	0
35	m	52	0	0	0	0
35	0	30	0	0	0	0
35	р	56	0	0	0	0
35	q	32	0	0	0	0
35	r	54	0	0	0	0
35	s	27	0	0	0	0
35	t	14	0	0	0	0
35	u	19	0	0	0	0
35	V	20	0	0	0	0
35	W	27	0	0	0	0
35	Х	10	0	0	0	0
35	У	20	0	0	0	0
35	Z	42	0	0	0	0
All	All	88150	0	52572	1	0

Continued from previous page...

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All (1) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:3:16:ILE:HD13	4:3:25:VAL:HG22	1.96	0.47

There are no symmetry-related clashes.



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	0	48/55~(87%)	48 (100%)	0	0	100	100
2	1	43/46~(94%)	43 (100%)	0	0	100	100
3	2	62/65~(95%)	60 (97%)	2(3%)	0	100	100
4	3	36/38~(95%)	36 (100%)	0	0	100	100
7	с	269/273~(98%)	261 (97%)	8 (3%)	0	100	100
8	d	204/209~(98%)	198 (97%)	6(3%)	0	100	100
9	е	199/201~(99%)	194 (98%)	5 (2%)	0	100	100
10	g	164/177~(93%)	158 (96%)	6 (4%)	0	100	100
11	h	39/149~(26%)	37 (95%)	2(5%)	0	100	100
12	i	139/142~(98%)	139 (100%)	0	0	100	100
13	j	121/123~(98%)	118 (98%)	3 (2%)	0	100	100
14	k	142/144~(99%)	139 (98%)	3 (2%)	0	100	100
15	1	132/136~(97%)	129 (98%)	3 (2%)	0	100	100
16	m	116/127~(91%)	112 (97%)	4 (3%)	0	100	100
17	0	112/115~(97%)	108 (96%)	4 (4%)	0	100	100
18	р	115/118 (98%)	114 (99%)	1 (1%)	0	100	100
19	q	101/103~(98%)	100 (99%)	1 (1%)	0	100	100
20	r	107/110~(97%)	106 (99%)	1 (1%)	0	100	100
21	S	91/100 (91%)	89 (98%)	2(2%)	0	100	100
22	t	89/104 (86%)	87 (98%)	2(2%)	0	100	100
23	u	91/94~(97%)	88 (97%)	3(3%)	0	100	100
24	V	73/85~(86%)	71 (97%)	2(3%)	0	100	100
25	W	75/78~(96%)	75 (100%)	0	0	100	100
26	x	58/63~(92%)	58 (100%)	0	0	100	100
27	У	56/59~(95%)	55 (98%)	1 (2%)	0	100	100



Contre	nucu jion	i previous page					
Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
28	z	52/57~(91%)	51 (98%)	1 (2%)	0	100	100
All	All	2734/2971~(92%)	2674 (98%)	60 (2%)	0	100	100

Continued from previous page...

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	0	46/49~(94%)	46 (100%)	0	100	100
2	1	37/38~(97%)	37~(100%)	0	100	100
3	2	51/52~(98%)	50~(98%)	1 (2%)	55	29
4	3	34/34~(100%)	33~(97%)	1 (3%)	42	15
7	с	216/218~(99%)	215 (100%)	1 (0%)	88	80
8	d	162/163~(99%)	161 (99%)	1 (1%)	86	75
9	е	165/165~(100%)	165 (100%)	0	100	100
10	g	130/138 (94%)	130 (100%)	0	100	100
11	h	32/114~(28%)	32 (100%)	0	100	100
12	i	115/116~(99%)	115 (100%)	0	100	100
13	j	104/104~(100%)	103 (99%)	1 (1%)	76	59
14	k	103/103~(100%)	103 (100%)	0	100	100
15	1	107/107~(100%)	106 (99%)	1 (1%)	78	63
16	m	98/103~(95%)	98 (100%)	0	100	100
17	О	99/100~(99%)	98~(99%)	1 (1%)	76	59
18	р	89/90~(99%)	89 (100%)	0	100	100
19	q	84/84~(100%)	84 (100%)	0	100	100
20	r	92/93~(99%)	92 (100%)	0	100	100
21	s	80/84~(95%)	80 (100%)	0	100	100
22	t	76/85~(89%)	75~(99%)	1 (1%)	69	47



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
23	u	77/78~(99%)	77~(100%)	0	100 100
24	v	56/63~(89%)	56 (100%)	0	100 100
25	W	67/68~(98%)	67~(100%)	0	100 100
26	х	54/55~(98%)	54 (100%)	0	100 100
27	У	48/49~(98%)	48 (100%)	0	100 100
28	Z	46/48~(96%)	46 (100%)	0	100 100
All	All	2268/2401 (94%)	2260 (100%)	8 (0%)	91 84

Continued from previous page...

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
22	t	24	LYS
17	0	87	LYS
13	j	58	LEU
8	d	32	ASN
15	1	6	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
7	с	37	ASN
17	0	115	ASN
26	Х	45	GLN
27	у	49	ASN

5.3.3 RNA (i)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
5	a	2689/2904~(92%)	296 (11%)	0
6	b	37/120~(30%)	2(5%)	0
All	All	2726/3024~(90%)	298 (10%)	0

5 of 298 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
5	a	10	А
5	а	34	U
5	a	45	G



Continued from previous page...

Mol	Chain	Res	Type
5	а	71	А
5	a	74	А

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

23 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Bos	Link	Bo	ond leng	ths	B	ond ang	les
	туре	Ullalli	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	5MU	a	1939	34,5	19,22,23	0.35	0	28,32,35	0.47	0
5	PSU	a	2580	$34,\!5$	18,21,22	1.06	1 (5%)	22,30,33	0.73	0
5	PSU	a	2504	34,5	18,21,22	0.93	1 (5%)	22,30,33	0.73	0
5	2MG	a	1835	5	18,26,27	1.04	2 (11%)	16,38,41	0.69	0
5	PSU	a	2605	5	18,21,22	0.95	1 (5%)	22,30,33	0.89	1 (4%)
5	PSU	a	2457	5	18,21,22	0.99	1 (5%)	22,30,33	0.68	0
5	PSU	a	955	5	18,21,22	0.90	1 (5%)	22,30,33	0.65	0
5	G7M	a	2069	5	20,26,27	1.09	2 (10%)	17,39,42	0.67	0
5	H2U	a	2449	5	18,21,22	0.57	0	21,30,33	0.70	0
5	OMC	a	2498	$5,\!33$	19,22,23	0.29	0	$26,\!31,\!34$	0.63	0
5	5MU	a	747	5	19,22,23	0.31	0	28,32,35	0.47	0
5	5MC	a	1962	5	18,22,23	0.36	0	26,32,35	0.69	0
8	MEQ	d	150	8	8,9,10	0.47	0	5,10,12	1.33	0
5	2MG	a	2445	5	18,26,27	1.09	2 (11%)	16,38,41	0.76	0
5	OMU	a	2552	5	19,22,23	0.25	0	26,31,34	0.44	0
15	4D4	1	81	15	9,11,12	0.45	0	8,13,15	1.06	1 (12%)
5	PSU	a	2604	5	18,21,22	0.96	1 (5%)	22,30,33	0.75	1 (4%)
5	6MZ	a	2030	5	18,25,26	0.71	0	16,36,39	0.96	1 (6%)
5	OMG	a	2251	34,5	18,26,27	1.03	2 (11%)	19,38,41	0.73	0
5	1MG	a	745	5	18,26,27	1.04	2 (11%)	19,39,42	0.47	0
5	PSU	a	746	5,33	18,21,22	0.99	1(5%)	22,30,33	0.66	0



Mol ′	Tune	Chain	Dec	Tink	Bo	Bond lengths			Bond angles		
	туре		nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
5	6MZ	a	1618	5	18,25,26	0.74	0	16,36,39	0.83	1 (6%)	
5	2MA	a	2503	34,5,33	17,25,26	0.99	1 (5%)	17,37,40	0.98	1 (5%)	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	5MU	a	1939	34,5	-	0/7/25/26	0/2/2/2
5	PSU	a	2580	34,5	-	0/7/25/26	0/2/2/2
5	PSU	a	2504	34,5	-	0/7/25/26	0/2/2/2
5	2MG	a	1835	5	-	0/5/27/28	0/3/3/3
5	PSU	a	2605	5	-	0/7/25/26	0/2/2/2
5	PSU	a	2457	5	-	0/7/25/26	0/2/2/2
5	PSU	a	955	5	-	0/7/25/26	0/2/2/2
5	G7M	a	2069	5	-	2/3/25/26	0/3/3/3
5	H2U	a	2449	5	-	0/7/38/39	0/2/2/2
5	OMC	a	2498	5,33	-	0/9/27/28	0/2/2/2
5	5MU	a	747	5	-	0/7/25/26	0/2/2/2
5	5MC	a	1962	5	-	0/7/25/26	0/2/2/2
8	MEQ	d	150	8	-	2/8/9/11	-
5	2MG	a	2445	5	-	0/5/27/28	0/3/3/3
5	OMU	a	2552	5	-	1/9/27/28	0/2/2/2
15	4D4	1	81	15	-	0/11/12/14	-
5	PSU	а	2604	5	-	0/7/25/26	0/2/2/2
5	6MZ	a	2030	5	-	2/5/27/28	0/3/3/3
5	OMG	a	2251	34,5	-	0/5/27/28	0/3/3/3
5	1MG	a	745	5	-	0/3/25/26	0/3/3/3
5	PSU	a	746	5,33	-	1/7/25/26	0/2/2/2
5	6MZ	a	1618	5	-	0/5/27/28	0/3/3/3
5	2MA	a	2503	34,5,33	-	1/3/25/26	0/3/3/3

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	$\operatorname{Ideal}(\operatorname{\AA})$
5	a	2580	PSU	C6-C5	4.23	1.40	1.35
5	a	2457	PSU	C6-C5	4.05	1.40	1.35
5	a	746	PSU	C6-C5	3.84	1.39	1.35
5	a	2604	PSU	C6-C5	3.77	1.39	1.35
5	a	2504	PSU	C6-C5	3.73	1.39	1.35



Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
5	a	2503	2MA	CM2-C2-N1	2.94	122.77	116.23
5	a	2030	6MZ	C2-N1-C6	2.69	118.90	116.59
5	a	1618	6MZ	C2-N1-C6	2.50	118.73	116.59
5	a	2604	PSU	C2'-C3'-C4'	-2.30	98.17	102.64
5	a	2605	PSU	C2'-C3'-C4'	-2.02	98.72	102.64

The worst 5 of 6 bond angle outliers are listed below:

There are no chirality outliers.

5 of 9 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	d	150	MEQ	NE2-CD-CG-CB
8	d	150	MEQ	OE1-CD-CG-CB
5	a	2030	6MZ	O4'-C4'-C5'-O5'
5	а	2069	G7M	C4'-C5'-O5'-P
5	a	2030	6MZ	C3'-C4'-C5'-O5'

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 345 ligands modelled in this entry, 342 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type Cha	Turne	Chain	Dec	Tink	Bon	Bond lengths			Bond angles		
	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2		
32	6UQ	a	3003	-	100,105,105	1.13	5 (5%)	132,164,164	1.43	17 (12%)	
31	3QB	a	3002	-	25,28,28	0.29	0	$29,\!40,\!40$	0.85	3 (10%)	
30	ACT	a	3001	-	3,3,3	0.96	0	3,3,3	0.79	0	



In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
32	6UQ	a	3003	-	2/2/35/35	11/47/211/211	0/11/11/11
31	3QB	a	3002	-	-	1/21/53/53	0/2/2/2

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	Observed(Å)	Ideal(Å)
32	a	3003	6UQ	ODC-CDA	-8.32	1.21	1.43
32	a	3003	6UQ	OCP-CCJ	3.17	1.44	1.40
32	a	3003	6UQ	OAY-CAX	3.14	1.45	1.41
32	a	3003	6UQ	CAD-CAC	2.77	1.44	1.40
32	a	3003	6UQ	OCG-CCB	2.02	1.46	1.41

All (5) bond length outliers are listed below:

The worst 5 of 20 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
32	a	3003	6UQ	CAX-OAY-CAZ	6.71	118.28	112.16
32	a	3003	6UQ	ODC-CDA-CDB	4.41	119.25	109.02
32	a	3003	6UQ	OBE-CBD-CBI	3.80	116.74	108.57
32	a	3003	6UQ	OBB-CBC-CBD	-3.79	97.29	101.85
32	a	3003	6UQ	OCK-CCE-CCF	3.67	117.31	110.45

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
32	а	3003	6UQ	CDA
32	а	3003	6UQ	CCN

5 of 12 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
32	а	3003	6UQ	CCM-CCN-CDA-ODC
32	а	3003	6UQ	OCT-CCN-CDA-ODC
32	a	3003	6UQ	CCO-CCN-CDA-ODC
32	a	3003	6UQ	CAF-CAE-ODD-CDE
32	a	3003	6UQ	CAC-CAD-CAJ-OAK

There are no ring outliers.

No monomer is involved in short contacts.



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and similar rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-16646. These allow visual inspection of the internal detail of the map and identification of artifacts.

Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections (i)

6.1.1 Primary map



6.1.2 Raw map



The images above show the map projected in three orthogonal directions.



6.2 Central slices (i)

6.2.1 Primary map



X Index: 400





Z Index: 400

6.2.2 Raw map



X Index: 400

Y Index: 400



The images above show central slices of the map in three orthogonal directions.



6.3 Largest variance slices (i)

6.3.1 Primary map



X Index: 403



Y Index: 391



Z Index: 467

6.3.2 Raw map



X Index: 403

Y Index: 348



The images above show the largest variance slices of the map in three orthogonal directions.



6.4 Orthogonal standard-deviation projections (False-color) (i)

6.4.1 Primary map



6.4.2 Raw map



The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.



6.5 Orthogonal surface views (i)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.0164. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.

6.6 Mask visualisation (i)

This section was not generated. No masks/segmentation were deposited.



7 Map analysis (i)

This section contains the results of statistical analysis of the map.

7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



7.2 Volume estimate (i)



The volume at the recommended contour level is 202 $\rm nm^3;$ this corresponds to an approximate mass of 183 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



7.3 Rotationally averaged power spectrum (i)



*Reported resolution corresponds to spatial frequency of 0.610 \AA^{-1}



8 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC (i)



*Reported resolution corresponds to spatial frequency of 0.610 ${\rm \AA^{-1}}$



8.2 Resolution estimates (i)

$\mathbf{B}_{\mathrm{assolution ostimato}}(\mathbf{\hat{\lambda}})$	Estimation criterion (FSC cut-off)			
Resolution estimate (A)	0.143	0.5	Half-bit	
Reported by author	1.64	-	-	
Author-provided FSC curve	-	-	-	
Unmasked-calculated*	1.96	2.54	2.01	

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 1.96 differs from the reported value 1.64 by more than 10 %



9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-16646 and PDB model 8CGK. Per-residue inclusion information can be found in section 3 on page 13.

9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.0164 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.



9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.0164).



9.4 Atom inclusion (i)



At the recommended contour level, 83% of all backbone atoms, 84% of all non-hydrogen atoms, are inside the map.



9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.0164) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8380	0.7990
0	0.7310	0.7800
1	0.9740	0.8780
2	0.9570	0.8670
3	0.8800	0.8240
a	0.8500	0.7980
b	0.8230	0.7590
с	0.9050	0.8470
d	0.8990	0.8420
е	0.7530	0.7830
g	0.3430	0.6350
h	0.4600	0.6520
i	0.9000	0.8430
j	0.8440	0.8130
k	0.8520	0.8280
l	0.8650	0.8260
m	0.9700	0.8720
0	0.8140	0.8000
р	0.9430	0.8710
q	0.8040	0.7980
r	0.8830	0.8380
S	0.7480	0.7720
t	0.7670	0.7840
u	0.6710	0.7400
V	0.6830	0.7160
W	0.8070	0.8130
Х	0.5720	0.7230
У	0.8440	0.8110
Z	0.8700	0.8310



1.0

