



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 24, 2021 – 02:16 PM EST

PDB ID : 3COK  
Title : Crystal structure of PLK4 kinase  
Authors : Atwell, S.; Burley, S.K.; Houle, A.; Leon, B.; Pelletier, L.A.; Sauder, J.M.;  
New York SGX Research Center for Structural Genomics (NYSGXRC)  
Deposited on : 2008-03-28  
Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.16  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.16


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1487 (2.26-2.26)
Ramachandran outliers	138981	1449 (2.26-2.26)
Sidechain outliers	138945	1450 (2.26-2.26)
RSRZ outliers	127900	1356 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	278	
1	B	278	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	A	277	-	-	X	-
2	SO4	A	278	-	-	X	-

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3857 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine/threonine-protein kinase PLK4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	234	1834	1179	313	327	15	0	0	0
1	B	234	1854	1194	312	333	15	0	2	0

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	SER	-	expression tag	UNP O00444
A	0	LEU	-	expression tag	UNP O00444
A	275	GLU	-	expression tag	UNP O00444
A	276	GLY	-	expression tag	UNP O00444
B	-1	SER	-	expression tag	UNP O00444
B	0	LEU	-	expression tag	UNP O00444
B	275	GLU	-	expression tag	UNP O00444
B	276	GLY	-	expression tag	UNP O00444

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0

- Molecule 3 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (three-letter code: ANP) (formula:  $C_{10}H_{17}N_6O_{12}P_3$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	27	10	6	9	2	0	0
3	B	1	31	10	6	12	3	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	44	Total	O	0	0
			44	44		
4	B	47	Total	O	0	0
			47	47		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Serine/threonine-protein kinase PLK4



- Molecule 1: Serine/threonine-protein kinase PLK4



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	84.52Å 102.15Å 63.06Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.25 27.16 – 2.25	Depositor EDS
% Data completeness (in resolution range)	97.7 (30.00-2.25) 97.2 (27.16-2.25)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.10 (at 2.24Å)	Xtrriage
Refinement program	REFMAC 5.0	Depositor
R, $R_{free}$	0.211 , 0.261 0.208 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	33.3	Xtrriage
Anisotropy	0.509	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 42.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3857	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	31.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 34.29 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.9861e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ANP, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.58	0/1871	0.75	3/2526 (0.1%)
1	B	0.56	0/1900	0.79	6/2567 (0.2%)
All	All	0.57	0/3771	0.77	9/5093 (0.2%)

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	43	ASP	CB-CG-OD2	7.42	124.98	118.30
1	B	43	ASP	CB-CG-OD2	6.39	124.05	118.30
1	B	239	ASP	CB-CG-OD2	6.00	123.70	118.30
1	A	213	ASP	CB-CG-OD2	5.81	123.53	118.30
1	B	153	ASP	CB-CG-OD2	5.69	123.42	118.30
1	A	260	ASP	CB-CG-OD2	5.44	123.19	118.30
1	B	251	ASP	CB-CG-OD2	5.38	123.14	118.30
1	B	226	ASP	CB-CG-OD2	5.15	122.93	118.30
1	B	192	ASP	CB-CG-OD2	5.03	122.83	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1834	0	1818	30	0
1	B	1854	0	1834	27	0
2	A	10	0	0	5	0
2	B	10	0	0	1	0
3	A	27	0	12	1	0
3	B	31	0	13	1	0
4	A	44	0	0	2	0
4	B	47	0	0	1	0
All	All	3857	0	3677	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:144:THR:HG21	2:B:302:SO4:O4	1.73	0.89
1:A:33:THR:HG22	1:A:35:LEU:H	1.46	0.80
1:B:144:THR:HG23	1:B:146:ASN:H	1.49	0.78
1:A:144:THR:HG22	1:A:148:ASN:H	1.51	0.76
1:A:144:THR:HG21	2:A:277:SO4:O3	1.91	0.71
1:B:115:PHE:CD1	1:B:149:ILE:HD12	2.30	0.67
1:A:246:ARG:NH1	2:A:278:SO4:O3	2.29	0.65
1:B:138:LEU:HD13	1:B:199:MET:HB2	1.78	0.64
1:B:33:THR:CG2	1:B:35:LEU:HB3	2.27	0.64
1:B:123:MET:CE	1:B:192:ASP:HB3	2.27	0.64
1:A:21:SER:HB3	1:A:22:PHE:HD2	1.64	0.63
1:B:144:THR:HG22	1:B:148:ASN:H	1.65	0.61
1:A:0:LEU:HD11	1:B:0:LEU:HD11	1.86	0.57
1:A:205:ILE:CD1	1:A:230:PRO:HG3	2.36	0.55
1:A:118:GLN:NE2	1:A:149:ILE:H	2.05	0.54
1:A:67:LYS:HD2	1:B:64:CYS:O	2.08	0.54
1:A:144:THR:HG23	1:A:146:ASN:H	1.73	0.53
1:A:214:THR:O	1:A:214:THR:HG22	2.09	0.52
1:B:68:HIS:HB3	1:B:71:ILE:HD12	1.93	0.51
1:A:96:MET:O	1:A:96:MET:HE3	2.10	0.51
1:B:96:MET:HG3	1:B:138:LEU:HG	1.93	0.50
1:A:98:ARG:NH1	3:A:301:ANP:O3'	2.44	0.50
1:B:217:ASN:HB3	1:B:220:ASN:HA	1.93	0.49
1:B:36:GLU:HG2	1:B:90:MET:HG3	1.93	0.49
1:A:21:SER:HB3	1:A:22:PHE:CD2	2.46	0.49
1:B:123:MET:HE1	1:B:192:ASP:HB3	1.95	0.49

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:89:GLU:OE2	4:B:328:HOH:O	2.20	0.48
1:A:246:ARG:NH1	2:A:278:SO4:S	2.84	0.48
1:A:59:GLU:HG3	4:A:317:HOH:O	2.13	0.48
1:A:190:GLU:C	1:A:192:ASP:H	2.18	0.47
1:A:30:SER:OG	1:A:33:THR:HB	2.14	0.47
1:B:123:MET:HE3	1:B:192:ASP:HB3	1.97	0.47
1:A:136:LEU:HA	4:A:343:HOH:O	2.15	0.46
1:B:112:ALA:O	1:B:116:MET:HB2	2.15	0.46
1:A:215:VAL:O	1:A:215:VAL:HG12	2.16	0.46
1:B:15:ASN:HB2	1:B:27:ARG:HH12	1.82	0.45
1:A:205:ILE:HD11	1:A:230:PRO:HG2	1.98	0.45
1:A:205:ILE:HD11	1:A:230:PRO:CG	2.46	0.45
1:B:33:THR:HG23	1:B:35:LEU:HB3	1.99	0.45
1:A:115:PHE:CD2	1:A:149:ILE:HD13	2.53	0.44
1:A:239:ASP:O	1:A:243:GLN:HG3	2.17	0.44
1:B:18:GLY:HA3	3:B:301:ANP:H4'	1.99	0.44
1:B:116:MET:CE	1:B:200:PHE:HB2	2.48	0.44
1:B:116:MET:HE1	1:B:200:PHE:HB2	2.00	0.43
1:A:116:MET:O	1:A:120:ILE:HD12	2.19	0.42
1:B:115:PHE:HD1	1:B:149:ILE:HD12	1.83	0.42
1:A:144:THR:CG2	2:A:277:SO4:O3	2.66	0.42
1:A:205:ILE:CD1	1:A:230:PRO:CG	2.97	0.41
1:B:25:VAL:HG22	1:B:40:LYS:HG3	2.01	0.41
1:B:248[B]:ASN:HD21	1:B:250:ALA:HB3	1.86	0.41
1:A:247:ARG:N	2:A:278:SO4:O1	2.42	0.41
1:A:47:MET:HE1	1:A:56:VAL:HG11	2.03	0.41
1:B:205:ILE:HD11	1:B:230:PRO:HG2	2.03	0.41
1:A:42:ILE:HG21	1:A:47:MET:HE2	2.03	0.40
1:B:123:MET:HE2	1:B:255:LEU:HD21	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	228/278 (82%)	223 (98%)	3 (1%)	2 (1%)	17	14
1	B	230/278 (83%)	223 (97%)	4 (2%)	3 (1%)	12	8
All	All	458/556 (82%)	446 (97%)	7 (2%)	5 (1%)	14	10

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	135	ASP
1	A	191	SER
1	B	135	ASP
1	B	153	ASP
1	B	220	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	196/248 (79%)	181 (92%)	15 (8%)	13	11
1	B	201/248 (81%)	182 (90%)	19 (10%)	8	6
All	All	397/496 (80%)	363 (91%)	34 (9%)	10	9

All (34) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	0	LEU
1	A	21	SER
1	A	27	ARG
1	A	30	SER
1	A	33	THR
1	A	61	LYS
1	A	67	LYS
1	A	139	SER
1	A	144	THR
1	A	149	ILE
1	A	156	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	196	LEU
1	A	246	ARG
1	A	255	LEU
1	A	256	SER
1	B	0	LEU
1	B	33	THR
1	B	35	LEU
1	B	49	LYS
1	B	90	MET
1	B	104	VAL
1	B	105	LYS
1	B	126	LEU
1	B	138	LEU
1	B	144	THR
1	B	149	ILE
1	B	160	LEU
1	B	196	LEU
1	B	221	LYS
1	B	223	VAL
1	B	226	ASP
1	B	233	LEU
1	B	255	LEU
1	B	256	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	118	GLN
1	A	148	ASN
1	B	57	GLN
1	B	65	GLN
1	B	159	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	SO4	B	303	-	4,4,4	0.16	0	6,6,6	0.22	0
2	SO4	A	278	-	4,4,4	0.20	0	6,6,6	0.11	0
2	SO4	A	277	-	4,4,4	0.28	0	6,6,6	0.40	0
3	ANP	B	301	-	29,33,33	3.09	7 (24%)	31,52,52	1.75	8 (25%)
2	SO4	B	302	-	4,4,4	0.19	0	6,6,6	0.47	0
3	ANP	A	301	-	24,29,33	2.39	4 (16%)	25,45,52	1.58	4 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ANP	B	301	-	-	3/14/38/38	0/3/3/3
3	ANP	A	301	-	-	0/9/32/38	0/3/3/3

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	301	ANP	PG-O1G	9.12	1.60	1.46
3	B	301	ANP	PB-O1B	8.65	1.59	1.46
3	A	301	ANP	PB-O1B	8.39	1.59	1.46
3	B	301	ANP	C4-N3	6.26	1.44	1.35
3	A	301	ANP	C4-N3	6.09	1.44	1.35
3	B	301	ANP	PG-N3B	5.09	1.76	1.63
3	B	301	ANP	PB-N3B	4.87	1.76	1.63

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	301	ANP	PA-O1A	2.63	1.60	1.50
3	B	301	ANP	PA-O1A	2.42	1.59	1.50
3	A	301	ANP	C2-N3	2.28	1.35	1.32
3	B	301	ANP	C2-N3	2.01	1.35	1.32

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	301	ANP	N3-C2-N1	-4.46	121.70	128.68
3	A	301	ANP	PA-O3A-PB	-3.97	119.96	132.56
3	B	301	ANP	C4-C5-N7	-3.95	105.28	109.40
3	A	301	ANP	N3-C2-N1	-3.65	122.98	128.68
3	B	301	ANP	PA-O3A-PB	-3.41	120.60	132.62
3	B	301	ANP	O1B-PB-N3B	-2.85	107.57	111.77
3	B	301	ANP	C2-N1-C6	2.66	123.30	118.75
3	B	301	ANP	C3'-C2'-C1'	2.46	104.68	100.98
3	A	301	ANP	C2-N1-C6	2.22	122.55	118.75
3	A	301	ANP	C4-C5-N7	-2.20	107.11	109.40
3	B	301	ANP	O3A-PB-N3B	2.13	112.50	106.59
3	B	301	ANP	O3G-PG-O1G	-2.07	108.26	113.45

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	301	ANP	PB-N3B-PG-O1G
3	B	301	ANP	PG-N3B-PB-O1B
3	B	301	ANP	PG-N3B-PB-O3A


There are no ring outliers.


5 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	278	SO4	3	0
2	A	277	SO4	2	0
3	B	301	ANP	1	0
2	B	302	SO4	1	0
3	A	301	ANP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In

addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	234/278 (84%)	-0.03	7 (2%) 50 53	19, 30, 46, 57	0
1	B	234/278 (84%)	0.03	10 (4%) 35 37	18, 31, 48, 58	0
All	All	468/556 (84%)	0.00	17 (3%) 42 44	18, 30, 47, 58	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	161	LYS	4.3
1	B	225	ALA	3.7
1	B	217	ASN	3.7
1	A	161	LYS	3.3
1	B	265	SER	3.2
1	A	223	VAL	3.0
1	B	222	VAL	3.0
1	A	250	ALA	3.0
1	A	212	THR	2.5
1	B	249	PRO	2.4
1	B	250	ALA	2.4
1	B	15	ASN	2.4
1	A	160	LEU	2.2
1	A	135	ASP	2.2
1	B	132	LEU	2.2
1	B	22[A]	PHE	2.1
1	A	211	ASP	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	ANP	B	301	31/31	0.85	0.18	19,41,69,69	0
3	ANP	A	301	27/31	0.90	0.15	19,41,60,61	0
2	SO4	A	277	5/5	0.94	0.13	49,50,51,52	0
2	SO4	B	302	5/5	0.95	0.15	47,48,48,50	0
2	SO4	B	303	5/5	0.96	0.11	68,68,68,69	0
2	SO4	A	278	5/5	0.97	0.28	32,32,33,33	5

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.