



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 9, 2020 – 09:58 PM BST

PDB ID : 1CPO
Title : CHLOROPEROXIDASE
Authors : Sundaramoorthy, M.; Poulos, T.L.
Deposited on : 1996-02-10
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION




The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	299	 92% 7%
2	B	6	 100%
3	C	2	 100%
4	D	2	 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	XYS	B	4	X	-	-	-
2	XYS	B	6	X	-	-	-
7	XYS	A	748	X	-	-	-

2 Entry composition i


There are 10 unique types of molecules in this entry. The entry contains 2784 atoms, of which 1 is hydrogen and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CHLOROPEROXIDASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	299	2323	1473	1	384	459	6	0	1	0

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-alpha-D-xylopyranose-(1-6)-[alpha-D-xylopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	6	68	38	2	28	0	0	0

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	2	28	16	2	10	0	0	0

- Molecule 4 is an oligosaccharide called beta-L-arabinopyranose-(1-2)-alpha-D-mannopyranose.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
4	D	2	Total	C	O	0	0	0
			20	11	9			

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is alpha-D-mannopyranose (three-letter code: MAN) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 11 6 5	0	0
6	A	1	Total C O 11 6 5	0	0
6	A	1	Total C O 11 6 5	0	0
6	A	1	Total C O 11 6 5	0	0
6	A	1	Total C O 11 6 5	0	0
6	A	1	Total C O 11 6 5	0	0
6	A	1	Total C O 11 6 5	0	0
6	A	1	Total C O 11 6 5	0	0
6	A	1	Total C O 11 6 5	0	0

- Molecule 7 is alpha-D-xylopyranose (three-letter code: XYS) (formula: C₅H₁₀O₅).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total C O 9 5 4	0	0


- Molecule 8 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: CHLOROPEROXIDASE

Chain A:  92% 7%




- Molecule 2: alpha-D-mannopyranose-(1-3)-alpha-D-xylopyranose-(1-6)-[alpha-D-xylopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain B:  100%




- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C:  100%



- Molecule 4: beta-L-arabinopyranose-(1-2)-alpha-D-mannopyranose

Chain D:  100%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	58.77Å 71.36Å 91.73Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 1.90 49.49 – 1.89	Depositor EDS
% Data completeness (in resolution range)	(Not available) (8.00-1.90) 88.3 (49.49-1.89)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.52 (at 1.88Å)	Xtrriage
Refinement program	X-PLOR	Depositor
R, R_{free}	0.182 , 0.224 0.178 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	16.2	Xtrriage
Anisotropy	0.482	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 59.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2784	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: YYS, NAG, MN, ARB, BMA, HEM, PCA, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.37	0/2392	0.60	0/3280

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2322	1	2137	13	0
2	B	68	0	57	0	0
3	C	28	0	25	0	0
4	D	20	0	17	0	0
5	A	14	0	13	1	0
6	A	99	0	90	1	0
7	A	9	0	8	0	0
8	A	1	0	0	0	0
9	A	43	0	30	0	0
10	A	179	0	0	0	0
All	All	2783	1	2377	13	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (13) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:27:ALA:HB2	1:A:94:LEU:HB3	1.83	0.59
1:A:0:PCA:HB3	1:A:297:ASN:HD22	1.67	0.59
1:A:253:PRO:HG2	1:A:256:LEU:HD13	1.89	0.53
1:A:1:GLU:HG2	1:A:16:PRO:HB3	1.91	0.53
1:A:80:GLU:HA	1:A:85:SER:O	2.10	0.52
1:A:27:ALA:HB1	1:A:28:PRO:HD2	1.91	0.50
1:A:10:TYR:O	5:A:512:NAG:H82	2.16	0.45
1:A:253:PRO:HB3	6:A:751:MAN:O4	2.17	0.45
1:A:147:HIS:HA	1:A:205:VAL:O	2.15	0.45
1:A:1:GLU:H	1:A:297:ASN:ND2	2.16	0.43
1:A:219:PHE:HA	1:A:220:PRO:HD3	1.78	0.42
1:A:53:LEU:HA	1:A:53:LEU:HD12	1.95	0.41
1:A:240:ALA:O	1:A:244:VAL:HG23	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	298/299 (100%)	288 (97%)	10 (3%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	252/251 (100%)	247 (98%)	5 (2%)	55 51

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	53	LEU
1	A	72	LEU
1	A	103	PHE
1	A	223	LEU
1	A	281	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	202	ASN
1	A	284	ASN
1	A	290	GLN
1	A	297	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	PCA	A	0	1	7,8,9	1.98	1 (14%)	9,10,12	1.90	3 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PCA	A	0	1	-	0/0/11/13	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	0	PCA	CD-N	4.85	1.47	1.34

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	0	PCA	OE-CD-CG	-3.90	119.96	126.76
1	A	0	PCA	CG-CD-N	-2.27	102.52	108.39
1	A	0	PCA	OE-CD-N	-2.20	119.72	124.86

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	0	PCA	1	0

5.5 Carbohydrates [i](#)

10 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	B	1	1,2	14,14,15	0.50	0	17,19,21	1.29	1 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	B	2	2	14,14,15	0.56	0	17,19,21	1.04	1 (5%)
2	BMA	B	3	2	11,11,12	0.77	0	15,15,17	0.92	1 (6%)
2	XYS	B	4	2	9,9,10	1.35	1 (11%)	10,12,14	1.85	2 (20%)
2	MAN	B	5	2	11,11,12	1.19	1 (9%)	15,15,17	1.18	1 (6%)
2	XYS	B	6	2	9,9,10	1.67	1 (11%)	10,12,14	1.67	2 (20%)
3	NAG	C	1	1,3	14,14,15	0.99	1 (7%)	17,19,21	1.04	1 (5%)
3	NAG	C	2	3	14,14,15	0.66	0	17,19,21	1.13	1 (5%)
4	MAN	D	1	1,4	11,11,12	1.02	0	15,15,17	1.39	2 (13%)
4	ARB	D	2	4	9,9,10	0.91	0	10,12,14	2.54	5 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	0/6/23/26	0/1/1/1
2	BMA	B	3	2	-	0/2/19/22	0/1/1/1
2	XYS	B	4	2	1/1/3/4	-	0/1/1/1
2	MAN	B	5	2	-	0/2/19/22	0/1/1/1
2	XYS	B	6	2	1/1/3/4	-	0/1/1/1
3	NAG	C	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	C	2	3	-	0/6/23/26	0/1/1/1
4	MAN	D	1	1,4	-	1/2/19/22	0/1/1/1
4	ARB	D	2	4	-	-	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	6	XYS	C4-C3	3.76	1.58	1.52
2	B	5	MAN	C4-C5	2.59	1.58	1.53
2	B	4	XYS	C4-C3	2.31	1.55	1.52
3	C	1	NAG	C3-C2	2.04	1.56	1.52

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	4	XYS	C5-O5-C1	4.63	118.64	111.52
4	D	2	ARB	C5-O5-C1	4.35	118.22	111.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	6	XYS	C5-O5-C1	3.78	117.34	111.52
4	D	2	ARB	O2-C2-C1	3.62	116.56	109.15
3	C	1	NAG	C1-O5-C5	3.36	116.74	112.19
2	B	1	NAG	C1-C2-N2	3.30	116.12	110.49
4	D	1	MAN	O3-C3-C2	3.29	116.28	109.99
4	D	2	ARB	O3-C3-C2	-2.99	104.27	109.99
2	B	6	XYS	C1-C2-C3	2.84	113.15	109.67
2	B	5	MAN	C1-O5-C5	2.81	116.01	112.19
3	C	2	NAG	C1-O5-C5	2.81	116.00	112.19
4	D	2	ARB	O2-C2-C3	-2.68	104.76	110.14
4	D	2	ARB	C1-C2-C3	2.67	112.95	109.67
2	B	4	XYS	O2-C2-C1	2.45	114.16	109.15
2	B	3	BMA	C1-O5-C5	2.30	115.31	112.19
2	B	2	NAG	C1-O5-C5	2.29	115.29	112.19
4	D	1	MAN	C1-C2-C3	-2.17	107.00	109.67

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	B	4	XYS	C2
2	B	6	XYS	C2


All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	D	1	MAN	O5-C5-C6-O6


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 1 is monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	NAG	A	512	1	14,14,15	0.75	0	17,19,21	1.31	2 (11%)
9	HEM	A	396	1,8	27,50,50	1.59	5 (18%)	17,82,82	1.29	2 (11%)
6	MAN	A	738	1	11,11,12	0.70	0	15,15,17	1.04	0
6	MAN	A	742	1	11,11,12	0.82	0	15,15,17	1.65	1 (6%)
7	XYS	A	748	1	9,9,10	1.45	2 (22%)	10,12,14	1.80	3 (30%)
6	MAN	A	750	1	11,11,12	0.93	0	15,15,17	1.09	1 (6%)
6	MAN	A	751	1	11,11,12	0.97	0	15,15,17	1.26	2 (13%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	MAN	A	783	1	11,11,12	1.30	1 (9%)	15,15,17	1.70	4 (26%)
6	MAN	A	752	1	11,11,12	0.84	0	15,15,17	1.37	2 (13%)
6	MAN	A	793	1	11,11,12	0.89	0	15,15,17	1.08	1 (6%)
6	MAN	A	741	1	11,11,12	0.74	0	15,15,17	1.23	1 (6%)
6	MAN	A	739	1	11,11,12	0.80	0	15,15,17	1.36	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	A	512	1	-	2/6/23/26	0/1/1/1
9	HEM	A	396	1,8	-	0/6/54/54	-
6	MAN	A	738	1	-	0/2/19/22	0/1/1/1
6	MAN	A	742	1	-	2/2/19/22	0/1/1/1
7	XYS	A	748	1	1/1/3/4	-	0/1/1/1
6	MAN	A	750	1	-	1/2/19/22	0/1/1/1
6	MAN	A	751	1	-	2/2/19/22	0/1/1/1
6	MAN	A	783	1	-	2/2/19/22	0/1/1/1
6	MAN	A	752	1	-	1/2/19/22	0/1/1/1
6	MAN	A	793	1	-	2/2/19/22	0/1/1/1
6	MAN	A	741	1	-	0/2/19/22	0/1/1/1
6	MAN	A	739	1	-	1/2/19/22	0/1/1/1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
9	A	396	HEM	C3B-CAB	-3.75	1.40	1.47
9	A	396	HEM	C3C-CAC	-3.66	1.40	1.47
9	A	396	HEM	C3B-C2B	-3.34	1.35	1.40
6	A	783	MAN	C2-C3	3.03	1.57	1.52
9	A	396	HEM	C3C-C2C	-2.74	1.36	1.40
9	A	396	HEM	CBC-CAC	2.33	1.44	1.29
7	A	748	XYS	O5-C5	2.06	1.46	1.42
7	A	748	XYS	C1-C2	2.02	1.56	1.52

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	742	MAN	C1-O5-C5	5.47	119.61	112.19
6	A	741	MAN	C1-O5-C5	3.94	117.53	112.19
7	A	748	XYS	C1-C2-C3	3.50	113.97	109.67
6	A	750	MAN	C1-O5-C5	3.35	116.73	112.19
5	A	512	NAG	C1-O5-C5	3.31	116.67	112.19
6	A	783	MAN	C1-O5-C5	3.26	116.61	112.19
6	A	751	MAN	C1-O5-C5	2.98	116.23	112.19
6	A	739	MAN	C1-C2-C3	2.93	113.27	109.67
6	A	739	MAN	C1-O5-C5	2.89	116.11	112.19
5	A	512	NAG	C1-C2-N2	2.86	115.37	110.49
9	A	396	HEM	CMC-C2C-C3C	2.76	129.84	124.68
7	A	748	XYS	C5-O5-C1	2.75	115.76	111.52
6	A	752	MAN	C1-O5-C5	2.72	115.88	112.19
6	A	783	MAN	C1-C2-C3	2.48	112.71	109.67
6	A	783	MAN	O4-C4-C5	2.47	115.42	109.30
9	A	396	HEM	CBD-CAD-C3D	2.29	116.70	112.48
7	A	748	XYS	O2-C2-C3	-2.20	105.72	110.14
6	A	783	MAN	C3-C4-C5	-2.17	106.36	110.24
6	A	752	MAN	O2-C2-C1	2.14	113.52	109.15
6	A	793	MAN	O3-C3-C2	-2.06	106.05	109.99
6	A	751	MAN	C1-C2-C3	2.05	112.19	109.67

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
7	A	748	XYS	C2

All (13) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	A	742	MAN	O5-C5-C6-O6
6	A	783	MAN	O5-C5-C6-O6
6	A	793	MAN	O5-C5-C6-O6
6	A	793	MAN	C4-C5-C6-O6
5	A	512	NAG	O5-C5-C6-O6
6	A	783	MAN	C4-C5-C6-O6
6	A	750	MAN	O5-C5-C6-O6
6	A	751	MAN	C4-C5-C6-O6
6	A	751	MAN	O5-C5-C6-O6
6	A	742	MAN	C4-C5-C6-O6
5	A	512	NAG	C4-C5-C6-O6
6	A	739	MAN	C4-C5-C6-O6
6	A	752	MAN	O5-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	512	NAG	1	0
6	A	751	MAN	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	298/299 (99%)	-0.33	0 100 100	5, 14, 31, 40	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	PCA	A	0	8/9	0.94	0.09	13,14,17,17	0

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	XYS	B	4	9/10	0.75	0.13	43,45,47,50	0
2	XYS	B	6	9/10	0.75	0.24	48,50,51,55	0
4	ARB	D	2	9/10	0.76	0.21	34,38,41,43	0
2	MAN	B	5	11/12	0.80	0.22	46,52,54,57	0
4	MAN	D	1	11/12	0.81	0.17	37,41,47,49	0
2	BMA	B	3	11/12	0.83	0.15	36,37,40,43	0
3	NAG	C	2	14/15	0.84	0.15	31,39,44,46	0
2	NAG	B	2	14/15	0.92	0.10	15,22,31,34	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	NAG	B	1	14/15	0.95	0.10	13,18,21,21	0
3	NAG	C	1	14/15	0.97	0.08	7,11,17,20	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain D:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
7	XYS	A	748	9/10	0.69	0.27	49,53,55,57	0
6	MAN	A	783	11/12	0.69	0.25	45,53,55,55	0
6	MAN	A	750	11/12	0.78	0.15	42,46,47,51	0
6	MAN	A	793	11/12	0.79	0.18	40,46,48,52	0
5	NAG	A	512	14/15	0.80	0.14	41,46,50,51	0
6	MAN	A	752	11/12	0.82	0.14	31,33,38,43	0
6	MAN	A	738	11/12	0.85	0.14	27,36,42,43	0
6	MAN	A	742	11/12	0.88	0.14	33,38,42,42	0
6	MAN	A	751	11/12	0.91	0.13	35,39,42,44	0
6	MAN	A	741	11/12	0.95	0.08	12,13,18,19	0
6	MAN	A	739	11/12	0.95	0.10	19,21,23,26	0
9	HEM	A	396	43/43	0.98	0.08	3,6,10,21	0
8	MN	A	301	1/1	0.99	0.07	14,14,14,14	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.