

wwPDB X-ray Structure Validation Summary Report (i)

Aug 21, 2020 - 09:17 PM BST

PDB ID	:	4CQM
Title	:	Crystal structure of heterotetrameric human ketoacyl reductase complexed
		with NAD and NADP
Authors	:	Venkatesan, R.; SahTeli, S.K.; Awoniyi, L.O.; Jiang, G.; Prus, P.; Kastoniotis,
		A.J.; Hiltunen, J.K.; Wierenga, R.K.; Chen, Z.
Deposited on	:	2014-02-19
Resolution	:	2.34 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity Mogul Xtriage (Phenix) EDS buster-report Percentile statistics Refmac CCP4 Ideal geometry (proteins) Ideal geometry (DNA, RNA)	· · · · · · · · · · · · · · · · · · ·	4.02b-467 1.8.5 (274361), CSD as541be (2020) 1.13 2.13.1 1.1.7 (2018) 20191225.v01 (using entries in the PDB archive December 25th 2019) 5.8.0158 7.0.044 (Gargrove) Engh & Huber (2001) Parkinson et al. (1996)
Ideal geometry (DNA, RNA) Validation Pipeline (wwPDB-VP)	:	Parkinson et al. (1996) 2 13 1
vandadon i ipenne (wwi DD vi)		2.10.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.34 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries}, { m resolution\ range}({ m \AA}))$				
R_{free}	130704	$2096 \ (2.36-2.32)$				
Clashscore	141614	2193 (2.36-2.32)				
Ramachandran outliers	138981	2159 (2.36-2.32)				
Sidechain outliers	138945	2160 (2.36-2.32)				
RSRZ outliers	127900	2067 (2.36-2.32)				

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	А	261	% • 84%	11%	5%
1	Е	261	84%	11%	5%
1	Н	261	81%	14%	• 5%
1	Ι	261	83%	12%	5%
1	L	261	80%	14%	6%
1	М	261	84%	11%	5%



Contri	nucu jion	i previous	puyc		
Mol	Chain	${f Length}$	Quality of chain		
1	P	961	%	100/	50/
T	1	201	82%	12%	5%
2	В	244		1604	E04
2	D	244	80%	15%	5%
0	a	014			
2	C	244	81%	13%	• 5%
2			2%		
2	F.	244	80%	18%	•
			7%		
2	G	244	68% 21%	•	8%
			%		
2	J	244	81%	17%	•
			2%		
2	Κ	244	81%	13%	5%
			2%		
2	Ν	244	77%	18%	5%
			3%		
2	0	244	75%	21%	•
			%		
3	D	261	83%	12%	5%

Continued from previous page...



2 Entry composition (i)

There are 8 unique types of molecules in this entry. The entry contains 29356 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Δ	248	Total	С	Ν	Ο	\mathbf{S}	0	0	0
	A	240	1783	1106	319	347	11	0	0	0
1	1 E	248	Total	С	Ν	Ο	S	0	1	0
			1775	1102	315	346	12	0		0
1	Ц	240	Total	С	Ν	Ο	S	0	1	0
L T	11	249	1785	1106	322	346	11	0	L	
1	т	249	Total	С	Ν	Ο	S	0	1	0
L T	L		1802	1121	322	348	11	0	L L	
1	т	246	Total	С	Ν	Ο	S	0	1	0
L T		240	1786	1109	322	344	11	0	L	
1	М	240	Total	С	Ν	Ο	S	0	2	0
		249	1819	1126	330	351	12	0	2	0
1	1 D	949	Total	С	Ν	Ο	S	0	1	0
	1	240	1785	1110	317	347	11			U

• Molecule 1 is a protein called ESTRADIOL 17-BETA-DEHYDROGENASE 8.

• Molecule 2 is a protein called CARBONYL REDUCTASE FAMILY MEMBER 4.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
2	В	232	Total 1680	C 1053	N 315	O 303	${f S}$ 9	0	0	0
2	С	232	Total 1667	C 1051	N 300	O 308	S 8	0	1	0
2	F	241	Total 1767	C 1107	N 327	O 323	S 10	0	1	0
2	G	224	Total 1628	C 1022	N 298	O 300	S 8	0	0	0
2	J	240	Total 1727	C 1085	N 318	0 314	S 10	0	0	0
2	K	231	Total 1649	C 1036	N 305	O 300	S 8	0	0	0
2	Ν	232	Total 1727	C 1083	N 327	O 308	S 9	0	3	0



Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
9	2 O 235	225	Total	С	Ν	Ο	S	0	0	0
		235	1640	1030	298	304	8	0	0	0

There are 56 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	-6	MET	-	expression tag	UNP Q8N4T8
В	-5	HIS	-	expression tag	UNP Q8N4T8
В	-4	HIS	-	expression tag	UNP Q8N4T8
В	-3	HIS	-	expression tag	UNP Q8N4T8
В	-2	HIS	-	expression tag	UNP Q8N4T8
В	-1	HIS	-	expression tag	UNP Q8N4T8
В	0	HIS	-	expression tag	UNP Q8N4T8
С	-6	MET	-	expression tag	UNP Q8N4T8
С	-5	HIS	-	expression tag	UNP Q8N4T8
С	-4	HIS	-	expression tag	UNP Q8N4T8
С	-3	HIS	-	expression tag	UNP Q8N4T8
С	-2	HIS	-	expression tag	UNP Q8N4T8
С	-1	HIS	-	expression tag	UNP Q8N4T8
С	0	HIS	-	expression tag	UNP Q8N4T8
F	-6	MET	-	expression tag	UNP Q8N4T8
F	-5	HIS	-	expression tag	UNP Q8N4T8
F	-4	HIS	-	expression tag	UNP Q8N4T8
F	-3	HIS	-	expression tag	UNP Q8N4T8
F	-2	HIS	-	expression tag	UNP Q8N4T8
F	-1	HIS	-	expression tag	UNP Q8N4T8
F	0	HIS	-	expression tag	UNP Q8N4T8
G	-6	MET	-	expression tag	UNP Q8N4T8
G	-5	HIS	-	expression tag	UNP Q8N4T8
G	-4	HIS	-	expression tag	UNP Q8N4T8
G	-3	HIS	-	expression tag	UNP Q8N4T8
G	-2	HIS	-	expression tag	UNP Q8N4T8
G	-1	HIS	-	expression tag	UNP Q8N4T8
G	0	HIS	-	expression tag	UNP Q8N4T8
J	-6	MET	-	expression tag	UNP Q8N4T8
J	-5	HIS	-	expression tag	UNP Q8N4T8
J	-4	HIS	-	expression tag	UNP Q8N4T8
J	-3	HIS	_	expression tag	UNP Q8N4T8
J	-2	HIS	-	expression tag	UNP Q8N4T8
J	-1	HIS	-	expression tag	UNP Q8N4T8
J	0	HIS	-	expression tag	UNP Q8N4T8
K	-6	MET	-	expression tag	UNP Q8N4T8



Chain	Besidue	Modelled	Actual	Comment	Beference
U V	F	IIIC			
N	-0	HIS	-	expression tag	UNP Q8N418
K	-4	HIS	-	expression tag	UNP Q8N4T8
K	-3	HIS	-	expression tag	UNP Q8N4T8
K	-2	HIS	-	expression tag	UNP Q8N4T8
K	-1	HIS	-	expression tag	UNP Q8N4T8
K	0	HIS	-	expression tag	UNP Q8N4T8
N	-6	MET	-	expression tag	UNP Q8N4T8
N	-5	HIS	-	expression tag	UNP Q8N4T8
N	-4	HIS	-	expression tag	UNP Q8N4T8
N	-3	HIS	-	expression tag	UNP Q8N4T8
N	-2	HIS	-	expression tag	UNP Q8N4T8
N	-1	HIS	-	expression tag	UNP Q8N4T8
N	0	HIS	-	expression tag	UNP Q8N4T8
0	-6	MET	-	expression tag	UNP Q8N4T8
0	-5	HIS	-	expression tag	UNP Q8N4T8
0	-4	HIS	-	expression tag	UNP Q8N4T8
0	-3	HIS	-	expression tag	UNP Q8N4T8
0	-2	HIS	-	expression tag	UNP Q8N4T8
0	-1	HIS	-	expression tag	UNP Q8N4T8
0	0	HIS	-	expression tag	UNP Q8N4T8

Continued from previous page...

• Molecule 3 is a protein called ESTRADIOL 17-BETA-DEHYDROGENASE 8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	248	Total 1772	C 1101	N 310	O 350	S 11	0	1	0

• Molecule 4 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula: C₂₁H₂₈N₇O₁₇P₃).





Mol	Chain	Residues		Ate	oms			ZeroOcc	AltConf
4	٨	1	Total	С	Ν	Ο	Р	0	0
4	A	L	44	21	$\overline{7}$	14	2	0	0
4	D	1	Total	С	Ν	Ο	Р	0	0
4	D	L	48	21	7	17	3	0	0
4	С	1	Total	С	Ν	Ο	Р	0	0
4	U	T	32	11	5	13	3	0	0
4	р	1	Total	С	Ν	Ο	Р	0	0
4	D	T	44	21	7	14	2	0	0
4	F	1	Total	С	Ν	Ο	Р	0	0
4		T	44	21	7	14	2	0	0
4	F	1	Total	С	Ν	Ο	Р	0	0
4	Ľ	T	48	21	7	17	3	0	0
4	C	1	Total	С	Ν	Ο	Р	0	0
4	G	T	27	10	5	10	2	0	0
4	н	1	Total	С	Ν	Ο	Р	0	0
-1	11	T	44	21	7	14	2	0	0
4	T	1	Total	С	Ν	Ο	Р	0	0
-1	L	T	44	21	7	14	2	0	0
4	Т	1	Total	С	Ν	Ο	Р	0	0
-1	J	T	48	21	7	17	3	0	0
4	K	1	Total	С	Ν	Ο	Р	0	0
-1	17	T	39	15	5	16	3	0	0
4	T.	1	Total	С	Ν	Ο	Р	0	0
-1	Ľ	T	44	21	7	14	2	0	0
	M	1	Total	$\overline{\mathrm{C}}$	N	Ō	Р	0	0
	IVI	L	44	21	7	14	2	0	0
	N	1	Total	С	N	Ο	Р	0	0
- ±	L 1 1	1	48	21	$\overline{7}$	17	3	U	



a 1	e		
Continued	from	previous	page
	9	1	1 5

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
4	0	1	Total	С	Ν	Ο	Р	0	0
4	4 0	L	48	21	7	17	3	0	0
4	D	1	Total	С	Ν	Ο	Р	0	0
4	Г	L	44	21	$\overline{7}$	14	2	0	0

• Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 6 3 3 \end{array}$	0	0
5	Е	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 6 3 3 \end{array}$	0	0
5	Ι	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 6 3 3 \end{array}$	0	0
5	М	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 6 & 3 & 3 \end{array}$	0	0

• Molecule 6 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	D	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 4 2 2 \end{array}$	0	0
6	Е	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 4 2 2 \end{array}$	0	0
6	F	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 4 2 2 \end{array}$	0	0
6	Ι	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 4 2 2 \end{array}$	0	0
6	Ι	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 4 2 2 \end{array}$	0	0
6	L	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 4 2 2 \end{array}$	0	0
6	М	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 4 2 2 \end{array}$	0	0
6	Р	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 4 2 2 \end{array}$	0	0

• Molecule 7 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	Н	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
7	М	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 4 2 2 \end{array}$	0	0
7	Р	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0

• Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	А	63	Total O 63 63	0	0
8	В	36	Total O 36 36	0	0
8	С	46	Total O 46 46	0	0
8	D	40	Total O 40 40	0	0
8	Е	66	Total O 66 66	0	0
8	F	49	Total O 49 49	0	0
8	G	24	$\begin{array}{ccc} \text{Total} & \text{O} \\ 24 & 24 \end{array}$	0	0
8	Н	36	Total O 36 36	0	0
8	Ι	60	Total O 60 60	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	J	48	Total O 48 48	0	0
8	K	47	$\begin{array}{cc} \text{Total} & \text{O} \\ 47 & 47 \end{array}$	0	0
8	L	56	Total O 56 56	0	0
8	М	93	Total O 93 93	0	0
8	Ν	52	$\begin{array}{cc} \text{Total} & \text{O} \\ 52 & 52 \end{array}$	0	0
8	О	38	Total O 38 38	0	0
8	Р	52	$\begin{array}{ccc} \text{Total} & \text{O} \\ 52 & 52 \end{array}$	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Chain A: 84% 11% 5% GLY GLY GLY GLY SER SER LYS GLU GLY • Molecule 1: ESTRADIOL 17-BETA-DEHYDROGENASE 8 Chain E: 84% 5% 11% GLY GLY GLY GLY GLY GLV GLY GLY GLY GLY GLY GLY GLY GLY GLY • Molecule 1: ESTRADIOL 17-BETA-DEHYDROGENASE 8 Chain H: 81% 14% • 5% GLY GLY GLY GLY GLV GLV GLV GLV GLV GLY PRO PRO • Molecule 1: ESTRADIOL 17-BETA-DEHYDROGENASE 8
- Molecule 1: ESTRADIOL 17-BETA-DEHYDROGENASE 8







• Molecule 1: ESTRADIOL 17-BETA-DEHYDROGENASE 8







K187









4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	87.50Å 238.41Å 87.69Å	Deperitor
$\mathrm{a,b,c,\alpha,\beta,\gamma}$	90.00° 94.18° 90.00°	Depositor
$\mathbf{P}_{\text{assolution}}(\hat{\mathbf{A}})$	49.25 - 2.34	Depositor
Resolution (A)	49.25 - 2.34	EDS
% Data completeness	$98.6 \ (49.25 - 2.34)$	Depositor
(in resolution range)	90.6~(49.25-2.34)	EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$0.61~({ m at}~2.34{ m \AA})$	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
B B.	0.203 , 0.235	Depositor
n, n_{free}	0.215 , 0.249	DCC
R_{free} test set	7402 reflections $(4.99%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	35.9	Xtriage
Anisotropy	0.476	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.34 , 38.8	EDS
L-test for $twinning^2$	$< L > = 0.47, < L^2 > = 0.30$	Xtriage
Estimated twinning fraction	0.118 for l,-k,h	Xtriage
Reported twinning fraction	0.120 for L,-K,H	Depositor
Outliers	2 of 148308 reflections (0.001%)	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	29356	wwPDB-VP
Average B, all atoms $(Å^2)$	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.08% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NAP, OAS, EDO, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bond	lengths	Bond angles		
	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.20	0/1803	0.37	0/2446	
1	Ε	0.20	0/1798	0.38	0/2442	
1	Н	0.20	0/1808	0.37	0/2456	
1	Ι	0.22	0/1826	0.39	0/2476	
1	L	0.20	0/1809	0.36	0/2454	
1	М	0.21	0/1845	0.38	0/2500	
1	Р	0.21	0/1808	0.37	0/2454	
2	В	0.20	0/1704	0.36	0/2302	
2	С	0.21	0/1689	0.37	0/2281	
2	F	0.21	0/1794	0.37	0/2421	
2	G	0.22	0/1645	0.41	0/2218	
2	J	0.20	0/1749	0.37	0/2364	
2	Κ	0.20	0/1668	0.37	0/2251	
2	Ν	0.20	0/1759	0.37	0/2369	
2	0	0.22	0/1658	0.37	0/2246	
3	D	0.20	0/1784	0.37	0/2421	
All	All	0.21	0/28147	0.37	0/38101	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1783	0	1780	16	0
1	Е	1775	0	1770	18	0
1	Н	1785	0	1771	25	0
1	Ι	1802	0	1811	20	0
1	L	1786	0	1802	22	0
1	М	1819	0	1839	18	0
1	Р	1785	0	1784	24	0
2	В	1680	0	1663	21	0
2	С	1667	0	1666	22	0
2	F	1767	0	1768	27	0
2	G	1628	0	1634	38	0
2	J	1727	0	1725	26	0
2	K	1649	0	1643	20	0
2	N	1727	0	1757	29	0
2	0	1640	0	1591	38	0
3	D	1772	0	1751	22	0
4	A	44	0	25	1	0
4	В	48	0	25	3	0
4	С	32	0	11	0	0
4	D	44	0	25	2	0
4	E	44	0	25	4	0
4	F	48	0	25	2	0
4	G	27	0	11	3	0
4	H	44	0	25	4	0
4	Ι	44	0	25	3	0
4	J	48	0	25	3	0
4	K	39	0	18	0	0
4	L	44	0	25	2	0
4	М	44	0	25	2	0
4	N	48	0	25	4	0
4	0	48	0	25	4	0
4	P	44	0	25	3	0
5	A	6	0	8	0	0
5	E	6	0	8	2	0
5	1	6	0	8	0	0
5	M	6	0	8	1	0
6	D	4	0	6	1	0
6	E	4	0	6	0	0
6	F	4	0	6	0	0
6	1	8	0	12	0	0
6	L	4	0	6	0	0
6	M	4	0	6	0	0
6	P	4	0	6	2	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	Н	4	0	3	0	0
7	М	4	0	3	0	0
7	Р	4	0	3	0	0
8	А	63	0	0	2	0
8	В	36	0	0	2	0
8	С	46	0	0	2	0
8	D	40	0	0	0	0
8	Е	66	0	0	0	0
8	F	49	0	0	2	0
8	G	24	0	0	1	0
8	Н	36	0	0	0	0
8	Ι	60	0	0	3	0
8	J	48	0	0	0	0
8	Κ	47	0	0	2	0
8	L	56	0	0	2	0
8	М	93	0	0	3	0
8	Ν	52	0	0	2	0
8	0	38	0	0	3	0
8	Р	52	0	0	1	0
All	All	29356	0	28209	358	0

Continued from previous page...

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 358 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:185:ARG:HD3	3:D:164:VAL:HG23	1.67	0.77
2:N:150:ALA:HB2	2:O:158:PHE:HB2	1.67	0.75
1:L:80:ARG:HA	1:L:83:ARG:HE	1.56	0.71
2:G:32:ILE:HG12	2:G:53:PHE:HB2	1.72	0.71
2:K:105:HIS:HA	2:K:109:LEU:HB3	1.73	0.70

There are no symmetry-related clashes.



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	244/261~(94%)	236~(97%)	7(3%)	1 (0%)	34	38
1	Ε	245/261~(94%)	238~(97%)	7(3%)	0	100	100
1	Η	246/261~(94%)	237~(96%)	8 (3%)	1 (0%)	34	38
1	Ι	246/261~(94%)	238~(97%)	7(3%)	1 (0%)	34	38
1	L	243/261~(93%)	236~(97%)	5 (2%)	2(1%)	19	20
1	М	247/261~(95%)	239~(97%)	8 (3%)	0	100	100
1	Р	245/261~(94%)	231 (94%)	14 (6%)	0	100	100
2	В	228/244~(93%)	217~(95%)	11 (5%)	0	100	100
2	С	229/244~(94%)	223~(97%)	6 (3%)	0	100	100
2	F	240/244~(98%)	228~(95%)	12~(5%)	0	100	100
2	G	218/244~(89%)	211 (97%)	7(3%)	0	100	100
2	J	238/244~(98%)	227~(95%)	11 (5%)	0	100	100
2	K	227/244~(93%)	218~(96%)	9 (4%)	0	100	100
2	Ν	231/244~(95%)	222~(96%)	9 (4%)	0	100	100
2	Ο	231/244 (95%)	224 (97%)	7 (3%)	0	100	100
3	D	$244/261 \ (94\%)$	237 (97%)	7 (3%)	0	100	100
All	All	3802/4040 (94%)	3662 (96%)	135 (4%)	5(0%)	51	62

All (5) Ramachandran outliers are listed below:

Mol	Chain	\mathbf{Res}	Type
1	А	203	ALA
1	L	203	ALA
1	Н	155	SER
1	Ι	155	SER
1	L	155	SER



5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	А	184/198~(93%)	181~(98%)	3~(2%)	62	74	
1	Ε	184/198~(93%)	182~(99%)	2(1%)	73	83	
1	Η	182/198~(92%)	179~(98%)	3~(2%)	62	74	
1	Ι	186/198~(94%)	182~(98%)	4 (2%)	52	63	
1	L	187/198~(94%)	185~(99%)	2(1%)	73	83	
1	М	192/198~(97%)	191~(100%)	1 (0%)	88	93	
1	Р	184/198~(93%)	181~(98%)	3 (2%)	62	74	
2	В	166/193~(86%)	162~(98%)	4 (2%)	49	59	
2	С	165/193~(86%)	160~(97%)	5(3%)	41	50	
2	F	178/193~(92%)	$172 \ (97\%)$	6 (3%)	37	46	
2	G	163/193~(84%)	152~(93%)	11 (7%)	16	17	
2	J	171/193~(89%)	170~(99%)	1 (1%)	86	92	
2	K	160/193~(83%)	156~(98%)	4 (2%)	47	58	
2	Ν	176/193~(91%)	175~(99%)	1 (1%)	86	92	
2	Ο	154/193~(80%)	153~(99%)	1 (1%)	86	92	
3	D	181/197~(92%)	179 (99%)	2 (1%)	73	83	
All	All	$2813/3127\ (90\%)$	2760 (98%)	53 (2%)	57	68	

5 of 53 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	G	21	LEU
2	G	70	LEU
2	Ν	235	LEU
2	G	28	ARG
2	G	34	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:



Mol	Chain	\mathbf{Res}	Type
2	G	50	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol T	Tune	Chain	Res	Link	Bond lengths			Bond angles		
	туре				Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	OAS	D	156	3	7,8,9	1.26	1 (14%)	$5,\!9,\!11$	1.25	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	OAS	D	156	3	-	0/5/7/9	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(A)	Ideal(Å)
3	D	156	OAS	OG-C1A	2.58	1.46	1.33

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
3	D	156	OAS	CB-OG-C1A	2.45	123.25	117.10

There are no chirality outliers.

There are no torsion outliers.



There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	156	OAS	2	0

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

31 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Tune	Chain	Dog	Tink	Bo	ond leng	ths	Bond angles		
	туре	Chain	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	EDO	Р	401	-	$3,\!3,\!3$	0.46	0	2,2,2	0.34	0
4	NAP	L	301	-	42,48,52	0.80	1(2%)	50,73,80	1.17	5(10%)
4	NAP	Р	301	-	42,48,52	0.81	1 (2%)	50,73,80	1.16	5(10%)
4	NAP	Ι	301	-	42,48,52	0.82	1 (2%)	50,73,80	1.17	5 (10%)
7	ACT	Н	501	-	$1,\!3,\!3$	1.36	0	0,3,3	0.00	-
7	ACT	М	501	-	$1,\!3,\!3$	1.26	0	0,3,3	0.00	-
6	EDO	F	401	-	$3,\!3,\!3$	0.46	0	2,2,2	0.33	0
7	ACT	Р	501	-	$1,\!3,\!3$	1.38	0	0,3,3	0.00	-
6	EDO	Е	402	-	3,3,3	0.47	0	2,2,2	0.31	0
4	NAP	E	301	-	42,48,52	0.80	1(2%)	50,73,80	1.20	5(10%)
6	EDO	D	401	-	$3,\!3,\!3$	0.46	0	2,2,2	0.35	0
4	NAP	Ο	301	-	45,52,52	0.81	1 (2%)	$56,\!80,\!80$	1.17	4 (7%)
4	NAP	F	301	-	45,52,52	0.80	1 (2%)	$56,\!80,\!80$	1.17	4 (7%)
4	NAP	J	301	-	45,52,52	0.81	1 (2%)	56,80,80	1.17	4 (7%)
4	NAP	Н	301	-	42,48,52	0.80	1 (2%)	50,73,80	1.17	5 (10%)
4	NAP	N	301	-	45,52,52	0.81	1 (2%)	56,80,80	1.17	4 (7%)
5	GOL	A	401	-	$5,\!5,\!5$	0.26	0	5, 5, 5	0.26	0
4	NAP	A	301	-	42,48,52	0.80	1 (2%)	50,73,80	1.17	5 (10%)



Mal	Tune	Chain	Dog	Link	Bo	ond leng	ths	B	ond ang	gles
	туре	Chain	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	EDO	М	402	-	$3,\!3,\!3$	0.46	0	2,2,2	0.32	0
4	NAP	М	301	-	42,48,52	0.81	1 (2%)	50,73,80	1.19	5 (10%)
5	GOL	Ι	401	-	$5,\!5,\!5$	0.25	0	5, 5, 5	0.27	0
4	NAP	D	301	-	42,48,52	0.80	1 (2%)	50,73,80	1.18	5 (10%)
6	EDO	L	401	-	$3,\!3,\!3$	0.48	0	2,2,2	0.30	0
5	GOL	Е	401	-	$5,\!5,\!5$	0.28	0	5, 5, 5	0.22	0
6	EDO	Ι	402	-	$3,\!3,\!3$	0.46	0	2,2,2	0.33	0
4	NAP	С	301	-	28,34,52	0.92	1(3%)	$34,\!53,\!80$	1.26	3 (8%)
4	NAP	G	301	-	25,29,52	0.98	1 (4%)	31,45,80	1.25	3 (9%)
4	NAP	K	301	-	36,42,52	0.84	1 (2%)	43,65,80	1.34	5 (11%)
4	NAP	В	301	-	45,52,52	0.82	1 (2%)	56,80,80	1.16	4 (7%)
5	GOL	М	401	-	5,5,5	0.27	0	5,5,5	0.33	0
6	EDO	Ι	403	-	$3,\!3,\!3$	0.45	0	2,2,2	0.33	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	EDO	P	401	-	-	0/1/1/1	-
4	NAP	L	301	-	-	$\frac{6/26/62/67}{}$	0/5/5/5
4	NAP	Р	301	-	-	18/26/62/67	0/5/5/5
4	NAP	Ι	301	-	-	10/26/62/67	0/5/5/5
6	EDO	F	401	-	-	0/1/1/1	-
6	EDO	Е	402	-	-	0/1/1/1	-
4	NAP	E	301	-	-	9/26/62/67	0/5/5/5
6	EDO	D	401	-	-	0/1/1/1	-
4	NAP	0	301	-	-	14/31/67/67	0/5/5/5
4	NAP	F	301	-	-	1/31/67/67	0/5/5/5
4	NAP	J	301	-	-	9/31/67/67	0/5/5/5
4	NAP	Н	301	-	-	10/26/62/67	0/5/5/5
4	NAP	N	301	-	-	8/31/67/67	0/5/5/5
5	GOL	А	401	-	-	2/4/4/4	-
4	NAP	А	301	-	-	9/26/62/67	0/5/5/5
6	EDO	М	402	-	-	0/1/1/1	-
4	NAP	М	301	-	-	9/26/62/67	0/5/5/5
5	GOL	Ι	401	-	-	4/4/4/4	_



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAP	D	301	-	-	13/26/62/67	0/5/5/5
6	EDO	L	401	-	-	0/1/1/1	-
5	GOL	Е	401	-	-	0/4/4/4	-
6	EDO	Ι	402	-	-	0/1/1/1	-
4	NAP	С	301	-	-	4/20/40/67	0/3/3/5
4	NAP	G	301	-	-	6/11/31/67	0/3/3/5
4	NAP	K	301	-	-	8/23/56/67	0/4/4/5
4	NAP	В	301	-	-	8/31/67/67	0/5/5/5
5	GOL	М	401	-	-	1/4/4/4	-
6	EDO	Ι	403	-	-	0/1/1/1	-

Continued from previous page...

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	$\operatorname{Ideal}(\operatorname{\AA})$
4	Ι	301	NAP	C5A-C4A	2.50	1.47	1.40
4	Е	301	NAP	C5A-C4A	2.50	1.47	1.40
4	А	301	NAP	C5A-C4A	2.50	1.47	1.40
4	В	301	NAP	C5A-C4A	2.48	1.47	1.40
4	К	301	NAP	C5A-C4A	2.48	1.47	1.40

The worst 5 of 71 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
4	J	301	NAP	PN-O3-PA	-4.00	119.11	132.83
4	М	301	NAP	PN-O3-PA	-3.54	120.66	132.83
4	В	301	NAP	C3D-C2D-C1D	3.47	106.20	100.98
4	Е	301	NAP	PN-O3-PA	-3.44	121.03	132.83
4	N	301	NAP	C3D-C2D-C1D	3.41	106.11	100.98

There are no chirality outliers.

5 of 149 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	L	301	NAP	C5D-O5D-PN-O3
4	L	301	NAP	O4D-C1D-N1N-C2N
4	Р	301	NAP	C5B-O5B-PA-O1A
4	Р	301	NAP	C5B-O5B-PA-O3
4	Р	301	NAP	C5D-O5D-PN-O1N

There are no ring outliers.



4	\cap	\cap	Мſ	
±	\cup	ખ	TAT	

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	Р	401	EDO	2	0
4	L	301	NAP	2	0
4	Р	301	NAP	3	0
4	Ι	301	NAP	3	0
4	Е	301	NAP	4	0
6	D	401	EDO	1	0
4	0	301	NAP	4	0
4	F	301	NAP	2	0
4	J	301	NAP	3	0
4	Н	301	NAP	4	0
4	N	301	NAP	4	0
4	А	301	NAP	1	0
4	М	301	NAP	2	0
4	D	301	NAP	2	0
5	Е	401	GOL	2	0
4	G	301	NAP	3	0
4	В	301	NAP	3	0
5	М	401	GOL	1	0

18 monomers are involved in 46 short contacts:

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.























































5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	# RSRZ > 2	$OWAB(Å^2)$	Q<0.9
1	А	248/261~(95%)	-0.12	2 (0%) 86 90	28, 44, 71, 124	0
1	E	248/261~(95%)	-0.24	0 100 100	28, 41, 64, 90	0
1	Η	249/261~(95%)	-0.09	1 (0%) 92 96	33, 52, 78, 90	0
1	Ι	249/261~(95%)	-0.16	1 (0%) 92 96	27, 44, 72, 111	0
1	L	246/261~(94%)	-0.23	1 (0%) 92 96	29, 44, 69, 92	0
1	М	249/261~(95%)	-0.23	1 (0%) 92 96	26, 40, 64, 103	0
1	Р	248/261~(95%)	-0.09	2 (0%) 86 90	31, 47, 72, 110	0
2	В	232/244~(95%)	-0.02	7 (3%) 50 60	30, 53, 87, 119	0
2	С	232/244~(95%)	0.13	4 (1%) 70 78	34, 54, 90, 130	0
2	F	241/244~(98%)	0.04	5 (2%) 63 73	30, 50, 88, 135	0
2	G	224/244~(91%)	0.53	16 (7%) 16 23	45, 73, 105, 123	0
2	J	240/244~(98%)	-0.01	3 (1%) 77 83	27, 51, 85, 118	0
2	Κ	231/244~(94%)	-0.01	4 (1%) 70 78	29, 49, 84, 127	0
2	N	232/244~(95%)	-0.08	4 (1%) 70 78	31, 45, 77, 102	0
2	Ο	$23\overline{5}/244~(96\%)$	0.25	8 (3%) 45 55	37, 62, 102, 136	0
3	D	$24\overline{7/261} \ (94\%)$	-0.15	2(0%) 86 90	31, 48, 74, 114	0
All	All	3851/4040 (95%)	-0.03	61 (1%) 72 80	26, 49, 86, 136	0

The worst 5 of 61 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	А	67	GLY	14.8
2	0	1	MET	9.3
2	G	51	LEU	6.0
2	F	194	LEU	5.2
2	В	47	GLY	4.9



6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B} ext{-factors}({ m \AA}^2)$	Q<0.9
3	OAS	D	156	9/10	0.85	0.17	$31,\!43,\!55,\!60$	0

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathrm{\AA}^2)$	Q < 0.9
6	EDO	L	401	4/4	0.73	0.16	$62,\!64,\!65,\!67$	0
5	GOL	Е	401	6/6	0.75	0.20	$62,\!66,\!71,\!82$	0
6	EDO	D	401	4/4	0.78	0.21	$60,\!60,\!62,\!63$	0
6	EDO	Р	401	4/4	0.87	0.22	$56,\!59,\!60,\!63$	0
4	NAP	0	301	48/48	0.88	0.19	$40,\!83,\!158,\!213$	0
4	NAP	G	301	27/48	0.88	0.14	56, 59, 105, 142	0
4	NAP	В	301	48/48	0.89	0.16	50,78,105,128	0
7	ACT	Н	501	4/4	0.90	0.15	$63,\!64,\!65,\!66$	0
4	NAP	N	301	48/48	0.90	0.23	$37,\!78,\!135,\!205$	0
5	GOL	А	401	6/6	0.92	0.16	$59,\!60,\!66,\!74$	0
4	NAP	С	301	32/48	0.92	0.14	47,58,147,147	0
6	EDO	М	402	4/4	0.92	0.19	$47,\!51,\!55,\!55$	0
4	NAP	J	301	48/48	0.92	0.14	$41,\!57,\!138,\!141$	0
6	EDO	Е	402	4/4	0.93	0.17	$47,\!49,\!51,\!52$	0
6	EDO	F	401	4/4	0.93	0.17	$47,\!54,\!58,\!61$	0
4	NAP	K	301	39/48	0.93	0.16	$35,\!49,\!160,\!161$	0
7	ACT	Р	501	4/4	0.93	0.20	$58,\!58,\!60,\!62$	0
5	GOL	M	401	6/6	0.93	0.24	47,51,57,70	0
6	EDO	Ι	403	4/4	0.93	0.13	$5\overline{4,}56,\!58,\!60$	0
7	ACT	М	501	4/4	0.94	0.13	$37,\!39,\!41,\!43$	0
4	NAP	Ĺ	301	44/48	0.94	0.14	40,50,61,209	0



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathrm{\AA}^2)$	Q<0.9
5	GOL	Ι	401	6/6	0.94	0.13	$52,\!59,\!66,\!67$	0
4	NAP	D	301	44/48	0.94	0.12	$35,\!52,\!61,\!82$	0
4	NAP	F	301	48/48	0.95	0.13	$33,\!48,\!80,\!127$	0
4	NAP	Р	301	44/48	0.96	0.12	$29,\!44,\!54,\!109$	0
4	NAP	М	301	44/48	0.96	0.13	$31,\!37,\!47,\!52$	0
4	NAP	Е	301	44/48	0.96	0.12	$24,\!39,\!48,\!54$	0
4	NAP	А	301	44/48	0.96	0.12	$34,\!43,\!48,\!55$	0
4	NAP	Н	301	44/48	0.97	0.10	$34,\!44,\!60,\!95$	0
4	NAP	Ι	301	44/48	0.97	0.11	$31,\!38,\!44,\!49$	0
6	EDO	Ι	402	4/4	0.97	0.15	34,40,45,47	0

Continued from previous page...

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







































6.5 Other polymers (i)

There are no such residues in this entry.

