



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 8, 2023 – 09:08 PM EDT

PDB ID : 8CRZ  
Title : Crystal structure of alien DNA CTSZZPBSBSZPPBAG in a tetragonal lattice  
Authors : Georgiadis, M.M.  
Deposited on : 2022-05-12  
Resolution : 1.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.32.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.32.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1223 (1.22-1.18)
Clashscore	141614	1286 (1.22-1.18)
RSRZ outliers	127900	1200 (1.22-1.18)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	8	
2	B	8	

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 416 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(\*CP\*TP\*(JSP)P\*(1W5)P\*(1W5)P\*(1WA)P\*(IGU)P\*(JSP))-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	A	8	164	79	27	51	7	0	0	0

- Molecule 2 is a DNA chain called DNA (5'-D(P\*(IGU)P\*(JSP)P\*(1W5)P\*(1WA)P\*(1WA)P\*(IGU)P\*AP\*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	B	8	173	80	36	49	8	0	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	39	Total	O	0	0
			39	39		
3	B	40	Total	O	0	0
			40	40		

### 3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA (5'-D(\*CP\*TP\*(JSP)P\*(1W5)P\*(1W5)P\*(1WA)P\*(IGU)P\*(JSP))-3')

Chain A: 



- Molecule 2: DNA (5'-D(P\*(IGU)P\*(JSP)P\*(1W5)P\*(1WA)P\*(1WA)P\*(IGU)P\*AP\*G)-3')

Chain B: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 42 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	46.50Å 46.50Å 42.45Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.79 – 1.20 25.99 – 1.20	Depositor EDS
% Data completeness (in resolution range)	99.6 (20.79-1.20) 99.6 (25.99-1.20)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	12.82 (at 1.20Å)	Xtrriage
Refinement program	PHENIX 1.13_2998	Depositor
R, $R_{free}$	0.184 , 0.200 0.190 , 0.204	Depositor DCC
$R_{free}$ test set	733 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	14.6	Xtrriage
Anisotropy	0.312	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.42 , 43.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	416	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	17.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.71% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 1WA, 1W5, JSP, IGU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.71	0/39	0.91	0/58
2	B	0.90	0/48	0.76	0/72
All	All	0.82	0/87	0.83	0/130

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	164	0	68	1	0
2	B	173	0	79	0	0
3	A	39	0	0	0	0
3	B	40	0	0	0	0
All	All	416	0	147	1	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (1) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:5:1W5:H2'	1:A:6:1WA:H8	2.03	0.41

There are no symmetry-related clashes.

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

#### 5.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

#### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

12 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	1W5	A	5	1	18,23,24	1.62	5 (27%)	20,33,36	2.28	5 (25%)
1	IGU	A	7	1	21,24,25	2.15	7 (33%)	30,35,38	1.24	3 (10%)
1	1W5	A	4	1	18,23,24	1.75	6 (33%)	20,33,36	2.31	5 (25%)
1	JSP	A	8	2,1	17,21,22	1.73	4 (23%)	20,30,33	0.75	1 (5%)
2	IGU	B	9	2,1	21,24,25	2.09	7 (33%)	30,35,38	1.26	3 (10%)
1	JSP	A	3	1	17,21,22	1.92	4 (23%)	20,30,33	0.96	1 (5%)
2	1WA	B	12	2	17,24,25	0.86	1 (5%)	19,35,38	1.15	2 (10%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	1WA	A	6	1	17,24,25	0.88	1 (5%)	19,35,38	1.02	1 (5%)
2	JSP	B	10	2	17,21,22	1.70	3 (17%)	20,30,33	0.92	0
2	1WA	B	13	2	17,24,25	0.95	1 (5%)	19,35,38	1.11	2 (10%)
2	IGU	B	14	2	21,24,25	2.50	9 (42%)	30,35,38	1.40	8 (26%)
2	1W5	B	11	2	18,23,24	1.60	4 (22%)	20,33,36	2.48	7 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	1W5	A	5	1	-	0/8/25/26	0/2/2/2
1	IGU	A	7	1	-	0/7/21/22	0/3/3/3
1	1W5	A	4	1	-	0/8/25/26	0/2/2/2
1	JSP	A	8	2,1	-	0/5/21/22	0/2/2/2
2	IGU	B	9	2,1	-	0/7/21/22	0/3/3/3
1	JSP	A	3	1	-	0/5/21/22	0/2/2/2
2	1WA	B	12	2	-	0/3/21/22	0/3/3/3
1	1WA	A	6	1	-	0/3/21/22	0/3/3/3
2	JSP	B	10	2	-	0/5/21/22	0/2/2/2
2	1WA	B	13	2	-	0/3/21/22	0/3/3/3
2	IGU	B	14	2	-	0/7/21/22	0/3/3/3
2	1W5	B	11	2	-	0/8/25/26	0/2/2/2

The worst 5 of 52 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	14	IGU	C5-N7	-5.68	1.28	1.39
1	A	7	IGU	C5-N7	-5.19	1.29	1.39
1	A	8	JSP	C1'-C1	5.01	1.57	1.50
2	B	9	IGU	C5-N7	-4.73	1.29	1.39
1	A	3	JSP	C1'-C1	4.72	1.57	1.50

The worst 5 of 38 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	11	1W5	C4-N3-C2	8.45	127.06	116.76
1	A	5	1W5	C4-N3-C2	8.03	126.55	116.76
1	A	4	1W5	C4-N3-C2	7.89	126.38	116.76
2	B	9	IGU	O4'-C1'-N9	3.97	114.97	107.86

Continued on next page...



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	11	1W5	C2'-C1'-C1	-3.86	109.19	114.72

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	5	1W5	1	0
1	A	6	1WA	1	0

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	2/8 (25%)	0.23	0 100 100	16, 16, 16, 17	0
2	B	2/8 (25%)	0.75	0 100 100	16, 16, 16, 18	0
All	All	4/16 (25%)	0.49	0 100 100	16, 16, 17, 18	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	IGU	B	14	22/23	0.90	0.11	14,17,21,25	0
1	JSP	A	3	20/21	0.91	0.10	14,17,20,22	0
1	1W5	A	4	22/23	0.94	0.09	12,16,20,29	0
2	1WA	B	13	22/23	0.95	0.08	12,15,19,22	0
2	1W5	B	11	22/23	0.95	0.09	12,16,21,29	0
1	1W5	A	5	22/23	0.96	0.09	12,14,18,29	0
2	1WA	B	12	22/23	0.96	0.07	11,14,19,23	0
1	IGU	A	7	22/23	0.96	0.09	11,12,16,19	0
2	IGU	B	9	22/23	0.96	0.09	10,12,15,16	0
1	1WA	A	6	22/23	0.97	0.09	11,12,15,16	0
2	JSP	B	10	20/21	0.97	0.08	11,15,18,19	0
1	JSP	A	8	20/21	0.97	0.09	10,12,17,18	0

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.