



Full wwPDB X-ray Structure Validation Report ⓘ

May 14, 2020 – 04:56 pm BST

PDB ID : 5CS6
Title : Crystal Structure of CK2alpha with Compound 3 bound
Authors : Brear, P.; De Fusco, C.; Georgiou, K.H.; Spring, D.; Hyvonen, M.
Deposited on : 2015-07-23
Resolution : 1.88 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.88 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	9470 (1.90-1.86)
Clashscore	141614	10282 (1.90-1.86)
Ramachandran outliers	138981	10152 (1.90-1.86)
Sidechain outliers	138945	10152 (1.90-1.86)
RSRZ outliers	127900	9303 (1.90-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	352	 3% 87% 6% 7%
1	B	352	 25% 81% 8% 10%

2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 5795 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Casein kinase II subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	327	2813	1801	493	507	12	0	7	0
1	B	316	2684	1719	473	481	11	0	2	0

There are 50 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-22	GLY	-	expression tag	UNP P68400
A	-21	SER	-	expression tag	UNP P68400
A	-20	MET	-	expression tag	UNP P68400
A	-19	ASP	-	expression tag	UNP P68400
A	-18	ILE	-	expression tag	UNP P68400
A	-17	GLU	-	expression tag	UNP P68400
A	-16	PHE	-	expression tag	UNP P68400
A	-15	ASP	-	expression tag	UNP P68400
A	-14	ASP	-	expression tag	UNP P68400
A	-13	ASP	-	expression tag	UNP P68400
A	-12	ALA	-	expression tag	UNP P68400
A	-11	ASP	-	expression tag	UNP P68400
A	-10	ASP	-	expression tag	UNP P68400
A	-9	ASP	-	expression tag	UNP P68400
A	-8	GLY	-	expression tag	UNP P68400
A	-7	SER	-	expression tag	UNP P68400
A	-6	GLY	-	expression tag	UNP P68400
A	-5	SER	-	expression tag	UNP P68400
A	-4	GLY	-	expression tag	UNP P68400
A	-3	SER	-	expression tag	UNP P68400
A	-2	GLY	-	expression tag	UNP P68400
A	-1	SER	-	expression tag	UNP P68400
A	0	GLY	-	expression tag	UNP P68400
A	1	SER	-	expression tag	UNP P68400
A	21	SER	ARG	engineered mutation	UNP P68400

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-22	GLY	-	expression tag	UNP P68400
B	-21	SER	-	expression tag	UNP P68400
B	-20	MET	-	expression tag	UNP P68400
B	-19	ASP	-	expression tag	UNP P68400
B	-18	ILE	-	expression tag	UNP P68400
B	-17	GLU	-	expression tag	UNP P68400
B	-16	PHE	-	expression tag	UNP P68400
B	-15	ASP	-	expression tag	UNP P68400
B	-14	ASP	-	expression tag	UNP P68400
B	-13	ASP	-	expression tag	UNP P68400
B	-12	ALA	-	expression tag	UNP P68400
B	-11	ASP	-	expression tag	UNP P68400
B	-10	ASP	-	expression tag	UNP P68400
B	-9	ASP	-	expression tag	UNP P68400
B	-8	GLY	-	expression tag	UNP P68400
B	-7	SER	-	expression tag	UNP P68400
B	-6	GLY	-	expression tag	UNP P68400
B	-5	SER	-	expression tag	UNP P68400
B	-4	GLY	-	expression tag	UNP P68400
B	-3	SER	-	expression tag	UNP P68400
B	-2	GLY	-	expression tag	UNP P68400
B	-1	SER	-	expression tag	UNP P68400
B	0	GLY	-	expression tag	UNP P68400
B	1	SER	-	expression tag	UNP P68400
B	21	SER	ARG	engineered mutation	UNP P68400

- Molecule 2 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is 1-(3-chloro-4-propoxyphenyl)methanamine (three-letter code: K82) (formula: C₁₀H₁₄ClNO).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	Cl	N	O	0	0
			13	10	1	1	1		
3	A	1	Total	C	Cl	N	O	0	0
			13	10	1	1	1		
3	A	1	Total	C	Cl	N	O	0	0
			13	10	1	1	1		
3	A	1	Total	C	Cl	N	O	0	0
			13	10	1	1	1		

- Molecule 4 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O P 5 4 1	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	198	Total O 198 198	0	0
5	B	27	Total O 27 27	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	64.50Å 68.85Å 335.87Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	167.94 – 1.88 167.94 – 1.88	Depositor EDS
% Data completeness (in resolution range)	99.2 (167.94-1.88) 99.9 (167.94-1.88)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.43 (at 1.88Å)	Xtrriage
Refinement program	BUSTER-TNT	Depositor
R, R_{free}	0.206 , 0.216 0.211 , 0.222	Depositor DCC
R_{free} test set	3111 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	35.5	Xtrriage
Anisotropy	0.564	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 56.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5795	wwPDB-VP
Average B, all atoms (Å ²)	77.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, K82, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.63	0/2888	0.63	0/3905
1	B	0.47	0/2756	0.62	0/3726
All	All	0.55	0/5644	0.63	0/7631

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2813	0	2751	18	0
1	B	2684	0	2627	23	0
2	A	12	0	9	0	0
2	B	4	0	3	0	0
3	A	52	0	0	1	0
4	A	5	0	0	0	0
5	A	198	0	0	4	1
5	B	27	0	0	0	0
All	All	5795	0	5390	39	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:35:ASN:OD1	1:B:37:ASP:HB2	1.74	0.85
1:A:296:GLU:HG3	3:A:407:K82:C2	2.27	0.64
1:B:255:TYR:HA	1:B:258:ILE:HG22	1.80	0.64
1:B:229:LYS:NZ	1:B:244[B]:ARG:NH1	2.50	0.59
1:B:84:ILE:HG23	1:B:152:ILE:HD13	1.84	0.59
1:A:236:HIS:HE1	5:A:655:HOH:O	1.87	0.58
1:A:72:PRO:O	1:A:73:VAL:HG23	2.04	0.57
1:A:286:HIS:HE1	5:A:671:HOH:O	1.88	0.56
1:A:50:TYR:HD1	1:A:73:VAL:HG21	1.72	0.55
1:B:258:ILE:HG13	1:B:263:ILE:HB	1.92	0.52
1:B:229:LYS:HZ1	1:B:244[B]:ARG:HH12	1.60	0.49
1:B:229:LYS:HZ2	1:B:244[B]:ARG:HH11	1.61	0.49
1:B:8:ARG:HH11	1:B:13:THR:HG21	1.77	0.49
1:A:50:TYR:CD1	1:A:73:VAL:HG21	2.49	0.48
1:A:285:VAL:HG22	1:A:293:VAL:HG11	1.95	0.48
1:B:229:LYS:HZ2	1:B:244[B]:ARG:NH1	2.11	0.48
1:A:70:LEU:HD13	1:A:78:ILE:HG12	1.96	0.48
1:B:96:THR:HB	1:B:114:GLU:HG3	1.96	0.48
1:A:118:ASN:HD22	1:A:164:ILE:H	1.61	0.48
1:B:54:PHE:CE2	1:B:69:ILE:HD12	2.49	0.47
1:B:229:LYS:HZ1	1:B:244[B]:ARG:NH1	2.13	0.46
1:B:96:THR:HB	1:B:114:GLU:CG	2.45	0.46
1:B:23:TYR:O	1:B:83:LYS:HE3	2.16	0.46
1:A:73:VAL:CG1	1:A:74:LYS:N	2.80	0.45
1:A:33:TRP:CE3	1:A:100:ILE:HG22	2.52	0.44
1:A:236:HIS:CE1	5:A:655:HOH:O	2.68	0.44
1:A:26:TYR:HE1	1:A:76:LYS:HG2	1.83	0.44
1:B:33:TRP:CZ3	1:B:100:ILE:HG22	2.53	0.43
1:B:165:ASP:HB3	1:B:170:LYS:HB3	2.01	0.43
1:A:90:GLY:HA3	1:B:33:TRP:HD1	1.83	0.43
1:A:118:ASN:ND2	1:A:164:ILE:H	2.17	0.42
1:A:225[A]:MET:HB2	1:A:225[A]:MET:HE3	1.82	0.42
1:B:190:VAL:HG11	1:B:205:ASP:HA	2.00	0.42
1:B:100:ILE:HG23	1:B:111:LEU:HD23	2.00	0.42
1:A:183:HIS:HB2	1:A:186:GLN:HE21	1.85	0.42
1:A:23:TYR:CE1	1:B:104:PRO:HB3	2.55	0.41
1:B:183:HIS:HB2	1:B:186:GLN:HE21	1.86	0.41
5:A:539:HOH:O	1:B:107:ARG:HD3	2.20	0.40
1:B:240:ASP:O	1:B:244[A]:ARG:HG2	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:681:HOH:O	5:A:681:HOH:O[4_597]	1.07	1.13

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	332/352 (94%)	320 (96%)	12 (4%)	0	100	100
1	B	314/352 (89%)	301 (96%)	13 (4%)	0	100	100
All	All	646/704 (92%)	621 (96%)	25 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	308/319 (97%)	308 (100%)	0	100	100
1	B	293/319 (92%)	291 (99%)	2 (1%)	84	83
All	All	601/638 (94%)	599 (100%)	2 (0%)	92	92

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	40	GLN
1	B	100	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	35	ASN
1	A	118	ASN
1	A	186	GLN
1	A	262	ASN
1	A	270	ASN
1	B	118	ASN
1	B	186	GLN
1	B	262	ASN
1	B	270	ASN
1	B	310	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	K82	A	405	-	13,13,13	0.17	0	16,16,16	0.32	0
4	PO4	A	408	-	4,4,4	2.55	1 (25%)	6,6,6	0.69	0
3	K82	A	406	-	13,13,13	0.16	0	16,16,16	0.39	0
3	K82	A	407	-	13,13,13	0.17	0	16,16,16	0.39	0
2	ACT	A	401	-	1,3,3	5.35	1 (100%)	0,3,3	0.00	-
2	ACT	A	402	-	1,3,3	8.36	1 (100%)	0,3,3	0.00	-
3	K82	A	404	-	13,13,13	0.16	0	16,16,16	0.39	0
2	ACT	B	401	-	1,3,3	4.61	1 (100%)	0,3,3	0.00	-
2	ACT	A	403	-	1,3,3	4.91	1 (100%)	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	K82	A	405	-	-	1/6/6/6	0/1/1/1
3	K82	A	406	-	-	0/6/6/6	0/1/1/1
3	K82	A	404	-	-	1/6/6/6	0/1/1/1
3	K82	A	407	-	-	1/6/6/6	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	402	ACT	CH3-C	8.36	1.59	1.48
2	A	401	ACT	CH3-C	5.35	1.55	1.48
2	A	403	ACT	CH3-C	4.91	1.55	1.48
2	B	401	ACT	CH3-C	4.61	1.54	1.48
4	A	408	PO4	P-O1	4.26	1.60	1.50

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	405	K82	C1-C2-O-C3
3	A	404	K82	C-C1-C2-O
3	A	407	K82	C5-C6-C7-N

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	407	K82	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	327/352 (92%)	0.12	12 (3%) 41 43	24, 38, 75, 105	0
1	B	316/352 (89%)	1.41	87 (27%) 0 0	46, 107, 200, 241	0
All	All	643/704 (91%)	0.75	99 (15%) 2 2	24, 69, 181, 241	0

All (99) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	125	TYR	8.2
1	B	257	TYR	7.7
1	B	50	TYR	6.9
1	B	33	TRP	6.2
1	B	281	TRP	6.2
1	B	327	VAL	5.7
1	B	5	VAL	5.5
1	B	43	ARG	5.2
1	B	42	VAL	5.2
1	B	307	TYR	5.1
1	B	273	LEU	4.8
1	B	296	GLU	4.8
1	A	124	LEU	4.8
1	B	272	ILE	4.7
1	B	300	PHE	4.7
1	B	45	LEU	4.6
1	B	41	LEU	4.5
1	B	251	THR	4.4
1	A	50	TYR	4.2
1	B	117	ASN	4.2
1	A	105	VAL	4.2
1	B	269	PHE	4.2
1	B	298	LEU	4.1
1	B	261	TYR	4.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	182	TYR	4.1
1	B	283	ARG	4.0
1	B	275	ARG	3.9
1	B	326	THR	3.8
1	B	49	LYS	3.8
1	B	170	LYS	3.8
1	B	290	GLN	3.7
1	A	328	VAL	3.5
1	B	265	LEU	3.4
1	B	116	VAL	3.4
1	B	259	ASP	3.4
1	B	227	PHE	3.4
1	A	126	GLN	3.3
1	B	4	PRO	3.3
1	B	166	HIS	3.3
1	B	255	TYR	3.2
1	B	243	VAL	3.2
1	B	278	ARG	3.2
1	B	252	GLU	3.1
1	B	256	ASP	3.0
1	B	274	GLY	3.0
1	B	323	TYR	3.0
1	A	104	PRO	3.0
1	B	48	GLY	3.0
1	B	19	ARG	2.9
1	B	104	PRO	2.9
1	B	258	ILE	2.9
1	B	277	SER	2.9
1	B	72	PRO	2.9
1	B	46	GLY	2.8
1	A	73	VAL	2.8
1	B	262	ASN	2.8
1	B	280	ARG	2.8
1	B	279	LYS	2.7
1	B	276	HIS	2.7
1	B	249	LEU	2.7
1	A	49	LYS	2.6
1	B	181	PHE	2.5
1	B	107	ARG	2.5
1	A	121	PHE	2.5
1	B	305	LEU	2.5
1	B	47	ARG	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	229	LYS	2.4
1	B	131	TYR	2.4
1	B	264	GLU	2.4
1	B	318	ALA	2.4
1	A	33	TRP	2.4
1	B	263	ILE	2.4
1	B	114	GLU	2.3
1	B	254	LEU	2.3
1	B	287	SER	2.3
1	B	73	VAL	2.3
1	B	299	ASP	2.3
1	B	232	PHE	2.2
1	B	324	PHE	2.2
1	B	30	VAL	2.2
1	B	167	GLU	2.2
1	B	250	GLY	2.2
1	B	319	MET	2.2
1	B	245	ILE	2.2
1	B	289	ASN	2.2
1	B	308	ASP	2.2
1	B	105	VAL	2.2
1	B	292	LEU	2.1
1	B	284	PHE	2.1
1	A	117	ASN	2.1
1	B	311[A]	SER	2.1
1	B	271	ASP	2.1
1	B	36	GLN	2.1
1	B	321	HIS	2.1
1	B	206	TYR	2.0
1	B	64	LYS	2.0
1	B	28	SER	2.0
1	B	102	LYS	2.0
1	B	242	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ACT	B	401	4/4	0.68	0.25	85,86,87,88	0
3	K82	A	404	13/13	0.82	0.18	43,46,57,59	0
3	K82	A	405	13/13	0.82	0.20	48,58,63,63	0
2	ACT	A	403	4/4	0.83	0.14	76,77,77,77	0
2	ACT	A	402	4/4	0.85	0.14	45,46,46,47	0
4	PO4	A	408	5/5	0.87	0.15	70,70,75,78	0
3	K82	A	407	13/13	0.91	0.18	63,65,69,70	0
3	K82	A	406	13/13	0.92	0.20	59,65,72,73	0
2	ACT	A	401	4/4	0.93	0.12	48,57,57,58	0

6.5 Other polymers [i](#)

There are no such residues in this entry.