



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 31, 2023 – 11:44 AM JST

PDB ID : 5CTT
Title : Crystal structure of human SART3/TIP110 NLS-mouse importin alpha complex
Authors : Park, J.K.; Kim, E.E.
Deposited on : 2015-07-24
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	447	 5% (poor fit), 86% (0-1 outliers), 8% (2 outliers), 5% (3+ outliers)
2	B	55	 4% (poor fit), 25% (0-1 outliers), 71% (2+ outliers)

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3874 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Importin subunit alpha-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	426	3260	2075	555	620	10	0	2	0

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	51	MET	-	initiating methionine	UNP P52293
A	52	GLY	-	expression tag	UNP P52293
A	53	SER	-	expression tag	UNP P52293
A	54	SER	-	expression tag	UNP P52293
A	55	HIS	-	expression tag	UNP P52293
A	56	HIS	-	expression tag	UNP P52293
A	57	HIS	-	expression tag	UNP P52293
A	58	HIS	-	expression tag	UNP P52293
A	59	HIS	-	expression tag	UNP P52293
A	60	HIS	-	expression tag	UNP P52293
A	61	SER	-	expression tag	UNP P52293
A	62	SER	-	expression tag	UNP P52293
A	63	GLY	-	expression tag	UNP P52293
A	64	LEU	-	expression tag	UNP P52293
A	65	VAL	-	expression tag	UNP P52293
A	66	PRO	-	expression tag	UNP P52293
A	67	ARG	-	expression tag	UNP P52293
A	68	GLY	-	expression tag	UNP P52293
A	69	SER	-	expression tag	UNP P52293
A	70	HIS	-	expression tag	UNP P52293
A	71	MET	-	expression tag	UNP P52293

- Molecule 2 is a protein called Squamous cell carcinoma antigen recognized by T-cells 3.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	16	142	82	37	23	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	595	LEU	-	expression tag	UNP Q15020
B	596	VAL	-	expression tag	UNP Q15020
B	597	PRO	-	expression tag	UNP Q15020
B	598	ARG	-	expression tag	UNP Q15020
B	599	GLY	-	expression tag	UNP Q15020
B	600	SER	-	expression tag	UNP Q15020

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	446	Total 446	O 446	0	0
3	B	26	Total 26	O 26	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	79.15Å 89.99Å 97.82Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.77 – 1.70 37.77 – 1.70	Depositor EDS
% Data completeness (in resolution range)	98.9 (37.77-1.70) 98.9 (37.77-1.70)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.85 (at 1.70Å)	Xtrriage
Refinement program	PHENIX 1.7.3_928	Depositor
R, R_{free}	0.191 , 0.219 0.195 , 0.196	Depositor DCC
R_{free} test set	3859 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	19.4	Xtrriage
Anisotropy	0.202	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 41.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3874	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.84% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.75	12/3321 (0.4%)	0.58	3/4526 (0.1%)
2	B	0.44	0/141	0.64	0/181
All	All	0.74	12/3462 (0.3%)	0.58	3/4707 (0.1%)

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	488	SER	CB-OG	-12.75	1.25	1.42
1	A	484	VAL	CB-CG1	-12.22	1.27	1.52
1	A	484	VAL	CB-CG2	-11.40	1.28	1.52
1	A	488	SER	CA-CB	-10.05	1.37	1.52
1	A	489	LEU	C-O	-8.96	1.06	1.23
1	A	488	SER	N-CA	-7.92	1.30	1.46
1	A	483	SER	C-O	-7.49	1.09	1.23
1	A	483	SER	CB-OG	-6.72	1.33	1.42
1	A	485	TYR	C-O	-6.58	1.10	1.23
1	A	487	ALA	C-O	-5.34	1.13	1.23
1	A	490	ASN	CB-CG	-5.29	1.38	1.51
1	A	483	SER	CA-CB	-5.16	1.45	1.52

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	488	SER	N-CA-CB	-7.25	99.63	110.50
1	A	485	TYR	N-CA-CB	5.51	120.51	110.60
1	A	489	LEU	CB-CG-CD2	-5.09	102.34	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3260	0	3338	34	0
2	B	142	0	153	3	0
3	A	446	0	0	12	2
3	B	26	0	0	3	0
All	All	3874	0	3491	37	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:348:LYS:NZ	3:A:503:HOH:O	1.95	0.97
1:A:485:TYR:HD2	1:A:486:LYS:N	1.72	0.87
1:A:88:ASN:ND2	3:A:504:HOH:O	2.06	0.87
1:A:72:GLY:N	3:A:506:HOH:O	2.10	0.84
1:A:247:ASP:OD1	3:A:505:HOH:O	2.09	0.70
1:A:287:GLU:OE2	1:A:291:LYS:HE3	1.92	0.69
1:A:386:LEU:HD21	1:A:425:LEU:HD13	1.75	0.69
1:A:485:TYR:CD2	1:A:486:LYS:N	2.61	0.69
2:B:639:GLU:OE1	3:B:701:HOH:O	2.12	0.67
2:B:639:GLU:OE2	3:B:702:HOH:O	2.12	0.66
1:A:123:LYS:HG2	3:A:524:HOH:O	2.02	0.59
1:A:477:GLN:OE1	1:A:489:LEU:HD22	2.02	0.59
1:A:485:TYR:CD2	1:A:485:TYR:C	2.75	0.58
1:A:485:TYR:HD2	1:A:485:TYR:C	2.08	0.56
1:A:73:THR:HA	1:A:76:TRP:NE1	2.23	0.53
1:A:172:ALA:HB2	3:A:809:HOH:O	2.08	0.53
1:A:366:ARG:NH2	3:A:507:HOH:O	2.14	0.52
1:A:458:GLU:H	1:A:458:GLU:CD	2.13	0.52
1:A:490:ASN:O	1:A:490:ASN:OD1	2.29	0.51
2:B:605:ARG:HG3	3:B:704:HOH:O	2.12	0.50
1:A:489:LEU:CD1	1:A:493:GLU:OE2	2.61	0.48
1:A:434:THR:OG1	3:A:508:HOH:O	2.19	0.48
1:A:213:LEU:O	1:A:258[B]:ARG:NH1	2.46	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:490:ASN:OD1	1:A:490:ASN:C	2.48	0.47
1:A:492:ILE:O	1:A:493:GLU:C	2.48	0.47
1:A:493:GLU:HA	1:A:497:SER:HB2	1.97	0.47
1:A:474:GLU:HG3	3:A:924:HOH:O	2.16	0.46
1:A:238:ARG:O	1:A:239:ASN:HB2	2.17	0.45
1:A:441:LEU:HA	1:A:444:ILE:HD12	2.00	0.44
1:A:426:MET:HE3	1:A:444:ILE:HG12	2.00	0.44
1:A:300:LYS:NZ	3:A:520:HOH:O	2.51	0.43
1:A:457:THR:O	1:A:461:SER:HB2	2.19	0.43
1:A:425:LEU:HG	1:A:440:ILE:HG23	2.03	0.41
1:A:235:ASN:OD1	3:A:501:HOH:O	2.22	0.41
1:A:489:LEU:O	1:A:490:ASN:C	2.48	0.41
1:A:285:ARG:HD2	3:A:624:HOH:O	2.22	0.40
1:A:434:THR:O	1:A:438:GLN:HG3	2.21	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:519:HOH:O	3:A:885:HOH:O[3_544]	1.77	0.43
3:A:632:HOH:O	3:A:699:HOH:O[3_544]	2.09	0.11

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	426/447 (95%)	422 (99%)	4 (1%)	0	100	100
2	B	12/55 (22%)	12 (100%)	0	0	100	100
All	All	438/502 (87%)	434 (99%)	4 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	359/375 (96%)	354 (99%)	5 (1%)	67	53
2	B	15/47 (32%)	14 (93%)	1 (7%)	16	4
All	All	374/422 (89%)	368 (98%)	6 (2%)	62	48

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	348	LYS
1	A	458	GLU
1	A	461	SER
1	A	485	TYR
1	A	489	LEU
2	B	605	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	88	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	426/447 (95%)	0.07	23 (5%) 25 28	13, 23, 61, 79	0
2	B	16/55 (29%)	0.87	2 (12%) 3 4	17, 33, 56, 61	0
All	All	442/502 (88%)	0.10	25 (5%) 23 26	13, 23, 61, 79	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	497	SER	10.6
2	B	600	SER	7.0
1	A	485	TYR	6.8
1	A	486	LYS	6.6
1	A	478	ARG	5.7
1	A	496	PHE	5.1
1	A	432	LYS	4.1
1	A	493	GLU	4.1
1	A	489	LEU	3.6
2	B	639	GLU	3.3
1	A	75	ASN	2.9
1	A	88	ASN	2.8
1	A	455	GLY	2.7
1	A	482	GLU	2.6
1	A	471	ASP	2.4
1	A	452	GLU	2.3
1	A	74	VAL	2.3
1	A	495	TYR	2.3
1	A	474	GLU	2.2
1	A	107	GLU	2.2
1	A	72	GLY	2.1
1	A	453	LYS	2.1
1	A	108	LYS	2.1
1	A	490	ASN	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	480	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.